

Morphology and Influence of Various Plant Growth Substances on Germination and Early Seedling Growth in *Macrotyloma uniflorum* (Lam.)

J.S. Chauhan^{1*}, Y.K. Tomar², Anoop Badoni¹, N. Indrakumar Singh¹, Seema Ali¹, Debarati¹, A.S. Rawat¹ and V.P.Nautiyal¹

1. Department of Seed Science & Technology, H.N.B. Garhwal Central University, Srinagar Garhwal, Uttarakhand-246 174 (India).
2. Department of Horticulture, H.N.B. Garhwal Central University, Srinagar Garhwal, Uttarakhand- 246 174 (India).

*Corresponding author: jaisinghchauahn@gmail.com

Abstract

The paper presents the results of studies on morphological characters, seed germination and the influence of different concentrations of plant growth substances on *Macrotyloma uniflorum* including the comparative growth patterns of the seedlings. This is one of the lesser known beans mainly cultivated in hilly areas and commonly grown up at 1800m above MSL. Seeds pre-soaked for 24h in various concentrations (0.1, 1.0 and 10 ppm) of GA₃, IBA and NAA respectively putting a separate control set soaked only in distilled water. The mean value of germination percentage, growth of root, shoot and cotyledonary expansion and biomass of seedlings were computed. The maximum germination percentage (99%) was observed through GA₃ 0.1ppm and NAA 1 ppm in comparison to control set (90%). Although highest elongation of shoot was observed under GA₃ 10 ppm (11.29 cm) and lowest under GA₃ at 1 ppm (10.45 cm) in comparison to control (7.50 cm) but the highest elongation of root was favoured by GA₃ at 0.1 ppm (4.46 cm) whereas, the minimum was observed under NAA at 10 ppm (1.28 cm). The maximum Cotyledon expansion was gained in GA₃ at 1ppm (5.36 cm) as compared to controlled set (2.94 cm). IBA 0.1 ppm resulted in highest dry weight of shoot (0.0183 g.) as compared to control (0.0172 g.) while dry weight of root was recorded highest under NAA at 10 ppm (0.0076 g). Lower concentrations of all the plant growth substances exhibited supportive response towards germination, growth of seedlings and development of cotyledons. The results of present findings show that applications of these plant growth substances if used at lower concentrations could enhance germinations and healthy seedling growth leading towards high production of *Macrotyloma uniflorum* to fulfill the demand of farming community. It is thus imperative that more identical studies be conducted on other legume species also generating data base so that the use of growth substances can be extended to produces healthy seedlings required to support large scale seed germination and plantations of those species. [Journal of American Science 2009;5(6):43-50]. (ISSN: 1545-1003).

Keyword: *Macrotyloma uniflorum*, Morphology, Plant Growth Substances, Seed Germination, Seedling Growth.

Introduction

Production of high quality seeds is primary source to the success of Indian agriculture. Every farmer is sensitive to need for the rapid uniform seedling emergence and establishment of an even and productive stand. Crop production relies heavily on high quality planting seeds. The latest ISTA rules (ISTA, 2008) contain seed testing protocols of a large number of species cultivated all over the world and it

forms the basic reference book for all kinds of seed testing activities and also for the international seed trade. Seeds are fundamental input to agriculture and natural ecosystem. The production of high percentage of viable seeds with capacity to germinate quickly is necessary for the propagation. Seed quality is very essential for optimum stand establishment and maximum yield potential. As a result, it is necessary

to have different seed testing parameters that permit rapid, objective and accurate evaluation of seed quality (S.C. Joshi *et al* 2009).

Horse gram (*Macrotyloma uniflorum* Lam) is a popular pulse, locally known as Gaheth belongs to the family Fabaceae that still remain an under exploited legume crop. It is usually grown up to the area at 1800msl. Horse gram seeds are rich in protein and consumed in majority by poorest section of the society. The fodder being rich in protein; it is widely used as a feed to animals and horses (B. G. Prakash *et al.* 2008). Dehusking, germination, cooking, and roasting have been shown to produce beneficial effects on nutritional quality of the legumes. Seeds and fruits of different species vary greatly in appearance, shape, size and ornamentation and structure of the embryo in relation to storage tissues. In some legumes seed coat colour is quite heterogeneous within a sample and such seeds often comprise a germplasm accession in a gene bank. Correlation of different seed coat colours with seed quality parameters may provide a non-destructive tool (especially useful in a gene bank) for sorting good quality seeds.

Germination represents a critical event in plant's life cycle and its timing largely predetermined the chances of survival of a seedling up to maturity (Thompson, 1973). Temperature is an important physical parameter of an environment, which

Material and Methods

The seeds of *Macrotyloma Uniflorum* were collected from Jakh village of District Tehri of Uttarakhand, India. The observation recorded on various morphological features of the seed such as shape, size, colour, ornamentation, length and width, fresh and dry weight of seed, floral characters along with length and width of pod, pods per plant and seed per pod etc. After thorough mixing, the whole lot was sampled and dried in open air for ten days and stored

$$\text{Moisture content (\%)} = \frac{\text{Fresh Weight} - \text{Dry weight}}{\text{Fresh weight}} \times 100$$

Seeds were pre-soaked for 24h under the different concentrations of Plant Growth Substances (0.1, 1.0 and 10 ppm) of GA₃, IBA and NAA. A control set was soaked only in distilled water. The seeds were placed on a wet filter paper in petridish and kept in

determines the success or failure of a species in a particular locality, which in turns depends mostly on the germinability of the seed. It is well known that the different population of the same species varies in their temperature and light requirements for germination. Germination requirements of a particular species are a result of the interaction of its genetic makeup with the environment and dormancy pattern of seeds of various plant species, which enable them to survive during adverse conditions (Wittington, 1973; Nikolaeva, 1977). The germination of seed is affected by hormonal secretion and enzymatic activity within the seeds. Gibberellins (GA₃) and Auxin- IBA (Indolebutyric acid), IAA (Indole acetic acid) control many behavioral functions in plants which act as chemical messengers influencing many patterns of plant development. These stimulate cell division and cell elongation and control enzyme secretions. Seed germination and seedling growth can be influenced by various concentrations of growth regulators i.e. GA₃ and IAA (Chauhan *et al.* 2009).

The present investigation is carried out to investigate the response of different concentrations of the Plant Growth Substances on seed germination, root, shoot and hypocotyls elongation and biomass production of seedlings. Some morphological features of seed of Horse gram have also been observed with germination processes.

at room temperature till the experimentation. The moisture content of seeds was determined by air oven method. Subsequently, the sample contained in glass and dried thermostatically at room temperature and weighed, then placed in hot air oven at 80⁰ C for 48hrs to find out the average dry weight and calculated using the following formula (Anonymous, 1976).

Seed Germinator. The mean value of germination percentage, growth of root, shoot and cotyledonary expansion and biomass of seedlings were computed on each alternate day till the final day of experimentation (Table 1).

Table 1. The effects of various concentrations of different plant growth substances on seed germination and seedling growth of *Macrotyloma uniflorum*.

Plant Growth Substances	Concentrations used		
GA ₃	0.1ppm	1ppm	10ppm
IBA	0.1ppm	1ppm	10ppm
NAA	0.1ppm	1ppm	10ppm
Control	Only distilled water		

Results and Discussion

Macrotyloma Uniflorum is an erect, sub-erect or trailing, densely hairy annual herb. The tap root produces a branched root system with smooth, rounded nodules. Seed is small somewhat gray to brown with pale fawn in colour sometimes with faint mottles or with small scattered black spots or with both. Seed size ranges 6-8 mm long and 3-4mm broad smooth of which 100 seed weight is recorded 3.65gm. The flowers are short and only 6-12 mm long cream - yellow with purple spot in auxiliary racemes with 2 appendages at base. The pods are 4-6 cm long and about 6 mm broad with 18-23 pods per plant bearing 6-7seeds per pod. Flowering and fruiting takes place between August to October (Table-2).

Correlation of different seed coat colours with seed quality parameters may provide a non-destructive tool (especially useful in a gene bank) for sorting

good quality seeds. Relationship between seed coat colour and quality parameters in horse gram (*Macrotyloma uniflorum*) was studied by Singh et al. (2009) in seeds separated in to pale brown, medium brown and blackish brown coloured fractions in 20 accessions of horse gram. Seed quality was assessed in terms of germination, seedling vigour and seed storability. The light coloured fractions consistently showed highest germination percentage and seedling vigor followed by medium and dark coloured fractions in all the accessions. Percentage seed moisture content and electrical conductivity were observed to be highest in the dark coloured seeds. Light coloured seeds showed better storability after three years of ambient storage whereas the medium and dark coloured seeds were poor in germination and showed significant decline. Colour of seeds can be used as a visual indicator of seed quality and storability in horse gram.

Table 2. The Morphological Features of *Macrotyloma uniflorum* Seeds

S. No.	Parameters studied	Range of Variation
1.	Shape of seed	Seeds are small and somewhat round
2.	Colour	Gray to brown with pale fawn in colour. Sometimes with faint mottles or with small scattered black spots or with both.
3.	Ornamentation	Smooth and ovoid
4.	Length of seed	6-8 mm
5.	Width of seed	3-4 mm broad
6	Seed fresh weight (100)	3.65 gm
	Seed dried weight	1.095 gm
7	Flowers	Short only 6-12 mm long. The flower is cream - yellow with purple spot in auxiliary racemes with 2 appendages at base.
8	Pod	Shortly stipitate, slightly curved and tomentose.
9	Length and width of pod	4-6 cm long and about 6 mm broad with a point about 6 mm long.
10	Pods per plant	18-23 pods
11	Seed per pod	Usually bear 6 or 7 seed per pod.
12	The sowing time of the seed	Last week of June to first week of July.
	Flowering and fruiting period	The plant attains flowers and fruits between Aug to Oct.

13	Harvesting	The crop is harvested, when the seeds turn to shining brown-grey color, generally during mid Oct to Nov.
----	------------	--

The observations of the experiment showed that GA₃ 0.1ppm and NAA 1ppm resulted in equal 99 percent germination which was followed by IBA10ppm (96%) while control set showed lowest germination (90%). GA₃ 1, GA₃10, IBA 0.1, and IBA-1ppm resulted in equal percent germination (98). Within the concentrations, the NAA 10ppm responded only 96 percent germination which is less than the control. (Table 3 and Fig. 1).

In case of the seedling growth, GA₃ at 10ppm attained maximum (11.29cm) shoot length which was followed by GA₃ at 1ppm (10.45cm) and IBA at 0.1ppm (9.23cm). Minimum shoot length was observed under NAA 10ppm (7.02cm) that was slightly lower than the control set (7.50cm). As far as root growth is concerned, it was observed highest under GA₃ at 0.1ppm (4.46) followed by NAA at 1ppm (4.07). The lowest performance with regards to the root length was recorded under NAA at 10ppm (1.28) that was very less than control (3.68).The

cotyledon expansion was observed highest under the influence of GA₃ 1ppm (5.36 cm) that was followed by GA₃ 10ppm(3.79, Fig-4). The control set exhibited (2.94 cm) a little higher than NAA-10 ppm (2.27).

The highest cotyledons development has enhanced under the GA₃ at 1ppm (5.36 ±0.5cm) concentration that was followed by GA₃ at 100ppm (5.36 ±0.5cm) and GA₃ 0.1 ppm(3.65 ±0.3). The lowest growth was recorded under the influence of NAA 10ppm (2.27 ±0.2) which is less than the control set (2.94 ±0.2, Fig-2). Dry weight of root biomass in case of controlled set was observed to be 0.0047±0.004g. The maximum dry weight of root was found in NAA at 10 PPM (0.0076 ±0.006g) and minimum dry weight of root was observed under GA₃ at 10 PPM (0.0021±0.002). The maximum fresh weight of shoot (0.0183 ±0.5) was observed under IBA at 0.1 ppm whereas, the dry weight of shoot under the control set has recorded higher (0.0172±0.2) than some of other concentrations (Table-4 and Fig.3).

Table 3: Effect of Various Concentrations of Plant Growth Substances on the Seed Germination and Seedling Development of *Macrotyloma uniflorum*.

Treatments	Germination %	Shoot Length (cm)	Root Length (cm)	Cotyledon Expansion (cm)
GA ₃ 0.1 ppm.	99	9.53 ±0.7	4.46 ±0.3	3.65 ±0.3
GA ₃ 1 ppm.	98	10.45 ±0.5	3.57 ±0.2	5.36 ±0.5
GA ₃ 10ppm.	98	11.29 ±0.8	3.69 ±0.5	3.79 ±0.2
IBA-0.1 ppm.	98	9.23 ±0.7	3.86 ±0.3	3.05 ±0.3
IBA -1 ppm.	98	8.34 ±0.7	3.95 ±0.3	2.91 ±0.2
IBA -10 ppm.	96	9.50 ±0.7	3.55 ±0.3	2.94 ±0.2
NAA- 0.1 ppm.	97	8.54 ±0.8	1.93 ±0.5	3.12 ±0.5
NAA -1 ppm	99	8.84 ±0.8	4.07 ±0.5	2.8 ±0.2
NAA -10ppm.	96	7.02 ±0.8	1.28 ±0.5	2.27 ±0.2
CONTROL	95	7.50 ±0.8	3.68 ±0.5	2.94 ±0.2



Table 4: Effect of Various Concentrations of Plant Growth Substances on the Shoot and Root Dry Weight of *Macrotyloma uniflorum*.

Treatments	Shoot Dry Weight(gm)	Root Dry Weight(gm)	Shoot: root ratio	Epicotyls length((cm)	No. of Roots
GA ₃ 0.1 ppm	0.0143 ±0.5	0.0053 ±0.005	1: 13	2.92	7
GA ₃ 1 ppm	0.0174 ±0.2	0.0029 ±0.002	1:0.8	2.62	9
GA ₃ 10ppm	0.0162 ±0.2	0.0021 ±0.002	1:0.2	3.52	9
IBA- 0.1 ppm	0.0183 ±0.5	0.0043 ±0.003	1: 4.1	3.46	8
IBA -1 ppm	0.0152 ±0.5	0.0036 ±0.003	1:3.6	3.44	8
IBA -10 ppm	0.0137 ±0.2	0.0053 ±0.005	1:3.5	1.87	8
NAA -0.1 ppm	0.0140±0.2	0.0045 ±0.004	1:0.6	2.45	7
NAA- 1 ppm	0.0143 ±0.2	0.0055 ±0. 005	1:40	2.58	7

NAA- 10 ppm	0.0175 ±0.5	0.0076 ±0. 006	1:1.8	0.29	6
Control	0.0172 ±0.2	0.0047 ±0. 004	1: 6.7	2.85	7

Heavy seeds generally have shown superior germination, survivorship, and seedling mass as also have been mentioned by Aaron M. Ellison (2001). The findings of Douglass H (1985) have summarized that seed size and germination requirements can be

determined to be useful characters for resolving systematic and phylogenetic problems. Archana and Shivana (1985) have studied the requirements for seed germination and seedling formation of a hemi-root parasite *Sopubia delphinifolia*.





Light was found to be essential for germination; none of the growth substances could

replace the light requirement. Seed must attain certain minimum specific moisture content, before they

germinate (Negbi *et al.*, 1966). GA₃ at 0.1ppm affected germination the most, leading 4% higher than the control. The more effective concentration has been proved to be GA₃ at 10 ppm with regards to shoot growth (11.29 ±0.8) at the cost of only 1% less germination. The same concentration resulted in almost equal growth of root (3.69 ±0.5) in comparison to control set. Response of lower and middle level of concentration such as GA₃ at 0.1ppm

and NAA at 1ppm has shown remarkable germination and seedling growth in *Macrotyloma*. GA₃ resulted in maximum shoot growth (11.29±0.8) in comparison to Control (7.50±0.8). These concentrations have shown good response on cotyledons development (5.36±0.5). As far as dry weight of shoot and root is concerned, GA₃ at10 ppm resulted less responsive in comparison to IBA 0.1 (0.0183±0.5) and GA₃ 1 ppm (0.0174±0.2).



The effect of seed-soaking for 24 hours with different Plant Growth Substances has been examined by M. Grzesik. (2006) on the growth of seedlings of Lathyrus

odoratus, Zinnia elegans, Matthiola incana and Antirrhinum majus. GA₃ improved the germination of the treated seeds. The seedlings of Zinnia, Matthiola

and *Antirrhinum* treated with NAA, GA₃ and GA₄₊₇ where higher, better branched and of better quality than the non-treated plants. Singh and Dara (1971) have shown the results on Influence of presoaking of seeds with gibberellins and Auxin on growth and yield attributes of wheat under high salinity, sodium adsorption ratio and boron levels. Similar efforts have been made through another experiment by M. Farooq et al. (2006) to investigate the possibility of rice seed invigoration by pre-sowing ethanol seed treatment. They revealed that employing ethanol treatments at lower concentrations can invigorate fine rice seeds. High temperature both delayed and inhibited the germinations of barley and radish seeds (Cavusoglu K & Kabar K 2007). Irfan Afzal et al., (2005) have shown the effects of seed soaking with plant growth regulators (IAA, GA₃, and Kinetin) on wheat emergence and seedling growth under normal and saline conditions and found their usefulness in increasing relative salt-tolerance. Among the 14 pre-sowing treatments, KNO₃ (150 min) and NaHClO₃ (30 min) significantly stimulated seed germination of *Angelica glauca* and reduced mean germination time under both laboratory and nursery trials, as well as developed seedling vigour under nursery conditions (Jitendra S. Butola and Hemant K. Badola 2004). A similar experiment was carried out by Gao Huan Zhang et al. (2002) in which Walnuts cv. Jianshi were soaked in water (control), IBA at 80 mg/kg, IAA at 100 mg/kg, NAA at 80 mg/kg, ABT root-growing powder at 1 g/kg, or 6-BA at 5 mg/kg. In this experiment, different concentrations of GA₃, IBA and NAA, the maximum germination percentage (100%) was recorded in seed treated with IBA 0.1 ppm, which shows that the lower concentrations of growth hormone shows better performance than higher which was similar to the results of James Chukwuma Ogbonna and P.G. Abraham (1989).

The results of the present findings are parallel to the findings of Parameswari and Srimathi (2008) to evaluate the influence of plant growth substances on seed germination and seedling quality characteristics of tamarind seeds. They revealed that the concentration of 100 and 200 ppm GA₃ were found to enhance both the seed germination and seedling vigour parameters, such as root length, shoot length, hypocotyl length, dry matter production and vigour index values. Seed fortification with GA₃ 100 ppm performed well even under nursery conditions recording a higher leaf number and stem circumference.

IBA proved good in seedlings biomass production. However, lower concentrations of all the plant growth substances exhibited supportive response towards

germination, growth of seedlings and development of cotyledons. The results of present findings shows that applications of these plant growth substances if used at lower concentrations could enhance germinations and healthy seedling growth leading towards the production of *Macrotyloma uniflorum* to fulfill the demand of farming community. It is thus imperative that more identical studies be conducted on other legume species also generating data base so that the use of growth substances can be extended to produces healthy seedlings required to support large scale seed germination and plantations of those species.

Corresponding Author:

Dr. J.S. Chauhan, Assoc. Professor,
Department of Seed Science & Technology
H.N.B. Garhwal Central University,
Srinagar Garhwal, Uttarakhand-246 174(India).
Phone: +911370267664
Email: jaisinghchauahn@gmail.com

References

- [1] Aaron M. Ellison. Interspecific and intraspecific variation in seed size and germination requirements of Sarracenia (Sarraceniaceae). American Journal of Botany. 2001; 88:429-437.
- [2] Archana Sahai and KR Shivanna. Seed Germination and Seedling Growth in *Sopubia delphinifolia*- a Hemi-root Parasite: Germination Requirements. Annals of Botany 1985; 55: 775-783.
- [3] BG Prakash, MB Guled and Asha M Bhosale. Identification of Suitable Horse gram Varieties for Northern Dry Zone of Karnataka. Karnataka J. Agric. Sci. 2008; 21(3): 343-345.
- [4] Cavusoglu K, Kabar K. Comparative effects of some plant growth regulators on the germination of barley and radish seeds under high temperature stress. Eurasia J BioSci. 2007; 1, 1, 1-10.
- [5] Chauhan JS, YK Tomar, N Indrakumar Singh, Seema Ali, Debarati. Effect of Growth Hormones on Seed Germination and Seedling Growth of Black Gram and Horse Gram. Marsland Press. Journal of American Science 2009;5(5):79-84. <http://www.americanscience.org>
- [6] Douglass H. Morse and Johanna Schmitt. Propagule size, dispersal ability, and seedling performance in Asclepiasiaceae. Oecologia. 1985; Volume 67, Number 3.
- [7] Gao HuanZhang, Bao XinMei, Hu XiaoXiang. Effect of plant growth regulators on seed germination

and seedling growth of walnut. Journal of Hubei Agricultural College. 2001; Jingzhou 434025, Hubei, China

[8] Irfan Afzal, Shahzad MA, Basra and Amir Iqbal. The effects of seed soaking with plant growth regulators on seedling vigor of wheat under salinity stress. Journal of Stress Physiology & Biochemistry, 2005; Vol. 1, No. 1, pp. 6-14.

[9] ISTA. International Rules for Seed Testing. ISTA, 2008; Bassersdorf, CH. Switzerland.

[10] James Chukwuma Ogonna and PG Abraham. Effect of Seed Pretreatment with some Plant growth regulator on germination, growth and yield of Cowpea (*Vigna sinensis*). Japan Jour. Crop. Sci. 1989; 58(4): 641-647.

[11] Jitendra S Butola and Hemant K Badola. Effect of pre-sowing treatment on seed germination and seedling vigour in *Angelica glauca*, a threatened medicinal herb. Current Science, 2004; vol. 87, no. 6, 25.

[12] M Farooq, Shahzad MA Basra, Hafeez-Ue-Rehman and Tariq Mehmood. Germination and early seedling growth as affected by pre-sowing ethanol seed treatments in fine rice. International Journal of Agriculture & Biology. 2006; 08 (1)19-22.

[13] M. Grzesik. Effect of Growth Regulators on the Seedling-Growth of *Lathyrus Odoratus*, *Zinnia Elegans*, *Matthiola Incana* and *Antirrhinum Majus*. Acta Horticulturae 2006; 251: III International Symposium on Growth Regulators in Ornamental Horticulture.

[14] Negbi M, Rushkin E and Koller D. Dynamics aspects of water relations germination of *Hirschfeldia incana* seeds plant cell Physiology. 1966; vol.7pp, 363-376.

[15] Nikolaeva MG. Factors controlling the seed dormancy pattern. In: The Physiology and Biochemistry of Seed Dormancy and germination. 1977; (Ed. AA Khan) pp5174. Elsevier.

[16] Parameswari K, Srimathi P. Influence of growth regulators on elite seedling production in tamarind (*Tamarindus indica*) Legume Research. 2008; Volume: 31, Issue: 4 Print ISSN: 0250-5371.

[17] Singh H and Dara BL. Influence of presoaking of seeds with gibberellins and Auxin on growth and yield attributes of wheat (*Triticum aestivum* L.) under high salinity, sodium adsorption ratio and boron levels. Indian J. Agr. Sci., 1971; 41: 998-1003.

[18] Singh N, Devi C, Kak A, Singh G, Kumari A, Mahajan RK. Influence of seed coat colour associated heterogeneity on quality and storability in horse gram (*Macrotyloma uniflorum*) Seed Science and Technology. 2009; Volume 37, Number 1, pp. 232-240(9)

[19] Sunil Chandra Joshi, N Indrakumar Singh and JS Chauhan. Soyabean Seed Quality Evaluation. Nature and Science, 2009;7(5), ISSN 1545-0740. <http://www.sciencepub.net>, naturesciencej@gmail.com

[20] Thompson PA. Effect of cultivation on the germination character of the Com cockle (*Agrostemma githago* L) Ann. Bot. 1973;36:133-154.

[21] Wittington WJ. Genetic regulation of germination. In: Seed Ecology, (ed. W. Heydecker) Proc. Nineteenth Ester School in Agric. Sci. 1973; Butterworth, Lond. Pp5-30.

12-08-2009