

## Hydro-Chemistry, Macro-Invertebrate Fauna And Fish Production Of Acadja Fishing Sites In A Tropical Lagoonal Ecosystem

Emmanuel, Babatunde Eniola<sup>1\*</sup>, Chukwu, Lucian Obinna<sup>1</sup> and Bakare, Sheriff Olaide<sup>1</sup>  
1 Department of Marine Sciences, University of Lagos, Akoka – Yaba, Lagos Nigeria.

\*[monetemi@yahoo.com](mailto:monetemi@yahoo.com)

**ABSTRACT:** An investigation of the hydrochemistry, macro – invertebrates fauna and fish production of three selected acadja fishing sites was carried out for six months (April – September, 2004) in the Lagos lagoon. Depth, Total suspended solid, total dissolved solid, nitrates, phosphates, sulphates, dissolved oxygen, biochemical oxygen demand and chemical oxygen demand recorded high or increasing values during the rainy season while transparency and salinity showed progressively lower values. The biological oxygen demand (BOD) and chemical oxygen demand (COD) ranged from 70.0 to 115.0mg/l and 20.0 to 75.0mg/l respectively. A total of twenty species and eighteen genera of Macro - invertebrate were recorded. The mollusks were the most abundant and diverse phyla, followed by the crustacean and annelids. The dominant species throughout the study were *Tympanoyonus fuscatus var radula*, *Balanus pallidus*, *Aloidis trigona* and *Mercierella enigmatica*. A total of 89.80kg (38.52%) fish was harvested in Site A, 71.30kg (30.59%) in Site B and 72.0 kg (30.89%) in Site C. *Sarotherodon melanotheron* and *Tilapia guineensis* contributed 36.75% of the total weight in Site A, 34.64% in Site B and 38.75% in Site C. Twelve fish species occurred in the Site A and C while eleven fish species occurred in fishing Site B with least species diversity. The profitability indices show that ‘acadja’ brush park fishing practice is a profitable business in the lagoon. [Journal of American Science 2009;5(7):42-48]. (ISSN: 1545-1003).

**Keywords:** Hydro – chemistry, macro – invertebrate, fish acadja, lagoonal ecosystem.

### INTRODUCTION

Ecological studies of benthos in the Lagos lagoon were initiated several decades ago by Webb and Hill (1958), also Sandison and Hill (1966) and Oyekan (1988) have also been major contributors. More recently are the investigations by Brown (1998) and Brown and Oyekan (1998). Oyekan (1988) put forward a classification of the benthic communities of the Lagos lagoon. The classification was based on the nature of deposit in which they occur and on the characteristics faunal element. Five communities were identified in Lagos lagoon: mangrove, pachymelina, estuarine, amphipod, venus and estuarine rock communities (Oyekan, 1988). The fish and the fisheries of the Lagos lagoon has been documented by authors like Fagade (1969); FAO (1969), Fagade and Olaniyan (1974), Udolisa and Solarin (1979), Solarin (1998), Solarin and Kusemiju (2003a&b), Emmanuel *et al.*, (2008a &b).

Over the years the Lagos lagoon has continued to be under intensifying pressure from pollution. According to Ajao *et al.*, (1996) coastal waters can be contaminated from both natural and anthropogenic sources of pollution. The anthropogenic sources include sand mining, industrial and domestic effluents, logging and timber transportation by water. The anthropogenic pollutants/waste has an attendant effect on the biodiversity of aquatic resources. Reduction in the composition and density of algae, zooplankton, benthic invertebrates, fish and fisheries resources has

been linked to the increasing menace over the years of aquatic pollution in the Lagos lagoon, Nigeria (Akpata *et al.*, 1993; Nwankwo, 1998; Solarin 1998; Emmanuel *et al.*, 2008a&b).

Lagos lagoon has supported decades of small-scale fisheries which have shown signs of over-exploitation. The fishing gears and crafts used include gillnet, stownets traps, liftnets, longline, basket traps etc (Fagade, 1969; Solarin, 1998; Emmanuel 2008; Emmanuel *et al.*, 2008a&b).

The term ‘acadja’ describes a family of installation of the fish-park type that is currently found in several of the West African coastal lagoon (Welcomme, 1972). Welcomme and Batley (1998) listed fish shelter as one of the techniques for stock enhancement. Acadja fish fence provide shade, hiding place, shelter as well as food for fishes. The acadjas also act as breeding, spawning or nursery areas for fishes and also act as point of attachments for algae and macrobenthic invertebrate. The sticks used for the acadja are progressively colonized by bacteria, fungi algae, nematodes, annelids, mollusk and other invertebrates forms which all serve as food for a wide away of fishes. Hence, the mechanism of fish aggregation is prompted by the shelter, security and colonization of the fish fences by plankton and invertebrates organism which in turn prompts the growth of a large number of small planktophagus and benthophagus fishes which feed on them. It takes about three or more months for the sticks to deteriorate



The influx of seawater from the sea give rise to brackish conditions in the lagoon and encourages the growth of mangrove which lines the edges of the many waters ways and cover those areas of low lying swampy grounds which are inundated at the high (Spring) tides.

### Sampling Stations

For this study, three strategic sites have been chosen for ecological assessment. Station A is a pollution relatively free site within the lagoon whereas Station B is a site where kitchen waste and organic materials from University of Lagos staff club and guest house are channeled into the lagoon. Household or domestic waste and sewage are prevalent in Station C, emanating from household at the edges and within the lagoon.

Collection of water sample and sediment, temperature were measured in-situ on the field with mercury in glass thermometer. Water samples were collected at each site in a 250ml water container between April and September 2002. Van-Veen grab (0.1m<sup>2</sup>) was used to collect the sediment. The water samples were then transported and stored in a refrigerator for further analysis. The sediment were emptied into a black polythene bag and stored in a deep freezer (<0°C).

### Physico-Chemical Parameters

Surface water samples were collected about 10cm the water surface. On each trip, temperature (°C) was recorded with mercury-in-glass thermometer and transparency (cm) measured by a 20cm diameter Secchi disc. Salinity (‰) was determined using the silver-nitrate-chromate method as suggested by Barnes (1980) and Nwankwo (1991) and pH measured in the field a Hackkit (Del, 2001). Dissolved oxygen (mg l<sup>-1</sup>) was measured by the Grifin oxygen meter (Model 40). Total suspended solids (TSS), (mg l<sup>-1</sup>) was estimated by the evaporation of a 100ml aliquot.

### Sediment Analysis

Sediment particle size analysis was done using the method described by Brown (1991) and total organic matter determined by the rapid ash method employed by Oyeneke (1975).

### Macrobenthic Invertebrate and Fish Collection

One of the grab was washed through a 0.5mm sieve in the field. The retained materials were preserved in 4% formalin and kept in labeled jars for further sorting and analysis in the laboratory. Fishes in the acadja sites were collected with enacting net (10mm-mesh polyamide) which was used to form a

complete fence. The debris within was removed and gradually the net was drawn together to entrap the fishes.

### (1) Statistical Analysis

All statistical methods used were adapted from Ogbeibu (2005). Macrobenthic community structure was estimated using the Margalef's species richness index (d) (Margalef, 1951); Shannon-Weiner information function (AS) (Shannon and Weaver, 1949) and Equitability index or Evenness (Lloyd and Ghellardi, 1964).

### (2) Economic and Profitability Analysis

The economic and profitability analyses as described by Ayinde and Aromolaran (1998) and Abdul *et al.*, (2004) were adopted. Net income of fisherman practicing "acadja" fishing was calculated for the six month. This analysis was used in the profitability analysis to calculate return to management, capital and family labour, return to investment, return on fixed cost or gross margin, rate of return on fixed cost, rate of return on variable cost and rate of return on labour cost.

Return to management, capital and family labour (RMCF):

RMCF = TVP – TFC – TVC less cost of capital borrowed and cost of family labour

Rate of Return to Investment (RRI):  $RRI = (RMCF/TC) \times 100$

Return on Fixed Cost (RFC) or Gross Margin (GM):

$RFC = TVP - TVC$

Rate of Return on Fixed Cost (RRFC):  $RRFC = (RFC/TFC) \times 100$

Rate of Return on Variable Cost (RRVC):  $RRVC = (RTVP - TFC/TVC) \times 100$

Rate of Return to Labour (RRLC):

$RRLC = TVP - \text{All costs except labour cost} / \text{total labour cost (man/day)} \times 100$

Where, TVP = Total value of product

TFC = Total fixed cost

TVC = Total variable cost

TC = Total cost

## RESULTS

The monthly variation in the physico-chemical parameter of water at the acadja fishing sites in the lagoon is shown in **Table 1 a&b**. The highest surface water temperature during the study was 32°C in May and the lowest was 28°C in July. pH values were essentially alkaline throughout the study period. pH range of 7.2 – 8.0 was recorded for water while that of sediment ranged between 7.0 and 8.0. A

neutral pH of 7.0 was recorded in Station A and B in July for sediment and water respectively. Total dissolved solid ranged between 10.0 and 24.0mg/l and the highest (24.0mg/l) was recorded in June while the lowest (10.0mg/l) was recorded in April, August and September. Total suspended solids in study areas ranged between 10.0 and 21.5mg/l. Turbidity was generally high for the months of May, June, July, August and September with highest value of 22.5mg/l in Station A. Salinity value was highest in April (20.5‰) while the lowest (2.0‰) was recorded in September. Chloride content of the study areas was fairly constant with a range of 10.0 – 18.0mg/l. The highest chloride value was observed in June and July. Low value of oil and grease was obtained in this study with maximum value of 0.25mg/l in July. The highest nitrate value (2.5mg/l) was recorded in April. Phosphate value was high in June with 4.5mg/l and the lowest (2.2mg/l) in July. Dissolved oxygen was high during the early part of the rainy season (April – June) with 6.5mg/l value at Station A and B in June and Station B in May and the lowest (3.0mg/l) in August (Station A & B) and September (Station C). A trend of high chemical oxygen demand was recorded in April, May, June and July with a peak in June (75mg/l). A range of 70 – 115.0mg/l was recorded for BOD in the study area with the maximum value noted at Station B in May and the lowest in Stations A and B in August and Station B in September. The sulphate level was very low with maximum value of 1.5mg/l. Total organic content (TOC) did not show any marked variation both in the station and the study period.

### Macrobenthic Invertebrates

A total of twenty (20) species of benthic were recorded in this study. The species constituted three phyla: Mollusca, Annelids and Crustacea (Table 2). The mollusca constituted fifteen species, the annelida crustacean a constituted two and three species respectively. The gastropod molluscs include *Semifusus morio*, *Pachymelania aurita*, *Thais haemistoma*, *Natica flammulata*, *Neritina glabrata*, *Nerita* sp, *Tympanotonus fuscatus*, var *radula*, *Terebra micans* and *Turritella unguilina*, while the bivalvea molluscs include *Aloidis trigona*, *Dreissena africana*, *Mactra* sp and *Cryphaea gasar*. The annelids were *Nereis succinea* and *Mercierella enigmatica*. The crustacean were represented by the common lagoon crab – *Callinectes amnicola* and the hermit crab – *Clibenarius africana* and *Balanus pallidus*, *Tympanotonus fuscatus* var *radula* was the most abundant organism during the study, followed by *T. fuscatus*, *Mercierella enigmatica* and *Balanus pallidus*. No specimen was recorded for Station A in

April. Figs 2 to 4 show the number of individual organisms, percentage abundance of classes and monthly variation in the abundance of major taxonomic group of macro – invertebrate in the acadja sites respectively.

Three attached form were recorded in this study these are *Balanus pallidus*, *Mercierella enigmatica* and *Cryphaea gasar*, they were found attached to shells of *Tympanotonus fuscatus* var *radula*. The crustacean and annelids forms (*Balanus pallidus* and *Mercierella enigmatica*) were the more frequent forms on *T. fuscatus* var *radula* shells. Lesser density of these organisms was also found to attach to shells of *Gryphaea gasar*. *Pallidus* was also found attached to stones located on the bottom of the study area. Juveniles of the *T. fuscatus* var *radula* were encountered from June, July, August and September.

### Fish species composition and production in the acadja sites

The fish species composition of the acadja fishing sites is shown in Table 3. A total of 89.80kg (38.52%) fish was harvested in Site A, 71.30kg (30.59%) in Site B and 72.0kg (30.89%) in Site C. *Sarotherodon melanotheron* and *Tilapia guineensis* contributed 36.75% of the total weight in Site A, 34.64% in Site B and 38.75% in Site C. Twelve fish species occurred in the Site A and C while eleven fish species occurred in fishing Site B with least species diversity. The production of the acadja fishing in Lagos lagoon was estimated in Table 4. The production in Site A was ₦19, 463 (US\$ 161.92), ₦15, 877 (US\$132.07).

In this study the abundance of benthos was related to fish production in the ‘acadja’ fishing sites. Table 5 shows that, at the end of six month of operation, a total of N23,000 was invested on three ‘acadja’ brush parks out of which N17,000 and N6,000 were variable and fixed costs respectively. Individual fisherman spent N3, 000 on seine-net rentage, N9, 000 on labour, N2, 000 on brush park maintenance and N3, 000 on canoe rentage. On fixed he spent N6, 000 on ‘acadja’ installation. An economic analysis indicated a fisherman net income of N28, 016. Profitability analysis shows that return to management, capital and family labour (RMCF) and Return on fixed cost (RFC) were N28, 016 and N34, 016 respectively. The rate of returns on investment (RRI), Fixed cost (RRFC), Variable cost (RRVC) and labour cost (RRLC) were 121.8%, 566.9%, 264.8% and 411.3% respectively (Table 6).

The performance of acadja fishery is illustrated in Fig. 5.

Table 1a: Monthly variations in the Physico-chemistry of water samples at the three acadja fishing sites in the Lagos lagoon

Parameters	April			May			June			July			August			September		
	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C
Water temperature	31	31	32	31	32	32	30	30	31	28	28	28	29	28	29	29	30	30
Water pH	7.5	7.2	7.5	7.5	7.2	7.2	7.5	7.5	7.2	8.0	7.0	7.5	8.2	7.4	7.6	8.2	7.5	7.5
TSS	12.0	10.5	12.5	12.5	10.5	10.5	20.0	20.0	21.5	8.5	10.5	10.5	10.0	10.0	12.0	10.0	15.0	15.0
TDS	15.0	12.0	10.0	14.0	15.0	15.5	24.0	22.0	22.5	12.0	11.0	10.5	10.0	12.0	15.0	10.0	15.0	15.5
Turbidity	10.0	9.5	8.2	12.5	10.5	9.5	22.5	21.5	20.0	11.5	10.5	10.0	10.0	10.0	12.5	10.0	15.0	18.5
Salinity	20.5	20.5	18.5	15.0	14.5	14.0	4.0	4.5	4.0	10.0	6.5	5.0	5.0	4.0	3.5	3.0	3.5	3.0
Chloride	10.0	10.0	12.0	-	10.0	12.0	15.0	18.0	0.0	18.0	12.0	15.0	15.0	15.0	12.0	10.0	12.5	12.0
Oil & grease	0.10	0.15	0.09	0.10	0.09	0.12	0.08	0.06	0.10	0.018	0.25	0.20	ND	ND	ND	ND	0.09	0.05
Nitrate	4.5	4.2	3.5	3.5	3.5	3.8	2.5	3.0	4.5	3.0	3.2	2.5	2.5	3.0	3.5	3.0	2.5	3.0
Phosphate	2.5	2.0	2.4	2.8	2.5	3.2	4.5	4.5	3.0	2.5	2.0	2.2	4.0	4.5	3.5	2.5	2.5	3.5
Dissolved O <sub>2</sub>	4.8	5.0	5.5	6.0	6.5	5.0	6.0	6.5	6.5	5.0	4.0	4.5	3.0	3.0	3.5	3.5	4.0	3.0
Chemical O <sub>2</sub> demand	60.0	55.0	65.0	70	65	75	75	70.0	75	65	70	72	35	20	25	40	35	30
Biochemical demand	100.0	95.0	90.0	110.0	115.0	112.5	100.5	95.5	85.0	95.5	85.0	120.0	70	75	95.5	80	75	100
Sulphate	1.5	0.06	0.60	0.15	0.10	0.09	0.10	0.12	0.18	0.15	0.12	0.10	0.09	0.15	ND	0.07	0.12	0.12
Depth	0.95	1.45	1.30	1.51	1.29	1.81	1.71	1.37	1.43	1.24	1.43	1.52	1.86	1.67	1.48	3.8	1.7	2.4
Transparency	0.85	0.68	0.73	0.91	0.65	0.6	0.73	0.56	0.52	0.26	0.33	0.35	0.36	0.35	0.3	0.4	0.35	0.4
Rainfall	159			221			372			296			85			189		

\*ND = Not detected

Table 1b: Monthly variations in the Physico-chemistry of sediment samples at the three Acadja fishing sites in the Lagos lagoon

Parameters	April			May			June			July			August			September		
	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C
Soil pH	7.4	7.5	7.4	7.6	7.5	7.2	8.0	7.6	7.4	7.0	7.2	7.4	7.5	7.4	7.2	8.0	7.5	7.5
Total Organic Matter	19.93	71.72	12.72	20.52	21.15	17.93	19.45	18.62	14.43	18.85	19.73	17.75	15.57	12.23	14.45	17.75	18.83	20.15



Table 2: Composition and abundance of macrobenthic invertebrates at three Acadja fishing sites in Lagos lagoon, Nigeria (April – September, 2002)

Taxa	April			May			June			July			August			September		
	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C	A	B	C
<b>Phylum - Mollusca</b>																		
<b>Class I – Galeodidae</b>																		
Family I – Galeodidae																		
* <i>Semifusus morio</i>								1										
Family II – Melaniidae																		
* <i>Pachymelania aurita</i>				6		8	3					10	3					
Family III – Muricidae																		
* <i>Thais haemostoma</i>											1							
Family IV – Naticacea																		
* <i>Natica flammulata</i>				1			1	1					1					
* <i>N. flannel</i>				1			1											
Family V – Neritidae																		
* <i>Neritina glabrata</i>			2				3						7	1				2
* <i>Nerita</i> sp							1						3	1				
Family VI – Potamididae																		
* <i>Tympanotonus fuscatus</i>						29		1		1		51	1					
* <i>T. fuscatus var radula</i>		31	60	66	19	91	61	26	75	41	24	56	4		14	14	38	35
Family VII – Terebridae																		
* <i>Terebra micans</i>																		
Family VIII – Turritellidae																		
* <i>Turritella unguina</i>				1	4	3			2	2				1				
<b>Class II – Bivalvia</b>																		
Family I – Aloididae																		
* <i>Aloidis trigona</i>			3	4	6		3		2	3		4	8	1			3	7
Family II – Dreissenacea																		
* <i>Dreissena africana</i>				3	1				1			7						
Family III – Mactridae																		
* <i>Mactra glabrata</i>				5				3	3									
Family IV – Ostreaceae																		
* <i>Gryphaea gasar</i>				10	1	4	10		1									
<b>Phylum – Annelida</b>																		
<b>Class – Polychaeta</b>																		
Family I – Nereidae																		
* <i>Nereis succinea</i>			4															
Family II – Serpulidae																		
* <i>Mecierella enigmatica</i>		2	4		9		21	31	58	2		53		13				3
<b>Phylum – Crustacea</b>																		
<b>Class I – Malacostraca</b>																		
* <i>Callinectes amnicola</i>				2														
* <i>Clibinarius africana</i>																		1
<b>Class II – Cirripedia</b>																		
* <i>Balanus pallidus</i>		3	5			385			45	6		34	5	21	5	12	3	34
<b>Total No. of Species</b>	0	3	6	10	6	6	10	5	8	7	1	7	8	6	1	2	4	5
<b>Total No. of Individuals</b>	0	36	78	99	40	520	105	62	187	56	24	215	32	38	14	26	46	80
<b>Ecological Indices</b>																		
SHANNON WEINER'S	0	0.48	0.89	1.28	1.39	0.82	1.34	0.99	1.29	1.01	0	1.66	1.89	1.08	0	0.69	0.65	1.12
MARGALEF'S INDEX	0	0.56	1.15	1.96	1.36	0.79	1.93	0.97	1.34	1.49	0	1.12	2.02	1.37	0	0.31	0.78	0.91




Fig.2: Number of Individual organisms at the Acadja Fishing Site




Fig. 3: Percentage Abundance of classes of macro-invertebrate fauna at Acadja Fishing Site at the Lagos Lagoon (April – September, 2002)




Fig. 4: Monthly variation in the abundance of major taxonomic groups




Fig. 5: Percentage weight of fish caught in acadja fishing sites

Table 3: Fish composition of the ‘Acadja’ fishing sites per six months

Family/Fish Species	Acadja					
	A(kg)		B(kg)		C(kg)	
		%		%		%
<b>Cichlidae</b>						
<i>Sarotherodon melanotheron</i> (Ruppell)	20.5	22.83	15.6	21.88	19.00	26.39
<i>Tilapia guineensis</i> (Bleeker)	12.5	13.92	9.10	12.76	8.90	12.36
<i>Hemichromis fasciatus</i> (Peters)	6.0	6.68	5.0	7.01	5.20	7.22
<b>Bagridae</b>						
<i>Chrysichthys nigrodigitatus</i> (Lacepede)	7.0	7.80	7.0	9.82	7.10	9.86
<i>Liza falcipinnis</i> (Valenciennes)	8.1	9.02	4.8	6.73	5.6	7.77
<b>Cynoglossidae</b>						
<i>Cynoglossus senegalensis</i> (Kaup)	5.6	5.90	5.0	7.01	5.1	7.10
<b>Elopidae</b>						
<i>Elops lacerta</i> (Cuvier & Valenciennes)	6.7	7.46	6.0	8.42	5.9	8.19
<b>Lutjanidae</b>						
<i>Lutjanus goreensis</i> (Valenciennes)	7.6	8.46	6.7	9.40	4.7	6.53
<i>Psettras sebae</i> (Cuvier)	3.0	3.34	1.0	1.40	2.1	2.92
<b>Carangidae</b>						
<i>Caranx hippos</i> (Linnaeus)	4.1	4.57	6.2	8.70	2.2	3.06
<i>Trichiurus lepturus</i> (Linnens)	3.1	3.45	0.0	0.0	1.2	1.67
<b>Portunidae</b>						
<i>Callinectes amnicola</i> (DeRochneburne)	5.6	6.24	4.9	6.88	5.0	6.94
<b>TOTAL</b>	<b>89.8</b>		<b>71.3</b>		<b>72.0</b>	
<b>Percentage total weight</b>	<b>38.52%</b>		<b>30.59%</b>		<b>30.89%</b>	

Table 4: Fish production in ‘Acadja’ Fishing Practice per six months

Fish Species	Station A		Station B		Station C	
	Wt of fish (kg/fishing /ha)	Price (₦)	Wt of fish (kg/fishing /ha)	Price (₦)	Wt of fish (kg/fishing /ha)	Price (₦)
<i>S. melanotheron</i>	20.50	4,305	15.6	3,276	19.0	3,990
<i>T. guineensis</i>	12.50	2,625	9.60	1,911	8.90	1,869
<i>H. fasciatus</i>	6.00	900	5.00	750	5.20	78.0
<i>C. nigrodigitatus</i>	7.00	2,100	7.00	2,100	7.10	2,130
<i>L. falcipinnis</i>	8.1	1,701	4.80	1,008	5.60	1,176
<i>C. senegalensis</i>	5.61	1,400	5.00	1250	5.10	1,275
<i>E. lacerta</i>	6.70	107	6.00	1260	5.90	1,239
<i>L. goreensis</i>	7.60	1,900	6.70	1,675	4.70	1,175

<i>P. sebae</i>	3.00	360	1.00	120	2.10	252
<i>C. hippos</i>	4.10	1,025	6.20	1,550	2.20	550
<i>T. lepturus</i>	3.10	620	0.00	0.00	1.20	240
<i>C. amnicola</i>	5.60	1,120	4.90	980	5.00	1,000
<b>TOTAL</b>	<b>89.80</b>	<b>19,463</b>	<b>71.30</b>	<b>15,817</b>	<b>72.0</b>	<b>15,676</b>

₦120.20 = US\$1

Table 5: Economic income analysis of Acadja brush park fishing practice

ITEMS	VALUE	
<b>A. INCOME</b>		
1. Sales of fish per six month	51,016.00	<b>51,016.00</b>
<b>B. COST</b>		
(i) Variable cost per six month		
(ii) Seine net (55mm) rent 3 times @ N1,000/day	3,000.00	
(iii) Labour (3 'acadja')	2,000.00	
(iv) Maintenance	9,000.00	
(v) Rentage of 2 canoe @ N500/day for 3 'acadja' for 3 days	3,000.00	<b>17,000.00</b>
<b>2. Fixed Costs</b>		
(i) Acadja installation (3)	6,000.00	<b>6,000.00</b>
<b>TOTAL COST</b>		<b>23,000.00</b>
<b>NET INCOME (A – B)</b>		<b>28,016.00</b>

₦120.22 = US\$1

Table 6: Profitability analysis of acadja fishing in Lagos lagoon

PROFITABILITY RATIOS	VALUE
Return to management capital and family labour (RMCF)	28,016
Rate of Return to Investment (%)	121.8
Return on Fixed Cost (RFC)/Gross Margin (GM)	34,016
Rate of Return on Fixed Gross (RRFC) (%)	566.9
Rate of Return on Variable Cost (RRVC) (%)	264.8
Rate of Return on Labour Cost (RRLC)	411.3

## DISCUSSION

The results of the physico-chemical and biological data in this study showed similarly with results from previous ecological investigations (Webb, 1958; Oyekan, 1988; Nwankwo *et al.*, 1994; Brown, 1998 and Emmanuel & Onyema, 2007). Depth, total suspended solids, chemical oxygen demand and biological oxygen demand gave high or increasing values during the rainy period (April to September), on the other hand transparency and salinity showed lower(ed) values. Rainfall associated with floodwater conditions have been shown to reduce transparency while increasing turbidity depth and the nutrient levels

(Nitrate, phosphate and sulphate) of the same water body (Nwankwo, 1991; Brown, 1998; and Chukwu & Nwankwo, 2004). This studied revealed the linkage among the biological abiotic factors and rainfall pattern in the lagoon. This correlates with the report of Hill and Webb (1958) that there is a close correlation between the salinity and other factors in the Lagos lagoon and harbour and the rainfall of the regions draining into the lagoons. Brown and Oyekan (1998) also attributed the variability in temporal distribution of physical, chemical and biological parameters in the Lagos lagoon to the two



seasons experienced yearly with alternative high and low salinity value being recorded in the lagoon.

Biological data from this investigation reveals reduction in the diversity of species compared with earlier works (Oyenekan, 1988; Akpata *et al.*, 1993; and Brown & Oyenekan, 1998). Ajao *et al.* (1996) identified sand mining, sand filling, industrial effluent discharge, oil wastes, domestic waste, sewage discharges among others as human related activities capable and presently destroying the sensitive coastal environment of Nigeria species.

The Pachymelania community as described by Oyenekan (1988) in the study of benthic macrofaunal communities in the Lagos lagoon seen to be the likely community encountered in all the three sites studied. According to the report, the community is characterised by sand and muddy sand deposit. Nominant member include *T. fuscatus*, *A. trigona*, *N. glabrata*, *N. succinea*, *P. Aurita* among others. Salinity and sediment type would seem to be the more important factors controlling the community structure and variability (Brown, 1998).

The occurrence of *N. succinea* in Site C in April in this study indicated that the site is polluted. This agreed with Akpata *et al.* (1993) that the presence of *N. succinea* in large numbers in particular site in the Lagos lagoon reveals that the areas has high organic pollutant load. Observed economically important benthic species utilized at the local level, which are highly valued by man as a source of food were: *T. fuscatus* var *radula*, *T. fuscatus* and *P. aurita*. All benthic macrofauna also serve as source of food for fish. Thus, they occupy a significance position in the aquatic food web (William and Feltmate, 1992; Mackie, 1998; Olomukoro and Ezemonye, 2007).

The profitability indices show that 'acadja' brush park fishing practice is a profitable business in the lagoon. Every N1 invested in the business generated N1.22 as profit. According to Ayinde and Aromolaran (1998) and Abudl *et al.* (2004) calculated rate of return to investment is acceptable only if the opportunity cost of capital and family labour utilised in agric-business is zero. For every N1 spent on hired labour about double the amount went back to the fisherman. According to Otubusin (2001) and Emmanuel (2009) economic viability of any fisheries business depends on the inter play of various components of the operation. In addition to this the fisherman should endeavour to have more acadja for more profit.

**\*Correspondence to:**

Dr Emmanuel, Babatunde Eniola  
Department of Marine Sciences  
University of Lagos, Akoka, Lagos, Nigeria.  
Cellular Phone: 234 – 802 – 853 – 945- 9  
Email: [monetemi@yahoo.com](mailto:monetemi@yahoo.com)

**REFERENCES**

- [1] Webb, J.E. and Hill, M.B. (1958). The ecology of Lagos lagoon. *Phil. Trans. Roy. Soc. Lond.* 241, B(683): 307 – 419.
- [2] Sandison, E.E., and Hill, M.B. (1966): The distribution of *Balanus pallidus* (Darwin) *Gryphaea gasar* Dautzenberg, *Mercierella enigmatica* (fauvel) and *Hydroides uncinata* (Philippi) in relation to Salinity in Lagos harbour and adjacent Greeks. *J. Animal Ecology* **35**: 235 – 250.
- [3] Oyenekan, J.A. (1988): Benthic macrofaunal communities of Lagos Lagoon, Nigeria. *Nig. J. Sci.* **21**: 45 – 51.
- [4] Brown, C.A.(1998). Distribution and population dynamics of an estuarine population of *Aloidis trigona* Hinds (Bivalvia). *Acta Hydrobiol.* 40(4): 227 -237.
- [5] Brown, C.A. and Oyenekan, J.A. (1998): Temporal variability in the structure of benthic macrofaun community of the Lagos lagoon and harbour, Nigeria. *Pol. Arch. Hydrobio.* **45(1)**: 45 – 54.
- [6] Fagade, S. O. (1969). Studies on the biology of some fishes and the fisheries of the Lagos lagoon. Ph.D. Thesis University of Lagos, Nigeria. 385pp
- [7] FAO (1969). Fisheries survey in the western and mid-western region of Nigeria. UNDP/FAO Rome SF:74/NIR6. 142pp.
- [8] Fagade S.O. and Olaniyan, C.I. O. (1974). Seasonal distribution of the fish fauna of the Lagos lagoon. Bulletin de l' I L F.A.N. T.XXXVI, Ser A. pp. 244 - 252.
- [9] Udolisa R.E.K and Solarin B.B. (1979). Design characteristics of cast nets and gillnets in Lagos lagoon, Nigeria. NIOMR Occ. Paper, 31: 24.
- [10] Solarin, B.B. (1998): Hydrobiology, fishes and fisheries of the Lagos Lagoon, Nigeria. Ph.D Thesis, University of Lagos. 256pp.
- [11] Solarin, B.B. and Kusemiju, K. (2003a). Fish shelters as fisheries enhancement techniques in the Lagos lagoon, Nigeria. *Nigerian Journal of Fisheries*.1: 57 – 61
- [12] Solarin, B.B. and Kusemiju, K. (2003b). The fishery and aspects of gillnet designs and operation in the Lagos lagoon, Nigeria. *Nigerian Journal of Fisheries*.1: 62 - 69
- [13] Emmanuel, B.E., Chukwu, L.O. and Azeez, L.O. (2008a). Cast net design characteristics, catch composition and selectivity in tropical open lagoon. *African Journal of Biotechnology*.7 (12): 2081 – 2089.
- [14] Emmanuel, B.E., Chukwu, L.O. and Azeez, L.O. (2008b). Gill net selectivity and catch rates of pelagic fish in tropical coastal lagoonal

- ecosystem. *African Journal of Biotechnology* . 7 (21), 3962-3971
- [15] Ajao, E.A., Oyewo, E.O. and Uyimadu, J.P. (1996): A review of the pollution of coastal waters in Nigeria. Nigerian Institute of Oceanography Technical Paper No. 107. 20pp.
- [16] Akpata. T.V.I., Oyenekan, J.A. and Nwankwo, D.I. (1993): Impacts of organic pollution on bacterial, plankton and benthic populations of Lagos lagoon, Nigeria. *International Journal of Ecology and Environmental Sciences*. **19**: 73 – 82.
- [17] Nwankwo, D.I. (1998). Seasonal changes in phytoplankton composition and diversity in Epe lagoon, Nigeria. *Acta Hydrobiol.* 40 (2): 83 - 92
- [18] Emmanuel, B.E. (2008) The Fishery and bionomics of the swimming crab, *Callinectes amnicola* (DeRocheburne, 1883) from a tropical lagoon and its adjacent creek, Southwest, Nigeria. *Journal of Fisheries and Aquatic Science* 3(2)114-125
- [19] Welcomme, R.L. (1972). An evaluation of the acadja method of fishing as practiced in the coastal lagoons of Dahomey (West Africa). *J. fish. Biol.* 4: 39 – 55.
- [20] Welcomme, R. L. and Bartley, D.M. (1998). An evaluation of present techniques for the enhancement of fisheries in Petr, T. (ed). *Inland fishery enhancements*. FAO Fisheries Technical Paper No. 374, 1-36
- [21] Nwankwo, D.I. (1991): Periphyton algae on fish fences “Acadja” in a tropical open lagoon, Nigeria. *Nig. J. Bot.*, **3**: 197 – 207.
- [22] Nwankwo, D.I., Abosede, A.O. and Abdulrasaq, Y. (1994): Floating timber logs as a substrate for periphyton algae in the Lagos Lagoon, Nigeria. *Pol. Arch. Hydrobiology* **4(4)**: 419 – 430.
- [23] Emmanuel, B.E. and Onyema, I.C. (2007). The plankton and fishes of a tropical creek in south-western, Nigeria. *Turkish Journal of Fisheries and Aquatic Science*.7: 105-114
- [24] Webb, J.E. (1958): The Ecology of Lagos Lagoon. 1: The Lagoons of the Guinea Coast. *Phil. Transc. Roy. Sco. Lond. Ser. B*. 307 – 318.
- [25] Brown, C.A. (1991). Community structure and secondary production of benthic macrofauna of the Lagos lagoon and harbour. M.Phil. Thesis. University of Lagos, Nigeria.
- [26] Oyenekan, J.A. (1975). A survey of the Lagos Lagoon benthos, with particular reference to the Mollusca M.Sc. *Dissertation* University of Lagos. 288pp.
- [27] Ogbeibu, A.E. (2005). *Biostatistics: A practical approach to research and data handling*. Mindex Publishing Company Limited., Nigeria.264pp
- [28] Margalef, D.R. (1951). Diversidad de especies en les comunideades natural *Public Institutte of Biologic Barcelona* 9 : 5 – 27.
- [29] Shannon, C.E and Wiener, W.( 1949) The Mathematical Theory of Communcation Urban. University of Illinois Press. 125.
- [30] Lloyd, M. and Ghellardi, R. G. (1964). A table for calculating the equitability components of species diversity. *J. Anim.Ecol.* 33: 217-225.
- [31] Ayinde, I.A. and Aromolaran, A.B. (1998): Economics of rabbit production in Abeokuta South Local Government Area of Ogun State, Nigeria. *Nigerian Journal of Animal Production* **25(1)**: 100 – 105.
- [32] Abdul, W.O., Omoniyi, I.T., Udolisa, R.E.K. and Alegbeleye, W.O. (2004): Economic potentials of “Ikern” brush park fishing practice in \_\_\_\_\_ lagoon, Ogun State, Nigeria. In 18<sup>th</sup> Annual Conference of the Fisheries Society of Nigeria (FISON) 8 – 12 Dec. 2003. Owerri Nigeria. 126 – 130.
- [33] Chukwu, L.O. and Nwankwo, D.I. (2004). The impact of land based pollution on the hydrochemistry and macrobenthic community of a tropical West African creek. *Ekologia (Asia)* **2(1)**: 1-9.
- [34] Hill, M.B. and Webb, J.E. (1958a): The lagoons of the Guinea Coast. In: *The ecology of the Lagos Lagoon*. Cambridge University, V. B. **241**: 301 – 419. [36] Williams, D.D. and Feltmate, B.W. (1992): *Aquatic insects*. CAB International, Wallingford.
- [35] Williams, D.D. and Feltmate, B.W. (1992): *Aquatic insects*. CAB International, Wallingford.
- [36] Mackie, G.L. (1998): *Applied Aquatic Ecosystem Concepts*. Course Pack, University of Guelph, Guelph.
- [37] Olomukoro, J.O. and Ezemonye, L.I.N. (2007): Assessment of the macroinvertebrate fauna of rivers in Southern Nigeria. *African zoology* 42(1): 1 – 11.
- [38] Otubusin, S.O. (2001): Economics of Small Scale tilapia production in floating net cages. Asset.
- [39] Emmanuel, B.E. (2009). The artisanal fishing gears, crafts technology and their efficiency in the Lekki lagoon, Nigeria. Ph.D Thesis, University of Lagos.268pp

7/9/2009