

Cyanobacteria of a Tropical Lagoon, Nigeria.

Adesalu, Taofikat Abosede¹, Nwankwo, Dike Ikegwu.²

¹Department of Botany and Microbiology, University of Lagos, Nigeria.

²Department of Marine sciences, University of Lagos, Nigeria.

boseadesalu@yahoo.com.

Abstract: Investigations for the first time into the blue green algae of Lekki lagoon were carried out for 24 months (June 2003- May 2005) at monthly intervals using standard plankton net of mesh size 55µm. One hundred and seventy nine species belonging to thirty genera were observed. The filamentous blue green algae *Oscillatoria* formed the most abundant genus making up twenty three species followed by *Phormidium* eighteen species. *Anabaena* and *Chroococcus* recorded thirteen species each while the genera, *Gleocapsa*, *Merismopedia* and *Microcystis* recorded ten, eight and twelve species respectively. Only one genus each of *Cyanosarcina*, *Calothrix* and *Scytonema* were encountered. Bloom forming species identified were *Microcystis aeruginosa*, *M. flos-aquae*, *M. wesenbergii* and *Anabaena flos-aquae*. In this study, thirty-nine new species were recorded for Lagos lagoon complex in which Lekki lagoon is one of it while *Cyanosarcina hueberliorum* is new record for Nigeria. [Journal of American Science 2010;6(4):193-199]. (ISSN: 1545-1003).

Keywords: Cyanophytes, tropical, bloom, Lagos lagoon complex

Introduction

The coastline of South Western Nigeria is a meandering network of lagoons and creeks of which Lagos lagoon with an area of 208sqkm is the largest (Nwankwo 1989). The geography and hydrology of various parts of Lagos lagoon complex in which Lekki lagoon is one it have been described by several workers. These include Lekki lagoon (Ikusemiju 1973); Lagos lagoon (Hill and Webb 1958) and harbour (Olaniyan 1957). Checklists of planktonic algae in some parts of Nigeria have been documented by different workers. For instance in the North, Holden and Green (1960) studied the phytoplankton of River Sokoto while Khan and Agugo (1990) studied Kongiri dam, Jos mine lakes was studied by Anadu et al. (1990).

In Southern region, studies include Opupe (1990,1991,1992) who studied Warri Forcados estuary phytoplankton, New Calabar river by Nwadiaro and Ezefili (1986). Biswas (1984, 1992) had report for eastern region while western region reports include that of Imevbore (1968) on Eleiyele reservoir, Egborge and Sagey (1979) on Ibadan freshwater ecosystem . Nwankwo (1988) studied the planktonic algae of Lagos lagoon, Nwankwo (1993) reported eight cyanobacteria bloom species of coastal waters in South Western Nigeria excluding Lekki lagoon, Nwankwo (1997) reported dinoflagellates list of Lagos lagoon. Adesalu and Nwankwo (2005, 2009) reported the diatoms of Olero creek and Lekki lagoon respectively, Wujek et al. (2003) studied the chrysophytes of Lekki lagoon while Kadiri (1989,

1993, 1999, 2000) reported the rich flora of *Micrasterias*, desmids, algae composition and euglenoids of Ikpoba reservoir respectively.

Of the entire aforementioned checklist, none specifically reported the cyanobacteria checklist in Nigeria coastal waters. The present study was undertaken to investigate the composition of cyanobacteria species of Lekki lagoon for possible biological monitoring since the lagoon is a source of fish supply for people of South Western states and beyond.

Description of study area

Lekki lagoon (Fig.1) a large expanse of shallow freshwater extends between Lagos and Ogun states. It covers an area of about 247km². A greater part of the lagoon is shallow (<3.0 m), while some areas are up to 6.4m deep. It lies between longitudes 4°00' E and 4°12' E and latitude 6°25' N and 6°37' N. The lagoon is fed by river Oni in the north eastern part, while rivers Osun and Saga flow into the north western part. Two peaks of rainfall are associated with this lagoon, a major peak in July and a lesser peak in September. There are two peaks of sunshine hours which approximately correspond to the equinoxes. The mainstay of communities that live around this environment is artisanal fishing.

TABLE 1 : AVERAGE DEPTH (M) AND GEOGRAPHICAL POSITION OF SAMPLING STATIONS




Fig.1: Lekki Lagoon showing sampling stations

Materials and methods

Collection of sample

Biological samples were collected monthly from twelve stations (Table 1) using Hydrobios plankton net of 55 μ m mesh size. For quantitative analysis 5litres of the water was concentrated. Biological samples were preserved in 4% unbuffered formalin. Identification was done using Olympus BX51 photomicroscope. Water samples were collected into clean plastic containers for chemical analysis while *in situ* measurements of temperature, transparency, pH and depth were made.

Physical and chemical analysis of water sample

The methods described by America Public Health Association (APHA 1998) were used for physical and chemical analysis. The air and surface water temperature were measured *in situ* with a simple mercury thermometer while the transparency was measured using a 20cm diameter Secchi disc. The depth was measured with a calibrated pole and the water pH determined using a Phillips pH meter (Model PW950). The chemical factors determined include Salinity, conductivity, dissolved oxygen (DO) and biological oxygen demand (BOD_5). Salinity was determined using the Silver Nitrate

STATIONS	Average depth	Longitudes	Latitudes
Emina 1	2.77	4°5.080E	6°32.754N
Emina II	1.61	4°7.511E	6°34.07N
Entrance of River			
Mosafejo	1.51	4°10.239E	6°35.344N
Entrance of River Oni	1.80	4°12.153E	6°35.090N
Iwopin 1	1.80	4°13.153E	6°32.309N
Iwopin 11	2.69	4°9.651E	6°32.137N
Imoki	2.17	4°10.048E	6°31.253N
Ise 1	1.81	4°13.413E	6°26.833N
Ise 11	2.41	4°9.788E	6°26.181N
Ebute lekki	1.29	4°5.353E	6°26.685N
Entrance of Omu creek	1.88	4°7.604E	6°28.867N
Lagoon centre	2.23	4°3.348E	6°28.577N

Chromate titration method as described by Barnes (1980) while Dissolved oxygen content was determined using a Griffin oxygen meter. Oxygen saturation was recorded in percentage. Biological Oxygen Demand is the measure of the amount of dissolved oxygen that could be depleted from the water body during natural biological assimilation or degradation of organic compounds by the organisms present especially bacteria. This was done after the dissolved oxygen had been measured using the standard method of biochemical consumption of oxygen in 5 days at 20°C while conductivity was determined using the HANNA instrument (HI8733), a wide range conductivity meter that has salinometer range in μ S. Conductivity values were recorded as $mScm^{-1}$ at 25°C (APHA 1998). The department The Federal Meteorological Department, Oshodi, Lagos kindly provided rainfall and sunshine hours data for the period of investigation (Table 2)

Nutrient determination

For nitrate determination, Hach Cadmium reduction method was used (APHA 1998). Phosphate-phosphorus is known to be important in a number of ways, one being that it facilitates the uptake of

nitrogen. It was determined by ascorbic acid method. The values obtained were recorded in milligrams per litre (mgL^{-1}) (APHA 1998) (Table 2)

Water chemistry

The physical and chemical characteristics of the study area are presented in Table 2. The mean pH of the water with a range of 7.41-7.46 indicated that the system is highly buffered. Conductivity which is the numerical expression of the ability of a solution to carry an electric current represents the total ions of water ranged between 0.47-0.56 μScm^{-1} the lowest value for phosphate-phosphorus, nitrate-nitrogen and sulphate were 2.42, 2.70 and 0.002 mgL^{-1} . Salinity recorded the least value of 0.40‰ while chloride ion had the highest value of 10.00 mgL^{-1} .

	Biological Oxygen demand	0.23	0.22	0.26	0.23	0.22	0.25
Chemical oxygen demand	0.36	0.30	0.26	0.25	0.24	0.29	
Oil and grease	0.03	0.02	0.02	0.02	0.00	0.01	
Turbidity (FTU)	8.59	9.25	10.00	7.46	9.57	8.87	
Total dissolved solids	9.48	10.29	10.65	8.51	11.29	11.50	
Sulphate	0.03	0.02	0.03	0.04	0.03	0.03	
Ca	10.46	11.11	11.92	12.07	11.57	10.71	
Fe	0.30	0.23	0.24	0.26	0.22	0.25	
Pb	0.01	0.08	0.25	0.08	0.08	0.01	
Hg	0.01	0.00	0.00	0.00	0.00	0.00	
Cu	0.05	0.02	0.02	0.02	0.03	0.06	
Ni	0.02	0.01	0.02	0.01	0.01	0.01	
Zn	0.02	0.01	0.01	0.01	0.01	0.02	

Table 2: Mean physico-chemical values for Lekki lagoon (concentrations in mg L^{-1}) (Stations A-L).

STATIONS	A	B	C	D	E	F
Surface Water tempemperature ($^{\circ}\text{C}$)	30.54	30.73	30.55	30.61	30.40	30.31
Air tempemperature ($^{\circ}\text{C}$)	29.68	29.60	29.60	29.71	29.51	29.44
Transparency (cm)	5.74	7.38	11.33	9.25	11.67	10.29
Total suspended solids	9.07	8.56	9.16	8.45	9.28	9.78
pH	7.42	7.43	7.41	7.43	7.44	7.46
Salinity %	0.47	0.47	0.45	0.45	0.40	0.55
Phosphate-phosphorus	2.53	2.70	2.42	2.50	2.56	2.61
Nitrate-nitrogen	2.73	2.93	3.42	2.98	2.54	3.85
Chloride	9.23	9.67	9.59	10.00	9.88	9.52
Conductivity (μScm^{-1})	0.56	0.52	0.56	0.47	0.47	0.48
Dissolved Oxygen	4.15	3.46	4.21	4.10	4.16	4.25
Biological Oxygen demand	0.23	0.22	0.26	0.23	0.22	0.25
Chemical oxygen demand	0.36	0.30	0.26	0.25	0.24	0.29
Oil and grease	0.03	0.02	0.02	0.02	0.00	0.01

STATIONS	G	H	I	J	K	L
Surface Water tempemperature ($^{\circ}\text{C}$)	30.35	30.23	30.38	30.63	30.70	30.55
Air tempemperature ($^{\circ}\text{C}$)	29.55	29.08	29.29	28.90	29.27	29.21
Transparency (cm)	11.08	10.42	14.42	7.30	10.04	12.33
Total suspended solids	11.76	12.71	8.39	11.50	8.87	8.87
pH	7.44	7.46	7.44	7.42	7.37	7.38
Salinity %	0.47	0.44	0.45	0.44	0.50	0.47
Phosphate-phosphorus	2.47	2.49	2.96	2.53	2.54	2.44
Nitrate-nitrogen	3.44	3.25	2.75	3.31	2.70	3.97
Chloride	9.40	9.17	9.31	9.31	9.54	9.99
Conductivity (μScm^{-1})	0.49	0.52	0.46	0.53	0.62	0.56
Dissolved Oxygen	4.25	4.19	4.17	4.16	4.18	4.17
Biological Oxygen demand	0.29	0.30	0.31	0.28	0.26	0.24
Chemical oxygen demand	0.34	0.32	0.30	0.31	0.31	0.25
Oil and grease	0.01	0.01	0.01	0.01	0.01	0.01

Turbidity (FTU)	8.37	7.99	8.42	10.29	9.15	8.69
Total dissolved solids	9.87	9.95	10.31	10.58	10.81	10.29
Sulphate	0.02	0.02	0.05	0.04	0.06	0.03
Ca	11.04	10.96	10.70	10.21	11.40	11.52
Fe	0.21	0.23	0.25	0.28	0.30	0.55
Pb	0.02	0.01	0.00	0.00	0.04	0.02
Hg	0.00	0.00	0.00	0.00	0.02	0.01
Cu	0.01	0.01	0.01	0.00	0.02	0.04
Ni	0.01	0.01	0.01	0.01	0.01	0.01
Zn	0.01	0.01	0.01	0.01	0.01	0.01

Results

Cyanobacteria analysis

In this study, cyanobacteria genera are arranged alphabetically within families and the species in alphabetical order within genera (Table 3).

Table 3: Cyanobacteria checklist at Lekki lagoon, Nigeria.

Division : Cyanophyta

Class : Cyanophyceae

Order: Chroococcales

Family 1: Chroococcaceae

Chroococcus decorticans

C. dispersus (V.Keiss) Lemm.

C. limnecicus Lemm.

**C. limnecicus var. subsalsus* Lemm.

C. minor

C. minutus (Kutz.) Rabenh.

C. palidus Nageli

C. prescottii Dr. & Daily

C. turgidus (Kutz.) Lemm.

C. turicensis (Nag.) Hangirg

C. varius A. Braun

Chroococcus sp 1

Chroococcus sp 11

**Cyanosarcina huebeliorum* Komarek & Anagnostidis

Dactylococcopsis raphidioides Hansg.

D. smithii Chodat & Chodat

Dactylococcopsis sp.

Family 2: Merismopediaceae

Agmenellum quadruplicatum

Agmenellum sp.

Aphanocapsa delicatissima West & West

A. elaschista West & West

A. elachista var conferta West & West

**A. nubilum* Nygaard

A.pulvrea

A. rivularis

Aphanocapsa sp.

**Aphanothece bullosa* var *major* Geitler

Aphanothece sp.

Merismopedia angularis Geitler

M. convoluta Breb.

M. elegans A.Br.

M. glauca (Ehr.) Nag.

M. major G.M.Smith

M. marsonii Lemm.

M. punctata Meyen.

M. tenuisima Lemm.

Family 3: Chaemasiphonaceae

Clastidium setigerum

Clastidium sp.

Family 4: Microscystaceae

**Gleocapsa alpicola* (Lyng.) Bornet

**G. arenaria* (Hass.) Rabenh.

G. biformis Novacek

G. compacta Kutz.

G. conglomerata Kutz.

**G. decorticans* (A.Br.) P.Richter

G. delicatissima

G. magma (Breb.) Kutz.

G. querternata (Breb.) Kutz

Gleocapsa sp.

Gleothece heufleri

Gleothece linearis Nag

Gleothece sp.

Microcystis aeruginosa Kutz.

M. aeruginosa var *elongata* Rao,C.B

**M. aeruginosa* var *major* (Witt.) Smith

M. elongata sp.nov.

**M. firma* (Kutz.) Dr. & Daily

M. flos-aquae (Witt.) Kirchner

M. paludosus

**M. pulvrea* (Wood) Forti

**M. ramosa* Bharadwaja

**M. robusta* (Clack) Nygaard

M. viridis (A. Br.) Lemm.

M. wesenbergii Kosinskaja

Order 2: Nostocales

Family: Nostocaceae

Anabaena azollae Strasburger

A. circinalis (Kutz.) Rabh.

**A. confervoides* Reinsch

A. constricta Lauter b.

A. cycadeae J.Reinsch

A. cylindrica Lemmermann

A. fircinalis

A. flos-aquae (Lyng.) Breb.

A. limnectica G.M.Smith

<i>A. spiroides</i> Lemm.	<i>Arthrospira</i> sp.
* <i>A. torulosa</i> (Carm.) Lagerh.	<i>Microcoleus codii</i> Fremy
<i>Anabaena</i> sp.1	<i>M. subtorulosus</i>
<i>Anabaena</i> sp 11	<i>M. willeana</i>
<i>Aphanabaena</i> sp	<i>Microcoleus</i> sp.1
<i>Calothrix</i> sp	<i>Microcoleus</i> sp 11
<i>Cylindrospermum catenatum</i> Ralfs	<i>Phormidium angustissimum</i> West & West
<i>Cylindrospermum majus</i> Kutz.	* <i>P. caeruleascens</i> Geitler
<i>Cylindrospermum</i> sp 1	* <i>P. chlorinum</i> Komarek
<i>Nostoc carneum</i>	<i>P. cortianum</i>
<i>Nostoc linkia</i>	<i>P. crouanii</i> Gomont
<i>N. muscorum</i> Agardh	<i>P. foveolarum</i> (Mont.) Gomont
<i>N. peltigerae</i> Letellier	* <i>P. insigne</i> Skuja
<i>N. sphaericum</i> Vauch.	* <i>P. laetevirens</i> Skuja
<i>Nostoc</i> sp.1	<i>P. luridum</i> (Kutz.) Gomont
<i>Nostoc</i> sp 11	* <i>P. luteum</i> Kosinskaja
Order 3: Oscillatoriales	* <i>P. molle</i> Palik
Family 1: Oscillatoriaceae	<i>P. nigro-viride</i> Gomont
<i>Lyngbya birgei</i> G.M.Smith	* <i>P. papyraceum</i> (Ag.) Gom.
<i>L. contorta</i> Lemm.	<i>P. retzii</i> (Ag.) Gomont
* <i>L. lagerheimia</i> (Mobius) Gom.	<i>P. tenue</i> (Menegh.)Gom.
<i>L. limnectica</i> Lemmermann	<i>P. tinctorium</i> Kutz.
* <i>L. martensiana</i> Menegh.	<i>Phormidium</i> sp 1
<i>L. versicolor</i> (Wattman)Gomont	<i>Phormidium</i> sp 11
<i>Lyngbya</i> sp	* <i>Planothrix clavarata</i> Skuja
<i>Oscillatoria acuminata</i> Gomont	* <i>P. cryptovaginata</i> Skacelova & Komarek
<i>O. acutissima</i> Kufferath	* <i>P. isothrix</i> Komarek
<i>O. agardii</i> Gomont.	<i>P. minor</i>
<i>O. angustissima</i> West & West	<i>P. planctonica</i>
<i>O. articulata</i>	<i>Planothrix</i> sp1
<i>O. brevis</i> Kutz.	<i>Planothrix</i> sp 11
<i>O. curviceps</i> Agardh	<i>Trichodesmium laucustre</i> Klebahn
<i>O. formosa</i> Bory.	<i>Trichodesmium</i> sp.
* <i>O.formosa</i> f. <i>edaghica</i> Novickova	Family 3:Pseudanabaenaceae
<i>O. germinata</i> Meneghini	* <i>Limnothrix planctonica</i> Geitler
<i>O. lacustris</i>	<i>Limnothrix</i> sp.
* <i>O. lemmermanni</i> Wolosz	* <i>Pseudoanabaena curta</i> Hollerbach
<i>O. limnectica</i> Lemm.	* <i>P. moniliformis</i> Komarek & Kling
<i>O. limosa</i> (Roth) Ag.	* <i>P. thermalis</i> Anagnostidis
<i>O. margaritifera</i> Kutzing (Gomont)	<i>Pseudoanabaena</i> sp
<i>O. minima</i>	<i>Spirulina filiformis</i>
<i>O. planctonica</i> Wolosz	<i>S. princeps</i> W.et G.S.West
* <i>O. rubescens</i> DC ex Gomont	<i>S. major</i> Geitler
<i>O. sancta</i> (Kutz.)Gom	<i>S. meneghiniana</i> Anagnostidis
* <i>O. simplissima</i> Gomont.	<i>S. tenerima</i>
<i>O. subrevis</i> Schmidle	<i>Spirulina</i> sp 1
<i>O. tenuis</i> Ag.	<i>Spirulina</i> sp 11
<i>Oscillatoria</i> sp.	* <i>Leptolyngbya hypolimnectica</i>
Family 2:Phormidiaceae	* <i>Leptolyngbya ocricana</i> Cardo
<i>Arthrospira fusiformis</i> Fott & Karim	<i>L. tenuis</i>

Leptolyngbya sp.

**Plantolyngbya brevicellularis* Cronberg
&Komarek

**P. minor* Komarek&Cronberg

**P. tallingii* Komarek & Kling

**P. minor*

Planktolyngbya sp.

Family:Schizotrichaceae

Schizothrix pulvinata

S. friesii (Ag.) Gomont

Schizothrix sp.

Scytonema sp.

Discussion

The cyanobacteria checklist reflects the influence of hydrological conditions of this area. The dominance of *Oscillatoria* throughout the season could be a pointer that the hydrology and salinity of the studied area favours its growth. The particular high diversity of blue-green observed in the lagoon could also be that the water chemistry favours growth of cyanophytes. Five bloom forming cyanophytes identified in this study include *Microcystis aeruginos*, *M. wesenbergii*, , *Anabaena flos-aquae*, *A. spiroides* and *Oscillatoria formosa*.

The variation in physical and chemical parameters observed during the study period may be as a result of the influence of weather conditions. For instance, the rainy season occurring between June and October, characterized by low transparency and pH; increased total suspended solids, higher turbidity and increased flood water condition which might have initiated stressful environmental condition and these conform with Dart and Stretton (1980) who stated that variations in water temperature could cause alterations in the pH due to changes in ionization and increased solubility or precipitation of bottom deposits. Nwankwo and Onitiri (1992) also pointed out that it is possible that rainfall triggers off flood situations which usually increases total solids, reduces transparency and consequently light penetration and also dislodges attached algal forms. The phytoplankton community and the physico-chemical parameters exhibited seasonal changes closely related to the pattern of rainfall. The presence or absence of any blue-green species may be due to the changing physical environment other than pollution (Nwankwo 1994).

Acknowledgements

Adesalu T.A. is grateful to Fulbright exchange program scholarship, University of Lagos, Nigeria; Central Michigan and Bowling Green state

Universities U.S.A.. Thanks are also due to Dr Akinsoji, O. my co-supervisor.

Corresponding author:

Adesalu, Taofikat Abosede
Department of Botany and Microbiology
University of Lagos, Akoka, Nigeria
E-mail: boseadesalu@yahoo.com.

References

1. Adesalu,T.A., Nwankwo, D.I. Studies on the phytoplankton of Olero creek and parts of Benin river, Nigeria. The Ekologia. 2005; 3(2):21-30.
2. Adesalu,T.A., Nwankwo, D. I. A checklist of Lekki lagoon diatoms. International Journal of Botany. 2009; (5) 2: 126-34.
3. American Public Health Association. Standard methods for the examination of water and waste water, 20th ed. American Public Health Association Incorporated, New York. 1998: 1270pp.
4. Anadu, D.I.,Obioma, A., Ejike, C. Water chemistry and plankton periodicity in two contrasting mine lakes in Jos,Nigeria. Hydrobiologia, 1990; 206:17-25
5. Biswas, S. Phytoplankton periodicity of Ogelube Lakes, Anambra State, Nigeria. During 1979-1980.Hydrobiologia, 1992; 246: 169-172.
6. Biswas, S. Phytoplankton of Opi Lakes, Anambra State, Nigeria.Verht.Int.Verein. Limnology, 1984; 22:1584-1589.
7. Dart, R. K., Stretton, R. J. Microbiological aspects of pollution control and environmental science. 6 Elsevier Scientific Publishers Amsterdam 1980.
8. Egborge, A. B. M., Sagay, E. G.. The distribution of phytoplankton and zooplankton in some Ibadan freshwater cosystems. Polskie Archiwum Hydrobiologii, 1979; 26: 23-335.
9. Hill, M. B., Webb, J. EThe ecology of Lagos lagoon II. The topography and physical features of Lagos harbour and Lagos lagoon. Phil. Trans. Royal Society London, 1958; 241: 319-417.
10. Holden, M. J., Green, J. The hydrology and plankton of the River Sokoto. Journal of Animal Ecology, 1960; 29: 65–84.
11. Ikusemiju, K. A study of catfishes of Lekki lagoon with particular reference to species of *Chrysichthys walkeri* (Bagridae) Ph.D Thesis University of Lagos, Nigeria, 1973.

12. Imevbore, A. M. A. Planktonic algae of Eleiyele reservoir, Nigeria. *Journal of Science*, 1968; 2: 85–90.
13. Kadiri, M.O. A rich flora of Micrasterias from Ikpoba reservoir, Nigeria. *Archiwum Hydrobiologii*, 1989; 116(3): 391-399.
14. Kadiri, M.O. Further desmids from Ikpoba reservoir (Nigeria) compared with other records from Africa. *Algological studies*, 1993; 71:23-35.
15. Kadiri, M. O. A comprehensive preliminary checklist of the algae of the Ikpoba reservoir Edo state,Nigeria. *Global journal of pure and applied sciences*, 1999; 5 (4): 485-491.
16. Kadiri, M.O. Seasonal trend in the chemical limnology of a shallow Nigerian man-made lake. *Acta Hydrobio*. 2000; 1 (42): 29-40.
17. Nwankwo, D. I., Onitiri, A.O. Periphyton (Aufwuchs) community on submerged aquatic macrophytes in Epe lagoon, Nigeria. *Journal of Agriculture Science and Technology*. 1992; 2 (2): 135- 41.
18. Nwankwo, D. I. A checklist of Nigerian marine algae. (Tarkwa Bay). *Nigerian Journal of Botany*, 1988; 1: 47–50.
19. Nwankwo, D. I. Cyanobacteria bloom species in coastal waters of South-western Nigeria. *Archiwum Hydrobiologie Supplement*, 1993; 90: 553–542.
20. Nwankwo D. I. Hydrochemical properties and bottom-dwelling of aLagos lagoon sewage disposal site. *Polskie Archiwum Hydrobiologia*, 1994; 41(1): 35-47.
21. Nwankwo, D. I. A first list of dinoflagellates (Pyrrophyta) from Nigerian coastal waters (creeks, estuaries,lagoons).*Polskie Archiwum Hydrobiologia*, 1997; 44(3): 313–321.
22. Olaniyan, C. I. O. The seasonal variation in plankton in Lagos harbour, Nigeria. Ph.D Thesis, University of London, London, 1957; 173pp.
23. Opute, F. I. Phytoplankton flora of the Warri/Forcados estuaries of Southern Nigeria.*Hydrobiologia*. 1990; 208:101-109.
24. Opute, F.I. A checklist of the freshwater, brackish and marine phytoplankton of the Warri/Forcados estuaries of southwestern Nigeria. *Nigerian Journal of Botany*.1991; 4: 227-254.
25. Opute, F.I. Contribution to the knowledge of algae of Nigeria 1: Desmids from the Warri/Forcados estuaries, Part 2. The genera *Euastrum* and *Micrasterias*. *Archiwum Hydrobiologii*. Supplement, 1992; 93: 73-92.
26. Wujek, D. E., Adesalu, T. A., Nwankwo D. I. Silica-scaled chrysophyceae and synurophyceae (chrysophyta) from Nigeria II Lake Lekki. *Tropical Freshwater Biology*. 2003/2004; 12/13:99-103.

9/9/2009