

Protective Effect of some Antioxidants against CCl₄-Induced Toxicity in Liver Cells from BRL3A Cell Line.Kamel, H.H.¹, Azza, H. Abd-El-Rahman¹, Walaa, M.S. Ahmed¹ and Amira, H. Mohamed²¹Clinical Pathology Dept., Fac. of Vet. Med. Beni-Suef University, Beni-Suef, Egypt²Clinical Pathology Dept., Fac. of Vet. Med. Cairo University, Giza, Egypt

Azza_abdelrahman92@yahoo.com

Abstract: An *in vitro* experiment was conducted to investigate the protective effect of ascorbic acid, mannitol and aminoguanidine at different concentrations against carbon tetrachloride induced toxicity and oxidative stress in hepatocytes cell line (BRL3A) from buffalo rats. Results were compared with those of vitamin E as standard hepatoprotective agent. Treatment of BRL3A with CCl₄ lead to generation of free radicals detected after two hours incubation using ESR technique and produced cell injury demonstrated by increased leakage of LDH, ALT and AST to the media. Exposure to CCl₄ caused apoptosis to cells but did not induce lipid peroxidation as tested by the TBARS technique. Treatment with vitamin E has significant hepatoprotective effect by lowering the leakage of intracellular enzymes, reducing the oxidation of proteins and decrease incidence of apoptosis. Ascorbic acid, mannitol and aminoguanidine were ineffective against CCl₄ toxicity.

[Kamel, H.H., Azza, H. Abd-El-Rahman, Walaa, M.S. Ahmed and Amira, H. Mohamed. **Protective Effect of some Antioxidants against CCl₄-Induced Toxicity in Liver Cells from BRL3A Cell Line.** Journal of American Science 2011; 7(2):283-296]. (ISSN: 1545-1003). <http://www.americanscience.org>.

Keywords: Protective Effect; antioxidants; CCl₄-Induced Toxicity; Liver; Cell; BRL3A Cell Line

1. Introduction:

The liver occupies a vital role in the main functions of the organism. It is particularly susceptible to chemically induced injury due to its extensive metabolic capacity and cellular heterogeneity. Oxidative stress occurs when there is an imbalance between reactive oxygen species (ROS) formation and scavenging by antioxidants. Excess generation of ROS can cause oxidative damage to biomolecules resulting in lipid peroxidation, mutagenesis and carcinogenesis (Khan and Sultana, 2009).

In vitro liver systems represent a better experimental approach to screen potential hepatotoxic compounds and to investigate the mechanism by which chemicals induce liver lesions (Kucera *et al.*, 2006). Liver cell lines are characterized by unlimited subcultivation and cell availability in large number) Guillouzo, 1998). The BRL3A cell line is an epithelial cell line from buffalo rat liver which is able to divide in the absence of serum.

Carbon tetrachloride (CCl₄) has long been known as a model toxicant and has been the focus of many *in vitro* and *in vivo* toxicological studies (Manibusan *et al.*, 2007). The liver is the major target organ of CCl₄ toxicity owing to its high content of cytochrome P-450 (Södergren *et al.*, 2001). Antioxidants are used to antagonize the deleterious action of free radicals and to protect hepatocytes from damage. Vitamin E (α-tocopherol) is considered the most important lipophilic antioxidant in biological tissues (Weber *et al.*, 2003). Ascorbic acid is an important dietary antioxidant. It significantly

decreases the adverse effect of reactive species such that can cause oxidative damage to macro-molecules such as lipids, DNA, and proteins (Alpsoy *et al.*, 2009). Mannitol is a polyol which develops an antioxidant activity by scavenging ROS (Mendoza *et al.*, 2007). A number of physiological functions have been ascribed to mannitol, including serving as a reserve carbon source, as an antioxidant, and to store reducing power (Ruijter *et al.*, 2003). Aminoguanidine is an irreversible inhibitor of the inducible form of nitric oxide synthase (iNOS), which also inhibits endothelial NOS (eNOS) and neuronal NOS (nNOS) (Mohamad *et al.*, 2009).

In the present study, *in vitro* antioxidant activities of ascorbic acid, mannitol and aminoguanidine were assessed in comparison with vitamin E against hepatotoxicity induced by carbon tetrachloride in liver cell line from rats.

2. Material and methods:**a. Material:-**

CCl₄ (99.5%), vitamin E "α-tocopherol acetate" and L (+) ascorbic acid "vitamin C" were purchased from Wako Pure Chemicals Co., Japan. D-mannitol and aminoguanidine hemisulfate were obtained from Sigma-Aldrich

Chemical Co., St. Louis, MO, USA. Ham's F-12K media was used for hepatocytes/cell line culturing (Wako Pure Chemicals Co., Japan). LDH cytotoxicity detection kit (Takara Company, Tokyo, Japan), mitochondria extraction kit (Sigma-Aldrich Chemical Co., St. Louis, MO, USA.), Oxyblot protein oxidation detection kit (Chemicon

International Co., USA), Qproteome for detection of Apoptosis and In situ cell death detection kit (Roche Company, Germany).

b. Methods:

1- Cell culture, liver cells injury induction and treatment with different antioxidants:-

Liver cell line BRL3A derived from buffalo rats (*Nissley et al., 1977*) was used. Approximately 1×10^5 cells/ml (5 ml final volume) were cultured into a 25 ml flask and incubated under standard conditions (37 °C and 5 % CO₂) in Ham's F-12K medium with L-Glutamine, phenol red and sodium pyruvate supplemented with 10% Fetal Bovine Serum (FBS) and penicillin and streptomycin as antibiotics. Carbon tetrachloride (7.5 µl) was added in the 25 ml headspace of the flask to a paper attached to the stopper. During incubation, CCl₄ evaporated and equilibrium was reached between the gas phase and the medium (*Azri et al., 1990*). Media samples were collected at 30 min, 1, 3, 6, 9, 12, 14, 24 and 48 hrs. Different antioxidants (vitamin E, ascorbic acid, mannitol and aminoguanidine) were added separately to the media just before the addition of CCl₄. Each agent was individually tested on the cells using a dose range. For vitamin E this was 100, 50 and 25 µM; for ascorbic acid 500, 250 and 125 µM; for either mannitol or aminoguanidine the dose range was 400, 200 and 100 µM. Media samples were collected at 14, 24 and 48 hrs.

2- Detection of Free radicals by Electron Spin Resonance (ESR) method:-

Electron spin resonance signals were measured by preincubation of the hepatocytes in the presence of 40 mM 5, 5-dimethyl-pyrroline-N-oxide (DMPO) (*Matsuki, et al., 1999*). After induction of CCl₄ toxicity, media samples were collected after 1, 2, 4 and 6 hrs. A control group was incubated with DMPO only. All ESR spectra were recorded at room temperature. AJEOL Model JES – FA 100 ESR spectrometer (JEOL Co., Tokyo Japan) was used. Spectrometer settings were; magnetic field 335.00±5 mT, sweep time 8.0 min, modulation frequency 100 kHz, modulation amplitude 0.2 mT, receiver gain 8×10^3 , time constant 0.03 S, microwave frequency 9.418GHz and microwave power 12 mW.

3- Leakage of intracellular enzymes as (Lactate dehydrogenase (LDH) and aminotransferase (ALT and AST) leakage in cell culture medium: Leakage of LDH enzyme in the cultured media was measured by a kit while the activities of ALT and AST were determined by using automatic multifunction-biochemical analyzer (DRI-CHEM 5500, Japan).

4- Measurement of Lipid Peroxidation: It was carried out by measuring thiobarbituric acid – reactive species (TBARS) according to *Kikugawa et al., (1992)*.

5- Protein oxidation detection:

Using Qproteome mitochondria extraction kit, mitochondria was extracted from control and treated cells after 24 hours incubation with CCl₄ to assess the formation of protein carbonyl groups as an index of protein oxidation. Protein concentration was determined by the Bradford assay (Bio-Rad) according to Bradford (1976). An OxyBlot protein oxidation detection kit (Chemicon International Co., USA.) was used according to the manufacturer's detailed protocol. Subsequently, 5 µL protein sample was added with 5 µL of 12% SDS and 10 µL of 1× DNPH solution into a tube. Ten microliters of 1× neutralization solution instead of the DNPH solution served as the negative control. Tubes were incubated at room temperature for 15 minutes. Neutralization solution (7.5 µL) was added to each tube. The DNP-derivatized protein samples were separated by polyacrylamide gel electrophoresis followed by Western blotting. Following electrophoresis and transfer to nitrocellulose membranes, the membranes were blocked in Tris buffered saline containing 0.1% Tween 20 and 1% bovine serum albumin for 1 hour at room temperature. Membranes were incubated overnight at room temperature with the primary antibody stock (1:150) and then incubated with secondary antibodies (1:3000) at room temperature for 1 hour. Blots were developed by an enhanced chemiluminescence detection system. Proteins that underwent oxidative modification (i.e., carbonyl group formation) were identified as a band in the derivatized sample, but not in the negative control. Levels of oxidatively modified proteins were quantified and expressed via measurement of optical density using molecular analyst alias program.

6- Analysis of oxidative DNA strand breaks by TUNEL assay:-

The principle of the test is that cleavage of DNA during apoptosis may yield double stranded as well as single strand breaks "nicks" which can be identified by labeling with modified nucleotides in an enzymatic reaction (TUNEL staining, according to Gavrieli et al., 1992). Cells were grown on a small cover glass which was placed inside the flask. After induction of CCl₄ toxicity, cells were washed in PBS and fixed with 4 % paraformaldehyde buffered saline for 1 h at 15-25 °C. Permeabilisation of the cells was carried out by incubation with 0.2% Triton X for 2 min. on ice (2-8 °C). Positive control cells were made by treatment with DNase I for 10 minutes to break up the DNA.

TUNEL reaction mixture (50 μ l) was added on sample, for the negative control, 50 μ l of label solution was added for 60 min. at 37 °C in a humidified atmosphere in the dark. The samples were analyzed under a fluorescence microscope using an excitation wavelength in the range of 450-500 nm. Converter AP (50 μ l) was then added on sample, and incubated in a humidified chamber for 30 min. at 37 °C. Substrate solution (100 μ l) was added to the slides which were incubated for 10 min. at 15-25 °C in the dark. Cells were washed in PBS between each of these steps. The slides were counterstained with haematoxyline and eosin, examined under light microscope, and the apoptotic cells were counted in 10 random fields.

c. Statistical analyses:-

Data are expressed as mean values \pm SE. Student's t-test was used for observations. One-way analysis of variance (ANOVA) was used to assess significant differences among treated groups followed by Tukey's test. Statistical analyses were performed using GraphPad prism software (GraphPad, Inc., California USA). Differences were considered statistically significant when $p < 0.05$.

3. Results

I- Toxic effect of CcL₄ on BRL3A cell line:-

a- Generation of free radicals

The ESR signals were observed after 2 hours incubation of BRL3A cell line with CcL₄ and DMPO (Fig.1, A) reached its peak at 4 hr and began to decline at 6 hr (Fig.1, D&E).

b. Leakage of intracellular enzymes:-

Activities of LDH, ALT and AST enzymes were increased significantly ($P < 0.05$) at 14 and 48 hrs. (Table, 1).

c- Lipids peroxidation:-

Lipids peroxidation revealed no significant change between control and treated cells with CcL₄ (Table, 1).

d -Oxidation of proteins:-

The amount of protein carbonyls was markedly increased in cells treated with CcL₄ compared with that of normal control group (Fig. 2, B).

e -Oxidative DNA strand breaks:-

Examination of control cells by the fluorescent microscope revealed no signals, while CcL₄ treated cells showed fluorescent signals. The percent of apoptotic hepatocytes stained with the TUNEL method was significantly greater in the

CcL₄-treated cells (4.2%) than that in the control group (0.66%) (Fig. 3) (Photo, 1).

II- Effect of different antioxidants against CcL₄-induced hepatotoxicity

in BRL3A cell line:-

A-Leakage of intracellular enzymes:-

Comparing with the CcL₄ treated group, All the used doses of vitamin E significantly ($p < 0.05$) reduced leakage of LDH enzyme to the media. Ascorbic acid at a dose of 500 μ M was able to reduce LDH enzyme leakage to the media after 24 and 48 hrs. Mannitol failed to show reduction in leakage of LDH except at a dose of 200 μ M at 48 hr, while aminoguanidine failed to show reduction in leakage of LDH. As regards to leakage of ALT and AST enzymes to the media, vitamin E treatment (100, 50 and 25 μ M) significantly ($p < 0.05$) reduced enzymes' leakage. Ascorbic acid, mannitol and aminoguanidine at all doses were unable to reduce the leakage of ALT and AST enzymes to the culture media compared with CcL₄ treated group throughout the experiment (Tables, 2,3 and 4).

b- Oxidation of proteins:-

Treatment with vitamin E showed marked reduction in the amount of protein carbonyls with all doses. Similar results were observed in Ascorbic acid treated groups. No changes in protein carbonyls were observed in groups treated with different doses of mannitol and aminoguanidine, (Figs. 4A & 4B),

c- Oxidative DNA strand breaks:-

Percentage of apoptotic cells stained with the TUNEL method was markedly decreased in vitamin E (VE) 100 μ M treated group. The percentage of positive TUNEL cells was 0.85% in control group, 3.8 % in CcL₄ group, 1.79 % in VE 100 μ M group, 2.1 % in VE 50 μ M and 2.3 % in VE 25 μ M treated group (Fig. 3B). Ascorbic acid (AA) did not reduce the incidence of apoptosis in cells treated with CcL₄. Percentage of positive apoptotic cells were 2.8% in AA 500 μ M, 2.5% in AA 250 μ M, 2.8% in AA 125 μ M and 3.2% in AA 62.5 μ M treated group. Treatment with different doses of mannitol (M) did not show protective effect against CcL₄ induced apoptosis in BRL3A cells. The percentages of positive TUNEL cells were 2.8% in M 400 μ M, 2.6% in M 200 μ M and 2.5% in 100 μ M (Fig. 3C). Aminoguanidine failed to reduce the incidence of apoptosis occurred in BRL3A cells due to CcL₄ toxicity. The percentage of apoptotic cells were 2.66% in AG 400 μ M, 2.42% in AG 200 μ M and 2.44% in AG 100 μ M group (Fig. 5).

Table (1):- LDH, ALT and AST enzyme activities & TBARS values of BRL3A cell line treated with CcL₄ at different incubation times.

Incubation time		LDH (U/ml)	ALT (U/L)	AST (U/L)	TBARS (μmol/L)
30 min	Control	0.026±0.001	1.667±0.333	9.000±1.723	0.006±0.001
	CcL ₄	0.032±0.002	2.000±0.577	8.000±1.000	0.012±0.002
1 h	Control	0.030±0.003	1.333±0.333	10.000±1.000	0.005±0.003
	CcL ₄	0.025±0.001	1.678±0.333	10.000±1.000	0.014±0.006
3 hs	Control	0.041±0.001	1.333±0.333	10.330±1.528	0.005±0.001
	CcL ₄	0.042±0.002	1.770±0.667	13.670±1.528	0.005±0.003
6 hs	Control	0.044±0.000	3.333±0.882	6.670±0.333	0.007±0.002
	CcL ₄	0.043±0.000	4.330±0.881	8.660±1.667	0.004±0.001
9 hs	Control	0.040 ±0.001	4.000±0.577	6.790±0.333	0.016±0.001
	CcL ₄	0.041±0.001	4.333±0.882	8.560±0.577	0.026±0.007
12 hs	Control	0.039±0.001	6.000±0.577	3.667±0.333	0.017±0.001
	CcL ₄	0.042±0.002	7.333±0.882	4.667±0.067	0.025±0.004
14 hs	Control	0.006±0.002	3.000±0.577	5.333±0.333	0.018±0.002
	CcL ₄	0.023±0.001**	7.335±0.333 **	.667±0.666*	0.024±0.006
24 hs	Control	0.013±0.003	4.333±0.333	6.300±0.330	0.022±0.003
	CcL ₄	0.028±0.001*	9.320±0.881**	11.67±0.333**	0.022±0.002
48 hs	Control	0.126±0.002	6.000±0.577	11.33±0.577	0.022±0.000
	CcL ₄	0.378±0.008**	12.67±1.453*	23.67±2.082**	0.031±0.004

(*) Significantly different from the control group at P < 0.05.

(**) Significantly different from the control group at P < 0.001.

Table (2): Activities of LDH, ALT and AST enzymes of BRL3A cell line treated with different doses of vitamin E, ascorbic acid, mannitol and aminoguanidine for 14 hr against toxicity of CcL₄.

Antioxidant	Groups	LDH (U/mL)	ALT (U/L)	AST (U/L)
Vitamin E (VE)	Control (0.5% ethanol)	0.005±0.002	4.33±0.333	3.670±0.333
	CcL ₄	0.077±0.003 ^a	9.67±0.333 ^a	11.330±1.202 ^a
	VE 100 μM+ CcL ₄	0.066±0.001 ^b	5.33±0.333 ^b	4.330±1.453 ^b
	VE 50 μM+ CcL ₄	0.068±0.000 ^b	6.33±0.333 ^b	4.667±1.202 ^b
	VE 25 μM+ CcL ₄	0.062±0.000 ^b	7.00±0.577 ^b	3.000±1.155 ^b
Ascorbic acid (AA)	Control	0.028±0.000	4.00±0.577	3.67±0.333
	CcL ₄	0.036±0.001 ^a	9.00±0.577 ^a	8.00±0.577 ^a
	AA 500 μM+ CcL ₄	0.033±0.001	7.67±0.333	6.00±0.577
	AA 250 μM+ CcL ₄	0.036±0.000	7.33±0.333	6.33±0.881
	AA 125 μM+ CcL ₄	0.034±0.001	7.00±0.577	7.00±0.577
Mannitol (M)	Control	0.009±0.000	3.33±0.667	4.33±0.333
	CcL ₄	0.030±0.002 ^a	7.33±0.333 ^a	9.00±0.577 ^a
	M 400 μM+ CcL ₄	0.049±0.002	6.67±0.667	9.67±0.333
	M 200 μM+ CcL ₄	0.041±0.003	6.00±0.577	9.33±0.333
	M 100 μM+ CcL ₄	0.039±0.000	5.33±0.333	8.33±0.333
Aminoguanidine (AG)	Control	0.009±0.000	3.33±0.667	3.67±0.333
	CcL ₄	0.022±0.000 ^a	7.33±0.330 ^a	10.67±0.333 ^a
	AG 400 μM+ CcL ₄	0.039±0.001	7.34±0.882	10.33±0.333
	AG 200 μM+ CcL ₄	0.028±0.002	6.33±0.660	9.67±0.333
	AG 100 μM+ CcL ₄	0.027±0.002	5.67±0.670	10.00±0.577

^(a) Significantly different from the control group at P<0.05.^(b) Significantly different from the CcL₄ treated group at P<0.05

Table (3):- Activities of LDH, ALT and AST enzymes of BRL3A cell line treated with different doses of vitamin E, ascorbic acid, mannitol and aminoguanidine for 24 hr against toxicity of CcL₄.

Antioxidant		groups	LDH (U/ml)	ALT (U/L)	AST (U/L)
Vitamin E (VE)		Control (0.5% ethanol)	0.004±0.000	8.000±0.577	7.670±0.333
		CcL₄	0.045±0.002 _a	15.330±0.882 _a	14.660±0.333 _a
		VE 100 μM+CcL₄	0.007±0.000 _b	6.330±0.881 _b	8.660±0.333 _b
		VE 50 μM+CcL₄	0.017±0.001 _b	7.000±1.13 _b	9.000±0.577 _b
		VE 25 μM+CcL₄	0.020±0.000 _b	10.000±0.577 _b	9.330±0.333 _b
Ascorbic acid (AA)		Control	0.029±0.000	5.00±0.577	4.67±0.333
		CcL₄	0.041±0.001 _a	10.33±0.333 _a	10.00±0.577 _a
		AA 500 μM+CcL₄	0.032±0.001 _b	10.00±0.577	9.00±0.577
		AA 250 μM+CcL₄	0.042±0.000	12.00±0.577	9.00±0.577
		AA 125 μM+CcL₄	0.042±0.001	9.00±0.577	9.33±0.333
Mannitol (M)		Control	0.027±0.001	4.00±0.577	5.67±0.333
		CcL₄	0.040±0.000 _a	9.00±0.577 _a	12.33±0.333 _a
		M 400 μM+CcL₄	0.073±0.000	7.33±0.33	11.30±0.667
		M 200 μM+CcL₄	0.069±0.004	7.33±0.33	11.33±0.333
		M 100 μM+CcL₄	0.047±0.001	9.33±0.33	10.67±0.333
Aminoguanidine (AG)		Control	0.027±0.001	4.00±0.577	5.00±0.577
		CcL₄	0.04±0.002 _a	9.00±0.577 _a	12.33±0.333 _a
		AG 400 μM+CcL₄	0.052±0.001	8.33±0.577	10.67±0.333
		AG 200 μM+CcL₄	0.045±0.000	8.00±1.15	11.33±0.333
		AG 100 μM+CcL₄	0.038±0.001	8.00±0.000	11.67±0.333

^(a) Significantly different from the control group at P<0.05.

^(b) Significantly different from the CcL₄ treated group at P<0.05

Table (4):- Activities of LDH, ALT and AST enzymes of BRL3A cell line treated with different doses of vitamin E, ascorbic acid, mannitol and aminoguanidine for 48 hr against toxicity of CCL₄.

Antioxidant	Groups	LDH (U/mL)	ALT (U/L)	AST (U/L)
Vitamin E (VE)	Control (0.5% ethanol)	0.004±0.00	6.000±0.577	18.000±0.577
	CcL ₄	0.031±0.001 _a	11.000±0.577 _a	25.000±0.577 _a
	VE 100 µM+ CcL ₄	0.004±0.000 _b	7.000±0.577 _b	12.000±0.577 _b
	VE 50 µM+ CcL ₄	0.012±0.000 _b	5.000±0.577 _b	14.000±0.577 _b
	VE 25 µM+ CcL ₄	0.014±0.000 _b	7.000±0.577 _b	13.000±0.577 _b
Ascorbic acid (AA)	Control	0.047±0.001	4.00±0.577	10.33±0.333
	CcL ₄	0.068±0.002 _a	8.00±0.577 _a	22.67±0.333 _a
	AA 500 µM+ CcL ₄	0.058±0.002 _b	8.00±0.577	20.00±0.577
	AA 250 µM +CcL ₄	0.070±0.001	7.00±0.577	21.33±0.333
	AA 125 µM+ CcL ₄	0.068±0.002	6.00±0.577	21.33±0.882
Mannitol (M)	Control	0.018±0.003	5.00±0.577	12.00±0.577
	CcL ₄	0.031±0.004 _a	8.00±0.577 _a	23.00±0.577 _a
	M 400 µM +CcL ₄	0.041±0.001	7.33±0.330	21.00±0.577
	M 200 µM +CcL ₄	0.045±0.001	5.00±0.577 _b	21.00±0.577
	M 100 µM +CcL ₄	0.048±0.001	6.00±0.577	20.67±0.333
Aminoguanidine (AG)	Control	0.017±0.003	5.00±0.577	9.00±0.577
	CcL ₄	0.030±0.004 _a	8.00±0.577 _a	22.00±0.577 _a
	AG 400 µM +CcL ₄	0.046±0.002	6.67±0.882	20.33±0.333
	AG 200 µM +CcL ₄	0.034±0.001	6.00±0.577	20.30±0.333
	AG 100 µM +CcL ₄	0.027±0.000	5.33±0.33	19.67±0.333

^(a) Significantly different from the control group at P<0.05.

^(b) Significantly different from the CcL₄ treated group at P<0.05




Fig. (1):- ESR spectra of supernatants of BRL3A cell line treated with CCL4 at different periods of incubation.

(Spectrum A) ESR spectra of the 40 mM DMPO adduct detected in BRL3A cell line. (Spectrum B) Same as in spectrum A, but cells were incubated with CCl₄ for one hour. The spectrum shown in C is from cells incubated 2 hours with CCl₄, while the spectrum in D after 4 hours incubation with CCl₄. Spectrum shown in E after 6 hour incubation period with CCl₄. Instrumental settings of a JEOL Model JES-FA 100 ESR spectrometer: magnetic field 335.000±5 mT, sweep time 8.0 min, modulation frequency 100 kHz, modulation amplitude 0.2 mT, receiver gain 8x10³, time constant 0.03 S, microwave frequency 9.418GHz and microwave power 12 mW.




Fig. (2,A):- Oxyblot of mitochondria of BRL3A in both control and CCl₄ treated groups using oxyblot technique.

Fig. (2,B):-Densitometric measurement of the oxyblot lanes of fig. (2,A)




Fig. (3):- Quantitative evaluation of TUNEL positive cells in BRL3A cells treated with CCl₄.




Photo (1):- Cell apoptosis of BRL3A cell line after exposure to CCl₄ as detected with TUNEL assay under fluorescent microscope.




Fig. (4,A):- Oxyblot of mitochondria of BRL3A incubated with CCl₄ and different dose of vitamin E (VE), ascorbic acid (AA), Mannitol (M) an aminoguanidine (AG).




Fig. (4,B):- Densitometric measurement of the oxyblot lanes of Fig. (3,A)




Fig (5):- Quantitative evaluation of TUNEL positive cells in BRL3A cells treated with ccl4 and different doses of vitamin E (VE), ascorbic acid (AA), Mannitol (M) and aminoguanidine (AG).

4. Discussion

Various pharmacological or chemical substances are known to cause hepatic injury. It is believed that the occurrence of drug/chemicals-induced liver injury is especially associated with oxidative stress and a cellular imbalance between the production and elimination of free radicals (Castro and Freeman, 2001). Overproduction of free radicals could directly injure hepatocellular membrane by lipid peroxidation, or other means, followed by a series of cascades of cellular events such as massive release of inflammatory mediators or cytokines,

which eventually lead to liver injury (Higuchi and Gores, 2003). Therefore, it is valuable to identify natural drugs or compounds that can antagonize the deleterious action of free radicals and act as an antioxidant to protect hepatocytes from damage.

In the present experiments, hepatocytes cultured cells were utilized. In an *in vitro* system, compounds affect the cells directly and continuously until the removal of compound-containing medium. In addition, when cells are cultured using single type of cells, there are no interactions from other interstitial cells (Kikkawa et al., 2006).

Cytotoxicity of CCL_4 has been selected as a model for induction of oxidative damage. Carbon tetrachloride is one of the most extensively studied hepatotoxicants (*Bhattacharjee and Sil, 2007*). It is now generally believed that CCL_4 hepatotoxicity depends on its reductive dehalogenation catalyzed by cytochrome P-450 enzyme in the endoplasmic reticulum of hepatic cells leading to the generation of an unstable complex trichloromethyl radical. The superoxide anion, H_2O_2 , and the hydroxyl radical (OH) are reactive oxygen species (ROS) mainly produced in mitochondria (*Wang et al., 2007*).

The possible protective effect of some antioxidants (ascorbic acid, mannitol and aminoguanidine) has been attempted on hepatocytes cultured cells treated with CCL_4 . Trials were compared to vitamin E known to be important lipophilic antioxidant in biological tissues. The activity of an antioxidant may depend on its reactivity towards particular radicals, its ability to concentrate near the critical target in the cell or its inhibitory action on radical formation (*Anderson and Phillips, 1999*).

Electron spin resonance (ESR) was employed to measure directly reactive oxygen species (ROS) production in the liver of CCL_4 -treated hepatocytes cell line. Results indicated that ROS production was highly elevated in the cultured media of the cell line at 4 h after CCL_4 treatment, thus further confirming that free radicals and oxidative damage certainly play a vital role in the pathogenesis of acute liver injury (*Wu et al., 2007*).

Cell viability depends directly on the structure of the membrane, and damage to the cellular membrane can be detected by enzyme leakage. It has been generally accepted that leakage of the cytosolic enzyme LDH correlates well with cellular viability, thus being a useful indicator of plasmatic membrane damage (*Grajeda-Cota et al., 2004*). In the present study, observations revealed that CCL_4 induced significant increase in the leakage of LDH, ALT and AST enzymes into the medium after 14 hr. The obtained results are in harmony with those reported by *Chandan et al. (2007)*. It is interesting to note that there was difference in the degree of leakage of ALT and AST enzymes, where AST enzyme leakage was higher than ALT enzyme leakage. *McQueen and Williams (1982)* reported differences in the extent of leakage of enzymes from the cytosolic fraction of hepatocytes. These differences could be due to differences in the tightness of binding of cytosolic enzymes to cellular organelles and in the size of the enzyme molecule (*Nakamura et al., 1985*). Moreover, AST is present in two isozymes, one located in the cytoplasm and the other in the mitochondria (*Latner,*

1975). The presence of enzymes outside the cell represents damage to the hepatic cell.

Involvement of lipid peroxidation in the mechanism of carbon tetrachloride-induced hepatotoxicity has been a point of controversy. The present results denoted that CCL_4 did not induce lipid peroxidation as there was no significant difference in the level of thiobarbituric acid-reactive substances (TBARS) between CCL_4 treated cells and control cells. These data are comparable to that of *Ikeda et al. (1998)* who reported that TBARS, which are widely used index of lipid peroxidation, don't increased significantly in the liver of rats treated with CCL_4 . Previous investigators reported absence of lipid peroxidative degradation products in mice after exposure to carbon tetrachloride and have used this evidence against the hypothesis that lipid peroxidation is an integral part of the events that cause tissue damage (*Lee et al., 1982*). On the contrary, *Krithika et al. (2009)* recently proved the involvement of lipid peroxidation on exposure to CCL_4 .

As regards to oxidation of proteins, the present study was able to detect that there was a significant increase in the formation of carbonyl proteins in the CCL_4 treated cells comparing to control cells. Free radical-mediated oxidation of proteins results in the formation of carbonyl groups in quantities that reflect the intensity of the oxidative stress (*Robinson et al.; 1999*). Protein carbonyl content is widely used as both a marker for oxidative stress and a measure of oxidative damage (*Luo and Wehr, 2009*). Carbonyl (CO) groups (aldehydes and ketones) are produced on protein side chains when they are oxidized. These moieties are chemically stable, which is useful for both their detection and storage (*Dalle-Donnea et al., 2003*). Protein oxidation may play a role in the pathogenesis of CCL_4 induced liver injury and that the accumulation of oxidized proteins may be an early indication of CCL_4 induced liver damage (*Sundari et al. 1997*). The usage of protein (CO) groups as a marker may have some advantages in comparison with lipid peroxidation products because the formation of protein bound (CO) groups seems to be a common phenomenon of protein oxidation and because of the relatively early formation and stability of oxidized proteins (*Dalle-Donnea et al., 2003*).

Apoptosis is characterized by morphological changes such as membrane "blebbing", nuclear condensation and fragmentation (*Ramage, et al., 2006*). These morphological changes are a result of a cascade of biochemical changes occurring within the cell, resulting in activation of caspases and DNA fragmentation (*Hengartner, 2000*). The present data indicate that along with oxidative damage, apoptosis

plays a crucial role in CcL₄ induced hepatotoxicity which was indicated by marked increase in DNA fragmentation (increase in the percent of TUNEL positive cells) in the CcL₄ treated cells compared to control one. Apoptosis represents one of the mechanisms of cell death after CcL₄ induced liver injury (Shi et al, 1998).

Effects of antioxidants on oxidative damage in vitro systems are variable. In most cases, endogenous antioxidant enzymes, particularly catalase, are very effective in preventing damage. Exogenous antioxidant chemicals, on the other hand, can have a pro-oxidant effect at high doses. The chemical composition of culture media and the doses of antioxidants that can be applied in culture systems are major factors in confounding interpretation of in vitro results (Anderson and Phillips, 1999).

Vitamin E antioxidant functions mainly in and around the membrane/lipid bilayers acting as the chain breaking antioxidants (Janero and Burghardt, 1989). Results of the present experiments showed that incubation of hepatocytes with vitamin E at concentrations of 100, 50 and 25 μM induced hepatoprotective effect against CcL₄ induced toxicity. This was manifested by an increase in the stability of the cell membrane and decreased leakage of cytoplasmic enzymes (LDH, ALT and AST). Although we cannot rule out a direct scavenging action of vitamin E on CcL₃ radical, inhibition of lipid peroxidation is not a likely explanation for vitamin E effects in the present study. Although the ability of CcL₄ to initiate lipid peroxidation in various biological systems has been equivocal, an important observation of the present study was that there was no evidence of CcL₄-initiated lipid peroxidation.

Ascorbic acid, in the present study, was ineffective against the elevation of enzymes leakage (LDH, ALT and AST). Also it was not able at any concentration to reduce the oxidation of protein. Ascorbic acid did not show any protective effect against DNA damage caused by CcL₄. Similar results were obtained by Winter et al. (2005) who reported that ascorbic acid was ineffective at reducing oxidative DNA damage produced by camphorquinone which generates reactive oxygen species and causes oxidative DNA damage in vitro. On the contrary Qin et al. (2006) found that ascorbic acid reduced myocyte apoptosis and ameliorated myocardial damage after acute myocardial infarction. Park and Lee (2008) reported that high ascorbic acid concentrations might act as a pro-oxidant due to its auto-oxidizing properties.

Mannitol is generally accepted as being a hydroxyl radical scavenger (Van Zandwijk, 1995). Previous studies have used mannitol to determine whether hydroxyl radicals are the predominant

species responsible for Copper mediated DNA damage (Li et al., 1995). However, these studies have found that mannitol is generally ineffective at preventing Cu-mediated oxidative damage in the presence of an ROS-generating system and at mannitol concentrations less than 100mM. The present results showed that mannitol did not effectively reduce CcL₄ induced hepatotoxicity. It showed no reduction in the leakage of cytoplasmic enzymes, protein oxidation or DNA damage. Winter et al. (2005) reported that mannitol treatments of 10.0, 5.0, 2.5, 1.0, and 0.5 mM did not influence the extent of DNA damage generated by camphorquinone. In explanation, it is possible that some ROS generated from CcL₄ are not effectively scavenged by mannitol. Additionally, mannitol may not be a very potent hydroxyl radical scavenger or mannitol may not have adequate access to the damaging ROS (Winter et al., 2005). Shinar et al. (1983) demonstrated that mannitol effectively scavenged hydroxyl radicals at concentrations greater than 1M.

In the present experiments, aminoguanidine (AG) treatment against CcL₄ induced hepatotoxicity was ineffective in reducing the leakage of enzymes, the oxidation of protein, and the DNA damage. A notable finding was that AG at a dose of 400 μM significantly increased leakage of LDH into the media comparing with CcL₄ group. Our results confirmed reports of other investigators (Ou and Wolff, 1993 and Skamarauskas et al., 1996) that AG has pro-oxidant activity where it generates H₂O₂ at a low rate in vitro.

In conclusion, the present study indicates that CcL₄ has a potential cytotoxic effect in BRL3A cell culture, and apoptosis represents one of the mechanisms of cell death after CcL₄-induced liver injury. There was no evidence of CcL₄-initiated lipid peroxidation. Vitamin E seems effective and provides complete protection against CcL₄ induced hepatotoxicity in culture cells. Ascorbic acid, mannitol and aminoguanidine were ineffective against CcL₄ toxicity. Aminoguanidine, in addition, generates H₂O₂ in vitro, so it has pro-oxidant activity.

Corresponding author

Azza, H. Abd-El-Rahman¹

Clinical Pathology Dept., Fac. of Vet. Med. Beni-Suef University, Beni-Suef, Egypt

5. References:

1. Alpsoy, L; Yildirim, A and Agar, G. (2009):- The antioxidant effects of vitamin A, C and E on aflatoxin B1-induced oxidative stress in human lymphocytes.

- Toxicology and Industrial Health, 25: 121-127.
2. Anderson D. and Phillips, B. (1999):- Comparative in vitro and In vivo effects of antioxidants. Food and Chemical Toxicology, 37: 1015-1025.
 3. Azri, S., Gandolfi, A. J., and Brendel, K. (1990):- Carbon tetrachloride toxicity in precision-cut rat liver slices. In Vitro Toxicology, 3: 127-138.
 4. Bhattacharjee, R. and Sil, P. (2007):- Protein isolate from the herb phyllanthus niruri modulates carbon tetrachloride-induced cytotoxicity in hepatocytes. Toxicology Mechanisms and Methods, 17: 41-47.
 5. Bradford, M. (1976):- A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. Analytical Biochemistry, 72: 248-254.
 6. Castro, L. and Freeman, B. (2001):- Reactive oxygen species in human health and disease. Nutrition, 17: 161-165.
 7. Chandan, B.; Saxena, A.; Shukla, S.; Sharma, N.; Gupta, D.; Suri, K. Suri, J.; Bhadauria, M. and Singh, B. (2007):- Hepatoprotective potential of Aloe barbadensis Mill against carbon tetrachloride induced hepatotoxicity. Journal of Ethnopharmacology, 111(3): 560-6.
 8. Dalle-Donnea, I.; Rossib, R.; Giustarinib, D. Milzania, A. and Colombo, R. (2003):- Review: Protein carbonyl groups as biomarkers of oxidative stress. Clinica Chimica Acta, 329: 23-38.
 9. Gavrieli, Y.; Sherman, Y. and Ben-Sasson S. (1992):- Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. The Journal of Cell Biology, 119: 493-501.
 10. Grajeda-Cota, P.; Ramírez-Mares, M. V. and González de Mejía, E. (2004):- Vitamin C protects against in vitro cytotoxicity of cypermethrin in rat hepatocytes. Toxicology in Vitro, 18: 13-19.
 11. Guillouzo, A. (1998):- "Liver cell models in in vitro toxicology." Environmental Health Perspectives, 106(2): 511-32.
 12. Hengartner, M.O. (2000):- The biochemistry of apoptosis. Nature, 407: 770-776.
 13. Higuchi, H. and Gores, G. (2003):- Mechanisms of liver injury: an overview. Current Molecular Medicine, 3: 483-490.
 14. Ikeda, K.; Toda, M.; Tanaka, K.; Tokumar, S. and Kojo, S. (1998):- Increase of lipid hydroperoxides in liver mitochondria and inhibition of cytochrome oxidase by carbon tetrachloride intoxication in rats. Free Radical Research, 28: 403-410.
 15. Janero, D. and Burghardt, B. (1989):- Cardiac membrane vitamin E and malondialdehyde levels in heart muscle of normotensive and spontaneously hypertensive rats. Lipids, 24: 33-38.
 16. Khan, T. and Sultana, S. (2009):- Antioxidant and hepatoprotective potential of Aegle marmelos Correa against CCl4-induced oxidative stress and early tumor events. Journal of Enzyme Inhibition and Medicinal Chemistry, 24 (2): 320-7.
 17. Kikugawa, K.; Kojima, T.; Yamaki, S. and Kosugi, H. (1992):- Interpretation of the thiobarbituric acid reactivity of rat liver and brain homogenates in the presence of ferric ion and ethylenediaminetetraacetic acid. Analytic biochemistry 202, 249-255.
 18. Kikkawa, R.; Fujikawa, M.; Yamamoto, T.; Hamada, Y.; Yamada, H. and Horii, I. (2006):- In vivo hepatotoxicity study of rats in comparison with in vitro hepatotoxicity screening system. The Journal of Toxicological Sciences, 31 (1): 23-34.
 19. Krithika, R.; Mohankumar, R.; Verma, R.; Shrivastav, P.; Mohamad, I.; Gunasekaran, P. and Narasimhan, S. (2009):- Isolation, characterization and antioxidative effect of phyllanthin against Ccl4- induced toxicity in HepG2 cell line. Chemico- Biological Interactions, 181(3): 351-8.
 20. Kucera, O.; Lotková, H.; Kriváková, P.; Rousar, T. and Cervinková, Z. (2006):- Model systems for study of toxic injury of hepatocytes in vitro.
 21. Ceskoslovenska Fysiologie, 55 (3): 103-10.
 22. Latner, A.L. (1975):- Enzymes. In: Saunders, W.B. Company (Eds.), Clinical Biochemistry Cap, 17: 547-549, Philadelphia, London, Toronto.
 23. Lee, P.; McCay, P. and Hornbrook, K. (1982):- Evidence for carbon tetrachloride-induced lipid peroxidation in mouse liver. Biochemical Pharmacology, 31(3): 405-9.
 24. Li, Y.; Kuppusamy, P.; Zweier, J. and Trush, M. (1995):- ESR evidence for the generation of reactive oxygen species from the copper-mediated oxidation of the benzene metabolite hydroquinone: role in DNA damage. Chemico-Biological Interactions, 94:101-20.
 25. Luo, S. and Wehr, N. (2009):- Protein carbonylation: avoiding pitfalls in the 2,4-

- dinitrophenylhydrazine assay. Redox Report, 14(4):159-66.
26. Manibusan M.; Odin, M. and Eastmond, D. (2007): Postulated carbon tetrachloride mode of action: a review. Journal of Environmental Sciences And Health part C Environmental Carcinogenesis and Ecotoxicology Reviews, 25 (3): 185-209.
 27. Matsuki, N.; Takanohashi, A.; Boffi, F.; Inanami, O.; Kuwabara, M. and Ono, K. (1999):- Hydroxyl radical generation and lipid peroxidation in C2C12 myotube treated with iodoacetate and cyanide. Free Radical Research, 31 (1): 1-8.
 28. McQueen, C. and Williams, G. (1982):- Cytotoxicity of xenobiotics in adult rat hepatocytes in primary culture. Fundamental and Applied Toxicology, 2: 139-144.
 29. Mendoza, G.; Alvarez A.; Pulido, M.; Molina, A.; Merino, G.; Real, R. Fernandes, P. and Prieto, J. (2007):- Inhibitory effects of different antioxidants on hyaluronan depolymerization. Carbohydrate Research, 342 (1): 96-102.
 30. Mohamad, N. A.; Cricco, G. P.; Sambuco, L. A, Croci, M.; Medina, V. A.; Gutiérrez, A. S.; Bergoc, R. M.; Rivera, E. S. and Martín, G. A. (2009):-
 31. Aminoguanidine impedes human pancreatic tumor growth and metastasis development in nude mice. World Journal of Gastroenterology, 15 (9): 1065–1071.
 32. Nakamura, T.; Fujii, T. and Ichihara, A. (1985):- Enzyme leakage due to change of membrane permeability of primary cultured rat hepatocytes treated with various hepatotoxins and its prevention by glycyrrhizin. Cell Biology and Toxicology, 1(4): 285-95.
 33. Nissley, S .P., Short, P .A., Rechler, M .M., Podskalny, J .M., and Coon, H .G .(1977):- Proliferation of buffalo rat liver cells in serum-free medium does not depend upon multiplication-stimulating activity (MSA). Cell, 11: 441–446.
 34. Ou, P. and Wolff, S. (1993):- Aminoguanidine :a drug proposed for prophylaxis in diabetes inhibits catalase and generates hydrogen peroxide in vitro. Biochemical Pharmacology, 46: 1139–44.
 35. Park, S.W. and Lee, S. M. (2008):- Antioxidant and prooxidant properties of ascorbic acid on hepatic dysfunction induced by cold ischemia/reperfusion. European Journal of Pharmacology, 580: 401–406.
 36. Qin, F.; Yan, C.; Patel, R.; Liu, W. and Dong, E. (2006):- Vitamins C and E attenuate apoptosis, -adrenergic receptor desensitization and sarcoplasmic reticular Ca²⁺ ATPase downregulation after myocardial infarction. Free Radical Biology and Medicine, 40: 1827–1842.
 37. Ramage, L.; Jones, A. and Whelan, C. (2006):- Induction of apoptosis with tobacco smoke and related products in A549 lung epithelial cells in vitro. Journal of Inflammation, 3: 3-14.
 38. Robinson, C.; Keshayarzian, A.; Pasco, D.; Frommel, T.; Winship, D. and Holmes, E. (1999):- Determination of protein carbonyl groups by immunoblotting. Analytical Biochemistry, 266(1): 48-57.
 39. Ruijter, G.; Bax, M.; Patel, H.; Flitter, S.; Van de vondervoort, P.; De vries, R.; Vankuyk, P. and Visser, J. (2003):- Mannitol is required for stress tolerance in Aspergillus niger conidiospores. Eukaryotic Cell, 2 (4): 690-698.
 40. Shi, J.; Aisaki, K.; Ikawa, Y. and Wake, K. (1998):- Evidence of hepatocyte apoptosis in rat liver after the administration of carbon tetrachloride. The American Journal of Pathology, 153 (2): 515-525.
 41. Shinar, E.; Navok, T. and Chevion M. (1983):- The analogous mechanism of enzymatic inactivation induced by ascorbate and superoxide in the presence of copper. The Journal Of Biological Chemistry 258: 14778–83.
 42. Skamarauskas, J.; McKay, A. and Hunt, J. (1996):- Aminoguanidine and its pro-oxidant effects on an experimental model of protein glycation. Free Radical Biology and Medicine, 21: 801–812.
 43. Södergren, E.; Cederberg, J.; Vessby, B. and Basu, S . (2001):- Vitamin E reduces lipid peroxidation in experimental hepatotoxicity in rats. European Journal of Nutrition, 40 (1):10-6.
 44. Sundari, P.; Wilfred, G. and Ramakrishna B. (1997):- Does oxidative protein damage play a role in the pathogenesis of carbon tetrachloride-induced liver injury in the rat? Biochimica et Biophysica Acta, 1362(2-3):169-76.
 45. Van Zandwijk N. (1995):- N-acetylcysteine (NAC) and glutathione (GSH): antioxidant and chemopreventive properties, with special reference to lung cancer. Journal of

- Cellular Biochemistry Supplement, 22: 24–32.
46. Wang, C. Y.; Ma, F. L.; Liu, J. T.; Tian, J. W. and Fu, F. H. (2007):-Protective Effect of Salvianic Acid A on Acute Liver Injury Induced by Carbon Tetrachloride in Rats." *Biological And Pharmaceutical Bulletin*, 30 (1): 44-47.
 47. Weber, L.W., Boll, M.and Stampfl, A. (2003):- Hepatotoxicity and mechanism of action of haloalkanes: carbon tetrachloride as a toxicological model. *Critical Reviews Toxicology*, 33 (2): 105-36.
 48. Winter, K.; Pagoria, D. and Geurtsen, W. (2005):- The effect of antioxidants on oxidative DNA damage induced by visible-light-irradiated camphorquinone/N,N dimethyl-p-toluidine. *Biomaterials*, 26: 5321–5329
 49. *Wu, Y.; Li, L.; Wen, T .and Li, Y .Q . (2007):- Protective effects of echinacoside on carbon tetrachloride-induced hepatotoxicity in rats. *Toxicology*, 232 (1-2): 50–56.

2010/23/11