

Toxigenic fungi and aflatoxin associated to nuts in Saudi Arabia

Mohamed Deabes¹ and Roquia Al- Habib²

Food Toxicology and Contaminants Department, National Research Centre, Dokki, Giza, Egypt¹ and Department of Biology, College of Arts and Science, Buraydah, Qassim University, king of Saudi Arabia².

Corresponding author: mydeabes@yahoo.com

Abstract: A survey was carried out in July-Sept., 2009 to obtain data on the occurrence of aflatoxin and the aflatoxins-producing potential of fungi isolated from nuts (almonds, peanuts, hazelnuts, pistachio nuts, Walnut and Cashew) in region of Qassim in Saudi Arabia. The samples were analyzed for aflatoxins by immune affinity column (IAC) clean-up with liquid chromatography and fluorescence detection. Percentages of positive samples with aflatoxins were 80, 80, 60, 40, 40, 20 % for pistachio, peanuts, walnuts, almonds, hazelnuts and cashews. Concentrations of aflatoxin B₁ were ranged between (38- 45, 11- 90, 41 -90, 0.3-3.6, 62-120 and 70-140 "µg/kg") respectively. 100% of samples showed variable incidence of fungal contamination. Fungi present in samples were *Penicillium spp*, *Aspergillus niger*, *A. flavus* and *Rhizopus spp*. Results showed that isolates of *A. flavus* were able to produce aflatoxins B₁, B₂, G₁, and G₂. The purpose of the survey was to determine levels of aflatoxins and to monitor the effectiveness of the controls in place to limit consumer exposure to aflatoxins.

[Mohamed Deabes and Roquia Al-Habib. Toxigenic fungi and aflatoxin associated to nuts in Saudi Arabia. Journal of American Science 2011;7(8):658-665]. (ISSN: 1545-1003). <http://www.americanscience.org>.

Keywords: exposure to carcinogens, aflatoxins, nuts, HPLC, Fungi, food contaminants.

1. Introduction

Mycotoxins; toxic secondary metabolites of fungi are biological in origin. Despite efforts to control fungal contamination, toxigenic fungi are ubiquitous in nature and occur regularly in worldwide food supplies due to mold infestation of susceptible agricultural products, such as cereal grains, nuts, and fruits. Thousands of mycotoxins exist, but only a few present significant food safety challenges. The natural fungal flora associated with foods is dominated by three genera of fungi; *Aspergillus*, *Fusarium*, and *Penicillium*, which except for the *Fusarium* plant pathogens, may include commensals as well as pathogens. Groundnut is frequently infected with fungi that produce mycotoxins during and after harvesting, which affect the quality and safety of human food (Martin *et al.*, 1999). They are toxic to humans and animals, cause significant reductions in crop yield and cause economic losses (Gourama and Bullerman, 1995; Gqaleni *et al.*, 1996). Their occurrence in various countries has been well documented (Bathnagar and Garcia, 2001). *Aspergillus flavus* and *Aspergillus parasiticus* are important contaminants of certain foods and animal feeds because of their ability to produce aflatoxins (Farr *et al.*, 1989). When these fungi invade and grow in commodities such as peanuts, corn and cottonseed, the resulting contamination with aflatoxins often makes the commodities unfit for consumption (Vardon, 2003). Aflatoxins are considered the most carcinogenic, mutagenic and teratogenic substances found naturally in foods and feeds (Conner, 1993). These metabolites cause liver damage to humans and

to most experimental animal species tested (Gradelet *et al.*, 1997). Consumption of mycotoxin-contaminated foods has been associated with several cases of human poisoning, or mycotoxicosis, sometimes resulting in death (Bathnagar and Garcia, 2001). The International Agency for Research on Cancer (IARC) has classified aflatoxin B₁ (AFB₁) as a group 1 carcinogen, primarily affecting liver (IARC, 1993). Natural occurrence of AF in nuts has been studied in various countries. According to a report from Mexico, 2.2% of pistachio nut samples analyzed contained AF higher than 20 ng/g (JECFA, 1998). In Sweden, 9.5% pistachio nut samples contained AFB₁ higher than 2 ng/g (Thuvander *et al.*, 2001). According to Ministry of Agriculture and Rural Affairs, Republic of Turkey, (2002) analysis of 523 pistachio nut samples in Turkey the mean of AFB₁ ranged 1–3.78 ng/g and the maximum level detected was 113 ng/g.

The present study was undertaken to determine the occurrence of fungi, can production of aflatoxins in nuts in Saudi Arabia.

2. Material and Methods

Methods:

Samples:

Thirty samples of nuts (almonds, hazelnuts, pistachio nuts, Cashew, Walnut and peanuts) were randomly collected from Saudi Arabia, Qassim region (Buraydah, Unayzah, Al-Ras, Al- Badaye and Al- Bukayriyah) during Juli to Sept at 2009.

Isolation of fungi associated with nuts.

Fungi associated with nuts were isolated according to methods of Lichtwardt *et al.*, (1958), Mislivec (1977) and Ichinoe *et al.* (1983). All nut samples were immersed in sodium hypochlorite solution (5%) as a sterilizer, for 3 minutes, rinsed 3 times in sterile distilled water then dried between sterile filter paper. Isolation of fungi was made by randomly taking 100 disinfested grains from samples of each location and directly plated in 20 Petri-dishes (5 grains per dish) on Czapek's agar medium, then incubated at 25°C for 4 days. Then, the percentage of the infected grains was determined and the fungal colonies that developed from the infected grains were counted, isolated, purified and maintained on a slant potato dextrose agar (PDA) medium for identification trials.

Identification of the fungal isolates:

All the isolated fungi were identified by studying the cultural characteristics, as well as the microsocial structures on Czapek's agar medium, according to Harigan and Margaret (1966), and according to the procedures of Gilman (1957), Barnnett and Hunter (1972) Numbers of the species of *Aspergillus* were classified according to the key published by Raper and Fennell (1965), while for species of other isolated fungi the "Manual of clinical mycology" by Count *et al.* (1954) was used.

Production of AFs by isolated *A. flavus* strain:

The ability of AFs production by *A. flavus* strains using liquid media (YES) was investigated according to Singh *et al.*, (1991).

Aflatoxin standard:

Preparation of aflatoxin standard was carried out according to the Association of Official Analytical Chemists, A.O.A.C.(2000). Crystals of aflatoxins B₁, B₂, G₁ and G₂ were diluted using benzene-acetonitrile (98: 2 v/v) to obtain a concentration of 8 to 10 µg/ml (stock solution).

Equipment and chromatographic conditions:

Extraction procedure

AFs were analyzed in nuts according to the method reported by Stroka, *et al.*, (2000). Briefly, 10 g of nut samples were added to 1 g of NaCl, then blended with 40 ml of methanol/water (80:20) and 20 ml of

cyclohexane for 3 min. After separation of the two phases, cyclohexane was eliminated. Extracts were filtrated on a Whatman filter paper No. 4. Immunoaffinity column clean-up An aliquot of 10 ml was diluted with 60 ml of PBS buffer (pH 7.4). An immunoaffinity column (IAC AflaTest, Vicam, USA) for AFs analysis was conditioned with 10 ml of PBS buffer by gentle syringe pressure at a flow rate of 5 ml/min. Then, the mixture of the filtrate diluted extract (70 ml) was applied to the IAC column (1–2 drops per second), followed by a washing with 20 ml of bi-distilled water and then dried with air. AFs were then slowly eluted from the IAC with 2 ml methanol into a glass vial.

Precolumn derivatization

The eluate was evaporated to dryness with a gentle stream of N₂ at 52 °C, redissolved with 100 µl of TFA for 3 min, re-evaporated to dryness with N₂ at 52 °C, and reconstituted in 500 µl of the mobile phase for LC analysis. TFA was added to AFs working standards in the same conditions as the extract samples to derivatize AFB₁, AFB₂, AFG₁ and AFG₂.

Determination of Aflatoxins A by HPLC.

High-performance Liquid Chromatography "HPLC" (Agilent 1100 series) equipped with a fluorescence detector (G 1321A) analysis was carried out with a liquid chromatograph equipped with solvent delivery systems (Agilent Technologies, Inc. 200 Regency Forest Drive, Suite 330 Cary, NC 27511 USA) system containing a G1322A Vacuum Degasser, a G1312A Binary pump and a reverse-phase analytical column packed with C₁₈ material (Agilent ZORBAX, DB- 5 µm, 150 mm × 4.6 mm). The mobile phase consisted of water: acetonitril: methanol (240:120:40) (Deabes, 2008). Separation was performed at 40 °C temperature at a flow rate of 1.0 ml min⁻¹; the injection volume was 50 µl for both standard solutions and sample extracts by auto sampler (G1329A). The detection was performed using fluorescence detector was operated at an excitation wave length of 360 nm and an emission wave length of 440 nm.

Quantitation: The mixed solutions of standard as well as sample extract after derivatisation were filtered through a 0.22 µm membrane filter and loaded (50 µl) into autosampler. The elution order of the four aflatoxins was G₂, B₂, G_{2a} (G₁ derivative), B_{2a} (B₁ derivative). AFs contents in samples were calculated from chromatographic peak areas using the standard curve.

3. Results and Discussion

Isolation and identification of fungi associated with nuts.

It is well known that some fungi grow on/ and in nuts and these fungi can deteriorate the stored nuts. Therefore, the present work, was started by isolation and identification of fungal species. Isolation of fungi was made by using agar as growing media.

Fungi isolation on Czapek's agar medium: Data in Table (1) represent the fungi associated with Czapek's agar medium from nuts purchased from retail markets in Qassim, Saudi Arabia during the summer season of 2009.

Table (1): Fungi associated with nut samples collected from Qassim region Saudi Arabia during 2009.


Commodities	No of samples	% of Infections	Total fungal Counts	Isolated Fungi		
				<i>Aspergillus</i>	<i>Penicillium</i>	<i>Rhizopus</i>
Almond	5	100	120	57.00	40.00	3.00
Hazelnut	5	100	153	60.00	32.00	8.00
Cashew	5	100	140	53.00	33.00	14.00
Walnuts	5	100	60	50.00	40.00	10.00
	5	100	160	57.00	33.00	10.00
	5	100	163	67.00	24.00	9.00

Table (2): Toxicity of *Aspergillus flavus* isolated from nuts and the ability for aflatoxins production in YES medium.

Kind of Nuts	Isolates of <i>Aspergillus</i> tested (No.)	Isolates of <i>A. flavus</i> (No.)	Isolates of <i>A. flavus</i> producing Aflatoxins (No.)	Isolates of <i>A. flavus</i> producing aflatoxins (%)	Total of Aflatoxins concentration mg	
					Min	Max
Almonds	40	25	22	88	20	40
Pistachio	60	42	33	78	12	60
Hazelnuts	65	32	27	84	16	50
Cashews	30	15	7	46	1.2	6
Walnuts	70	50	40	80	30	70
Peanuts	90	62	58	93	65	150

Table (3): Survey of aflatoxins in nuts collected from Qassim, Saudi Arabia during summer , 2009

Kind of nuts	No. of samples	No. of positive samples	% of positive samples	Aflatoxins concentration $\mu\text{g}/\text{kg}$							
				AFB1		AFB2		AFG1		AFG2	
				Min	Max	Min	Max	Min	Max	Min	Max
Almonds	5	2	40	38.00	45.00	11.00	13.00	0.00	2.30	0.00	4.20
Pistachio	5	4	80	11.00	90.00	2.00	70.00	1.10	55.00	1.90	25.00
Hazelnuts	5	2	40	41.00	55.00	12.90	18.00	3.10	6.00	5.20	8.40
Cashews	5	1	20	0.30	3.60	0.10	2.30	0.00	0.00	0.00	0.00
Walnuts	5	3	60	62.00	120.00	13.00	33.00	5.30	11.00	7.90	17.00
Peanuts	5	4	80	70.00	140.00	18.00	71.00	2.20	10.10	3.10	13.40

**Figure (1): High Performance Liquid Chromatogram of a positive sample of aflatoxins (AFG₁, AFB₁, AFG₂ and AFB₂)**




Figure (2): High Performance Liquid Chromatogram of a standard of aflatoxins (AFG₁, AFB₁, AFG₂ and AFB₂)




Figure (3): High Performance Liquid Chromatogram of a negative sample of aflatoxins (AFG₁, AFB₁, AFG₂ and AFB₂)

Data in Table (1) showed the percentage of infected nuts kernel and the percentage of the fungal genera which isolated in summer of 2009. Data revealed that, the percentage of fungal infected samples was 100% in all nuts samples collected from retail markets in Qassim region. The percentage averages of *Aspergillus*, *Pencillium* and *Rhziopus* genera infected were 57, 40, and 3.0 % in almond, 60, 32 and 8 in Pistachio, 53, 33 and 14 % in hazelnut, 50,40 and 10% in Cashew, 57, 33 10% in walnuts, 67,24 and 9% in

peanut collected from Qassim region, respectively. Infection of these samples with fungi may be due to the bad storage conditions in these region. Results also indicated that, fungi genera were *Aspergillus*, *Penicillium* and *Rhziopus* in all nut samples. It is known that, food species are common substrates for *A. flavus*, and subsequent aflatoxin production in these food stuffs is almost always due to poor drying, handling, or storage (Arim, 1995). Jimenez *et al.*, (1991) reported moulds and mycotoxins in almonds,

peanuts, hazelnut and pistachio nuts and detected aflatoxins at up to 95 mg/kg in the tested samples. The predominant fungi present in the samples were *A. flavus*, *A. niger*, *A. glaucus* Link ex Grey and *Penicillium* spp. In this concern, Vaamonde *et al.*, (2003) reported that, the incidence of aflatoxigenic *A. flavus* strains was higher in peanuts (69%) than in wheat (13%) or soybeans (5%) while the ratio of CPA producers *A. flavus* isolated from all substrates was very high (94% in peanuts, 93% in wheat and 73% in soybeans). Isolates of *A. flavus* able to produce simultaneously aflatoxins type B and CPA were detected in all substrates, suggesting the possibility of co-occurrence of these toxins. While, Mphande *et al.*, (2004) found that, the account of *Aspergillus* 41% in all the isolates and 98% in peanut samples. the account of *Aspergillus* spp. was 35% of all the isolates, while *Aspergillus niger* being the most prevalent (20.4%). *Aspergillus flavus* / *parasiticus* were also present and accounted for 8.5% of all the isolates, with *A. flavus* accounting for the majority of the *A. flavus* / *parasiticus* identified. Regional differences in aflatoxin contamination of crops may be attributable to climatic conditions and to agricultural practices that increase susceptibility of plants to invasion by *A. flavus* (Pildaina *et al.*, 2004).

Occurrence of mycotoxic fungi isolated from nuts :

Mycotoxins are secondary metabolites produced by specific filamentous fungi that contaminate agricultural commodities. Mycotoxins produced by an active growing mold. Molds can grow actively without mycotoxin formation. Therefore, these study was carried out to isolate mycotoxic fungi namely *A. flavus* producing aflatoxins.

From Table (2) it could be indicated that, *A. flavus* was the most mycotoxins- producing *Aspergillus* spp, represented, 88,78, 84, 46, 80 and 93% of *Aspergillus* spp isolates from almonds, pistachio, hazelnuts, cashews, walnuts, and peanuts collected from Qassim region, respectively were able to produce the aflatoxins. Results of Table (2) showed that, averages the range of total aflatoxins 20-40, 12-60, 16-50, 1.2-6, 30-70 and 65-150 mg /L of the yeast extract sucrose (YES) medium were produced by *A. flavus* isolates almonds, pistachio, hazelnuts, cashew, walnuts and peanuts samples collected from Qassim region, respectively. It is worthy to mention that, the current investigation to detect the aflatoxins (B_1 , B_2 , G_1 and G_2) levels and /or frequencies in the nut. Furthermore, Mphande *et al.*, (2004) investigated 32 isolates of *A. flavus* for their ability of mycotoxin production. They found that, 11 isolates did not produce detectable aflatoxins, 8 isolates produced only aflatoxins B_1 and B_2 , and 13 produced all four aflatoxins (B_1 , B_2 , G_1 and G_2) in varying amounts. When the raw peanut samples

(n=120) were analyzed for total aflatoxins, 78% contained aflatoxins at concentrations ranging from 12 to 329 $\mu\text{g}/\text{kg}$. A review of monitoring studies on the occurrence of aflatoxins in food products has demonstrated that aflatoxins are still being found frequently in food products at levels that are of significant concern for consumer protection (Scott and Lawrence, 1997 and Stroka and Anklam, 2002). The occurrence of aflatoxins in dried fruits and nuts was surveyed in the study by Lutfullah and Hussain, (2011) in Pakistan. They found the percentage of contamination for total aflatoxins in the samples such as in; dried apricot (20%), dates (10%), dried figs (50%), dried mulberries (26%) and raisins (20%), while in apricot kernels (26%), almonds without shell (30%), walnuts with shell (40%), walnuts without shell (70%), peanut with shell (40%), peanuts without shell (50%), pistachios with shell (20%), pistachios without shell (50%) and pine nuts with shell (20%). The highest contamination levels of aflatoxins were found in one peanut sample (14.5 mg/kg) and one pistachio sample (14 mg/kg). Molds of the genus *Aspergillus* frequently decay the kernel of pistachio nuts (Mojtahedi *et al.*, 1979). On the other hand, pistachio nuts are among the commodities with the highest risk of AF contamination (Pittet, 1998).

The obtained results of Table (3) and Figures (1,2 and 3) indicated that 80, 80, 60, 40, 40, 20 % for pistachio, peanuts, walnuts, almonds, hazelnuts and cashews samples collected from Qassim region, respectively were contaminated with aflatoxins B_1 and B_2 , aflatoxins G_1 and G_2 . The highly percentage of aflatoxins was found in both Pistachio and peanuts 80% with a concentrations of aflatoxins (AFB_1 , AFB_2 , AFG_1 and AFG_2) ranged from (11-90, 2-70, 1.1-55 and 1.9-25 " $\mu\text{g}/\text{kg}$ ") and (70-140, 18-71, 2.2-10.1 and 3.1-13.4 " $\mu\text{g}/\text{kg}$ " respectively). Figures 1, 2 and 3 shows the HPLC chromatogram of aflatoxins (G_1, B_1, G_2 and B_2) separation of standards and both positive and negative nuts sample, respectively.

Mycotoxin contamination in some edible dry fruits and nuts has been reported by Zohri and Abdel-Gawad (1993), Abdel-Hafez and Saber, (1993) and Singh *et al.*, (2001). Aflatoxins were detected in 90% of hazelnut samples (25–175 mg/kg) and 75% of walnut samples (15–25 mg/kg). In a survey of peanut products in North America, 19% of 1416 samples examined were contaminated with an average level of 1 $\mu\text{g}/\text{kg}$ (Stoloff 1977), In Thailand 49% of 216 samples contained AFB_1 at an average level of 424 $\mu\text{g}/\text{kg}$ (Shank *et al.* 1972). Aflatoxins are present in food chain consumption of aflatoxin in many parts of the world varies from 0 to 30 000 ng/kg/day (Denning, 1987). Some factors such as high temperature and low moisture can result in cracks in the seed and subsequent invasion by the fungus. Temperature and moisture are

the dominant factors that affect aflatoxin contamination of corn. Environmental conditions most favorable for maximum growth and aflatoxin production by *A. flavus* are temperatures greater than 30°C, maximum relative humidity of greater than 85%, and water activity of 0.98 to 0.99 (Payne *et al.*, 1988). Thus, *A. flavus* can infect with proper moisture/temperature conditions during storage almost any stored product (Payne, 1992). Aflatoxin formation in groundnut is favored by prolonged end of season drought and associated elevated temperature (Rachaputi, *et al.*, 2002). Singh and Shukla (2008) investigated fungal infection and mycotoxin contamination in fresh and stored kernels of walnuts collected from different localities of Uttaranchal (India). They found the species of *Alternaria*, *Aspergillus* and *Penicillium* were predominant. Also, they found that, Thirty-nine percent of *Aspergillus flavus* isolates were toxigenic and produced up to 2170 mg/l of aflatoxin B₁ in the liquid media. Aflatoxin B₁ was the most common mycotoxin encountered as a natural contaminant in stored samples.

The current investigation was carried out to detect and determine the aflatoxins quantitatively (AFB₁, AFB₂, AFG₁ and AFG₂) and contamination levels in the nuts samples collected from retail markets in Qassim, region.

Conclusion

Studies were carried out to investigate the mycoflora profile and aflatoxin contamination in red nut kernels (almonds, peanuts, hazelnuts, pistachio nuts, walnut and cashew). *Aspergillus flavus*, *A. niger* and *Penicillium* sp the predominant fungi. The concentration of aflatoxin B₁ was high in peanut and walnut kernels as well as in the toxigenic strains of *A. flavus* isolated from these kernels. There is an urgent need to check the use of contaminated kernels and their products for edible purposes, aflatoxins are potent carcinogens which pose the risk of mycotoxicoses for the consumers.

References:

1. Abdel-Hafez, A.I.I., Saber, S.M. Mycoflora and mycotoxin of hazelnut (*Corylus avellana* L.) and walnut (*Juglans regia* L.) seeds in Egypt. Zentralblatt für Mikrobiologie 1993: 148; 137–147
2. AOAC. Official methods of analysis of the Association of Analytical Chemists. Edited by H. William 2000: 17th Ed., Vol II, Association of Official Analytical Chemists, Arlington VA
3. Arim, R.H. Present status of the aflatoxin situation in the Philippines. Food Addit. Contam 1995: 12; 291-296.
4. Barnett, H.L. and B.B. Hunter. Illustration genera of imperfect fungi. Burgess Publishing Company, Minn-Eapolis 15 min. U.S.A. 1972: 225.
5. Bathnagar, D. and Garcia, S.. Aspergillus. In: Labbe', R.G., Garci'a, S. (Eds.), Guide to Foodborne Pathogens. John Wiley and Sons, New York 2001; pp. 35– 49.
6. Conner, D.E. Conner, Naturally occurring compounds. In: P.L. Davidson and A.L. Branen, Editors, *Antimicrobials in Foods*, Marcel Dekker, New York 1993; 441–468.
7. Count, N.F.; F.S. Martin; D.T. Smith; R.F. Baker and J.L. Gallaway. Manual of clinical mycology. Philadelphia, W.B. Saunders Company 1954.
8. Deabes, M.M. Studies on the control of some mycotoxins contamination in some foods. Ph.D. Thesis, Food Science & Technology Dept., Faculty of Agriculture, Al-Azhar Univ., Cairo, Egypt 2008.
9. Denning, D. W. Aflatoxin and human diseases, p. 175-209. In D. M. Davies and H. deGlanville (ed.), Adverse drug reactions and acute poisoning reviews 1987: vol. 4. Oxford University Press, Oxford, U.K.
10. Farr, D.F., Billis, G.F., Chamuris, G.P., Rossman, A.Y. Fungi on plants and plant products in the United States. American Phytopathological Society Press, St. Paul, MN 1989; 1– 152.
11. Gilman, J.E.. A manual of soil fungi" low state college press, Anes, Iowa, U.S.A. 1957; 392 .
12. Gourama, H., Bullerman, L.B.. Antimycotic and antiaflatoxigenic effect of lactic acid bacteria—a review. Journal of Food Protection 1995: 57; 1275– 1280.
13. Gradelet, S., Astorg, P., Le, A.M., Berges, R., Suschetet, M.. Modulation of aflatoxin B₁ carcinogenicity, genotoxicity and metabolism in rat liver by dietary carotenoids: evidence for a protective effect of CYP1A inducers. Cancer Letters 1997:114; 221–223.
14. Gqaleni, N.; Smith, J.E. and Lacey, J. Co-production of aflatoxins and cyclopiazonic acid in isolates of *Aspergillus flavus*. Food. Addit. Contam 1996:13(6):677-685.
15. Harrigan, W.F. and Margaret, E. McCance. Laboratory methods of Microbiology. Department Bacteriology the West of Scotland, Agricultural Collage. Auclin. Cruine Ayr, Scotland 1966.
16. Ichinoe, M.E.; H.A. Kurata; Y.B. Sugtura and Y.S. Uemo. Chemotaxonomy of *Gibberella zea* with special reference to production of trichothecenes and zeralenone. Applied and Environmental Microbiology 1983: 46; 1364-1369.
17. JECFA. Forty-ninth meeting of the joint FAO/WHO expert committee on food additives. In: Safety Evaluation of Certain Food Additives and Contaminants: Aflatoxins WHO Food Additives Series 40. WHO, Geneva 1998; 359–469.
18. Jimenez, M., Mateo, R., Querol, A., Huerta, T., Hernandez, E.,. Mycotoxins and mycotoxigenic

- moulds in nuts and sunflower seeds for human consumption. *Mycopathologia* 1991;115;121–127.
19. Lichtwardth, R.W.; G.L. Basrn and L.H. Tiffang. Fungus growth in shelled corn as affected by moisture. *J. Agr. Res.* 1958: 56; 291-307.
 20. Luttfullah G and Hussain, A. Studies on contamination level of aflatoxins in some dried fruits and nuts of Pakistan. *Food Control* 2011;22; 426-429.
 21. Martin, J., Ba, A., Dimanche, P., and Schilling, R.. How groundnut contamination can be controlled? *Work in Senegal.Agriculture et Development* 1999: 23; 58–67.
 22. Mojtahedi, H., Rabie, C.J., Lubben, A., Steyn, M., Danesh, D. Toxic aspergilli from pistachio nuts. *Mycopathol* 1979: 67; 123–127.
 23. Mphande, F.A.; Siame,B.A. and Taylor, J.E.. Fungi, aflatoxins, and cyclopiazonic acid associated with peanut retailing in Botswana. *J Food Prot* 2004: 67(1); 96-102.
 24. Mislivec, P.B.. Direct plating versus dilution planting in qualitatively determining the mold flora of dried beans and soybeans. *Journal Association of Official Analytical Chemists* 1977: 60;741-743.
 25. Ministry of Agriculture and Rural Affairs, Republic of Turkey. Data on Aflatoxins in hazelnuts, pistachios, walnuts and almonds during 1998–2002.
 26. Pittet, A. Natural occurrence of mycotoxins in foods and feeds – an updated review. *Revue. Med. Vet* 1998: 149; 479–492.
 27. Payne, G. A. Aflatoxin in maize. *Crit Rev Plant Sci* 1992;10(5);423–440.
 28. Payne, G. A.; Hagler, W. M. and Adkins. C. R.. Aflatoxin accumulation in inoculated ears of field-grown maize. *Plant Dis.* 1998: 72; 422–424.
 29. Raper, J.P. and D.T. Fennell. The genus *Aspergillus williams* and *Wilkins*. Baltimorr, U.S.A 1965.
 30. Rachaputi, N .R.; Wright, G .C. and Kroschi, S.. Management practices to minimise pre-harvest aflatoxin contamination in Australian groundnuts. *Austr. J. Exp. Agric* 2002: 42; 595-605.
 31. Scott , P. M. and Lawrence, G. A.. Determination of Aflatoxins in beer. *Journal of AOAC International* 1997: 80; 1229–1234.
 32. Shank, R. C.; Gordon, J. E.; Wogan, G. N.; Nondasuta, A. and Subhamani, B. Dietary aflatoxins and human liver cancer. III. Field survey of rural Thai families for ingested aflatoxins. *Food Cosmet. Toxicol.* 1972: 10; 71-84.
 33. Singh, K., Frisbad, J.C., Thrane, U., Mathur, S.B.. An illustrated manual on identification of some seed-borne *Aspergilli*, *Fusaria*, *Penicillia* and their mycotoxins. Danish government. Insitute of Seed Pathology for Developing Countries, Hellerup 1991; 8, 12.
 34. Singh, P. K. and Shukla, A.N. () Survey of mycoflora counts, aflatoxin production and induced biochemical changes in walnut kernels *Journal of Stored Products Research* 2008;44;169–172.
 35. Stroka, J. and Anklam, E. New strategies for the screening and determination of aflatoxins and the detection of aflatoxin-producing moulds in food and feed. *Trends in Analytical Chemistry* 2002;21;90–95.
 36. Stroka, J., Anklam, E., Jorissen, U., and Gilbert, J. Immunoaffinity column cleanup with liquid chromatography using post-column bromination for determination of aflatoxins in peanut butter, pistachio paste, fig paste, and paprika powder: Collaborative study. *Journal of AOAC International* 2000: 83; 320–340.
 - 37 Singh, P.K., Khan, S.N., Harsh, N.S.K., Pandey, R.. Incidence of mycoflora and mycotoxins in some edible fruits and seeds of forest origin. *Mycotoxin Research* 2001: 17; 46–58.
 38. Stoloff, L.. Aflatoxins - an overview. In "Mycotoxins in Human and Animal Health", eds. J.V. Rodricks, C.W. Hesseltine, and M.A. Mehlman. Park Forest South, Illinois: Pathotos Publishers 1977; 7-28.
 39. Thuvander, A., Moller, T., Barbieri, H.E., Jansson, A., Salomonsson, A.C., Olsen, M. Dietary intake of some important mycotoxins by the Swedish population. *Food Addit. Contam* 2001;18;696–706.
 40. Vaamonde, G.; Patriarca, A.; Fernandez Pinto, V.; Comerio, R. and Degrossi, C.. Variability of aflatoxin and cyclopiazonic acid production by *Aspergillus section flavi* from different substrates in Argentina. *Int J Food Microbiol* 2003: 88(1); 79-84.
 41. Vardon, P.J. Mycotoxins: risks in plant, animal and human systems, potential economic costs of mycotoxins in the United State. *Cast Task Force* 2003, pp. 136– 142. Report No. 139.
 42. Zohri, A.A., Abdel-Gawad, K.M. Survey of mycoflora and mycotoxins of some dried fruits in Egypt. *Journal of Basic Microbiology* 1993;33;279-288.

8/2/2011