

**The Protective role of brown alga (*Sargassum crassifolia*) against the degenerative toxic effects induced by Nimbecidin in muscles of Nile Tilapia, *Oreochromis niloticus* L.**

Hawazin H. Mutawie\* and Asmaa M. Hegazi\*\*

\*Department of Biology, Faculty of Applied Sciences, Umm Al-Qura University, Makkah, Saudi Arabia.

Corresponding author : e-mail [hmutawie@uqu.edu.sa](mailto:hmutawie@uqu.edu.sa)

\*\*Department of Aquatic Pathology, National Institute of Oceanography and Fisheries, Al-Qanater, Cairo, Egypt. e-mail [dr.amhegazy@gmail.com](mailto:dr.amhegazy@gmail.com). [sabsak@yahoo.com](mailto:sabsak@yahoo.com) for acceptance

**Abstract:** This study was undertaken to evaluate the potential effect of the brown alga *Sargassum crassifolia* against biochemical and histological alterations in the muscles of fresh water fish, *Oreochromis niloticus*, exposed to 1/10 LC<sub>50</sub> (0.03 ppm) of Nimbecidin (azadirachtin). *S. crassifolia* was added to the basal diet of fish in different proportions (0, 10 and 25%) and the experiment lasted for 4 weeks. Exposure of fish to Nimbecidin induced significant decrease in the activity of reduced glutathione, catalase and in the total protein content of fish muscles. Also many histopathological changes including degeneration and necrosis, disorganized myofibers with chromophobic cytoplasm, haemocytic infiltration, oedema and inflammations were noticed in Nimbecidine - treated fish. **Exposing** fish to Nimbecidin and *Sargassum crassifolia* (10% and 25%) led to marked improvement in the examined biochemical parameters together with the histological structure of muscles. This improvement was more obvious at high concentration of *Sargassum crassifolia*. In conclusion, the results of the present work indicated that the brown alga *Sargassum crassifolia* had ameliorative effect against muscle damage induced by Nimbecidin and this may be mediated by its potent antioxidant activities.

[Hawazin H. Mutawie and Asmaa M. Hegazi. **The Protective role of brown alga (*Sargassum crassifolia*) against the degenerative toxic effects induced by Nimbecidin in muscles of Nile Tilapia, *Oreochromis niloticus* L.** Journal of American Science 2011;7(12):422-431]. (ISSN: 1545-1003). <http://www.americanscience.org>.

**Key words:** *Sargassum crassifolia*, *Oreochromis niloticus*, antioxidant, neem, Nimbecidin, muscles, Glutathione, Catalase, Total protein.

## 1- Introduction

Marine algae are rich sources of structurally new and biologically active metabolites. In recent years, there have been many reports of macroalgae derived compounds that have a broad range of biological activities, such as antiviral (Ehresmann *et al.*, 1977), antimicrobial (Devi *et al.*, 2008), anti-inflammatory (Dar *et al.*, 2007; Kang *et al.*, 2008 and Ananthi *et al.*, 2010), anticarcinogenic and antioxidant **activities** (Kim *et al.*, 2005; Athukorala *et al.*, 2006; Kukovinets and Kislitsyn, 2006; Devi *et al.*, 2008; Iwai, 2008; and Ananthi *et al.*, 2010).

Free radicals scavengers and antioxidants can reduce lipid peroxidation and the generation of ROS. The importance of antioxidants in human health has become increasingly clear due to spectacular advance in understanding the mechanisms of their reaction oxidants. Furthermore, interest in employing antioxidants from natural sources to increase the shelf-life of foods is considerably enhanced by consumer preference for natural ingredients and concerns about toxic effects of synthetic antioxidants (Heo, *et al.*, 2005). During the last years, several authors reported that certain species of brown and red algae contained multitude of bioactive compounds that have antioxidant properties (Matanjun *et al.*, 2008, Plaza *et al.*, 2008). Recently, the antioxidant activity of the brown algae

*Sargassum crassifolia* inhibiting Jeddah corniche, Saudi Arabia, was reported (Al-Amoudi *et al.*, 2009).

Neem; *Azadirachta indica* (*A. indica*), is one of the most promising medicinal **plants**, having a wide spectrum of biological activity, well known for its insecticidal properties. Neem leaf extract and neem-based products were successfully implied in fish farming as antiparasitic and antibacterial agent (Harikrishnan *et al.*, 2003 and Mousa *et al.*, 2008) and as control of fish fry predators (Dunkel and Ricilards, 1998). Martinez and Souza (2002) reported that aqueous extract of neem leaves and other neem-based products have been extensively used in fish-farms as alternative for the control of fish parasites and fish fry predators such as dragonfly larvae. Although neem extract is considered of low toxicity towards non-target aquatic life, water extracts of the bark of the neem plant caused respiratory problems in *Tilapia zilli* (Omeregie and Okpanachi, 1997), while long exposure to low concentrations of the crude extract of *A. indica* delayed the growth of this cichlid fish (Omeregie and Okpanachi, 1992). Such results indicate that neem extracts added to water may cause disturbances on fish. Consequently it is important to recognize the effects that the employ of these products used to prevent the appearance of diseases may have in different parameters of aquatic life (Rábago-Castro *et al.*, 2006).

The present work was aimed to study the effect of *Sargassum crassifolia*, a brown alga from Saudi Arabia coasts, against the toxicity induced by Nimbecidin, natural plant insecticide derived from neem plant, in the muscles of the fresh water tilapia, *O. niloticus*.

## 2- Materials and methods

### 2.1. Nimbecidin

Nimbecidin®, a commercial natural pesticide derived from neem tree, *Azadirachta indica*. The active ingredient of nimbecidin is azadirachtin (Aza) which makes about 0.03% of its chemical composition and the other relevant ingredients are neem oil (90.57%), Hydroxyel (5.00%), Epichlorohydrate (0.50%), Aromax (5.90%).

### 2.2. Algal materials

*Sargassum crassifolia* (brown alga) was collected along Jeddah corniche, about 22 km through the longitudinal direction of N-W. The collected samples were carefully cleaned from epiphytes and washed several times with tap and distilled water then air-dried, chopped into small pieces and powdered and preserved in a desiccator for the determination of protein, fat and ash contents.

#### 2.2.1. Crude protein content

Crude protein was determined by measuring the total nitrogen content of the sample multiplied by the empirical factor 6.25. This method accomplished briefly in three stages: digestion by sulfuric acid, distillation and titration (Hepher *et al.*, 1983).

#### 2.2.2. Crude lipid content (ether extract)

Crude lipid content was determined by weighing the filter paper containing the dried algal sample and then transferred to Soxhlet apparatus using petroleum ether at 60-80 °C for 12 hrs. The sample with filter paper was dried and reweighed, the difference between sample weights indicates the total lipid content in the sample (AOAC, 1995).

#### 2.2.3. Ash content

Ash content was determined by weighing the crossable containing the dried sample and then transferred to a muffle furnace at 550 °C for 8 hrs. The crossable containing the sample was reweighed, the difference between sample weights indicates the ash content (AOAC, 1995).

### 2.3. Experimental animals

Nile tilapia (*Oreochromis niloticus*) with an average weight  $14 \pm 1.2$ g were obtained from a fish hatchery in Abbassa, Abo-Hammad, Sharkia, Egypt and transferred immediately to the laboratory in sacs

filled with fresh water and supplied with continuous system of aeration. In the laboratory, fish were kept in 3.3 m<sup>3</sup> aquaria filled with dechlorinated tap water and supplied with continuous system of filtration and aeration. Temperature was maintained at  $21 \pm 2$  °C and dissolved oxygen values at  $7.0 \pm 0.5$  mg/L. Fish were fed with wheat bran (3% of total fish weight/day) and acclimatized for 15 days before the beginning of the experiments.

### 2.4. Toxicity test of Nimbecidin pesticide

Following acclimatization, ten groups of ten fishes were placed in 100 L aerated aquaria and exposed to a series of Nimbecidin concentrations that would permit the computation of the 96 hrs LC<sub>50</sub> value and mortalities of fish were recorded daily.

### 2.5. Experimental Design:

In a four week experiment, the fish were assigned to 6 groups of 50 fish each:

Group 1 (G1): Animals of this group were used as controls and fed on basal diet without *S. crassifolia*.

Group 2 (G2): The diet of this group consisted of the basal diet (90%) mixed with *S. crassifolia* (10%).

Group 3 (G3): Animals of this group were fed on basal diet (75%) mixed with *S. crassifolia* (25%).

Group 4 (G4): Animals of this group were exposed to 0.03 ppm (1/10 LC<sub>50</sub>) of Nimbecidin and fed on normal basal diet.

Group 5 (G5): Animals of this group were exposed to 0.03 ppm (1/10 LC<sub>50</sub>) of Nimbecidin and fed on basal diet (90%) mixed with *S. crassifolia* (10%).

Group 6 (G6): Animals of this group were exposed to 0.03 ppm (1/10 LC<sub>50</sub>) of Nimbecidin and fed on basal diet (75%) mixed with *S. crassifolia* (25%).

At termination of 2<sup>nd</sup> and 4<sup>th</sup> experimental weeks, samples of fishes from all groups were dissected and muscle samples were removed.

### 2.6. Biochemical analysis

For biochemical study, muscle samples from all fish groups were washed in saline solution and homogenized (10% w/v) in ice-cold normal saline. The homogenate was centrifuged at 10 000xg for 20 min at 4 °C and the resultant supernatant was used for biochemical assay according to the method described by Chitra *et al.* (1999).

#### 2.6.1. Antioxidants: Reduced glutathione (GSH) and catalase

The biochemical assays of reduced glutathione (GSH) and catalase were determined according to the methods described by Ellman (1959), Aebi *et al.*

(1974) and Okhawa *et al.* (1979) respectively. The activity of glutathione was expressed as micromoles per wet gram of tissue and catalase was expressed as IU per mg protein.

## 2.6. 2. Total protein content

The total protein content of muscles was estimated by the method of Bradford (1976) and expressed as mg per gram of tissue.

## 2.7. Histological study

For histological study, samples of trunk muscles from control and treated fish were removed, rapidly fixed in Bouin's fluid and routinely processed for paraffin embedding. Sections were cut at 5  $\mu$ m and stained with haematoxylin and eosin.

## 2.8. Statistical analysis

Biochemical results were subjected to one-way analysis of variance (ANOVA) and represented as mean  $\pm$ SD of eight fishes per group. Differences in mean values between groups were assessed by Tukey's test and were considered statistically different at  $p < 0.05$ .

## 3- Results

### 3.1. Effect of Nimbecidin on the mortality of *Oreochromis niloticus*

The toxicity of Nimbecidin against *O. niloticus* was evaluated to determine the median lethal and sublethal concentrations. Observation of the mortality rate over a period of 96 hours of *O. niloticus* exposed to various concentrations of Nimbecidin showed that the percentage of mortality of fishes depended on the concentrations of Nimbecidin. The  $LC_{50}$  of Nimbecidin for *O. niloticus* was found to be 0.33 ppm.

### 3.2. Analysis of algal material.

Results of analysis of the chemical composition of *S. crassifolia* were 15.31  $\pm$  0.36 mg/g dry weight (for crude protein), 1.86  $\pm$  1.5 mg/g dry weight (for crude lipid) and 35.34  $\pm$  1.04 mg/g dry weight (for ash).

## 3.3. Biochemical analysis

### 3.3.1. Antioxidants

#### 3.3.1.1. Reduced glutathione (GSH) :

Data in Table (1) shows that there was a significant reduction ( $p < 0.05$ ) in the activity of the mitochondrial antioxidant reduced glutathione (GSH) in the muscles of fish treated with Nimbecidin compared with that of control. Treating animals with Nimbecidin together with *S. crassifolia* (10% and 25%) for 2 & 4 weeks induced significant increase ( $p < 0.05$ ) in glutathione (GSH) when compared with control

animals (group 1) or animals treated with Nimbecidin alone (group 4).

### 3.3. 1.2. Catalase :

Significant decrease ( $p < 0.05$ ) in catalase was recognized in fish exposed to Nimbecidin (groups 4) compared to that in control animals (group 1) or in animals fed with 10% and 25% of *S. crassifolia* (group 3 & 4 respectively). When animals treated with both Nimbecidin plus 10% *S. crassifolia* for 2 weeks, no significant change was recorded in catalase activity compared with that in fish exposed to Nimbecidin alone. Significant increase in the activity of catalase was recorded in animals treated with Nimbecidin plus 25% *S. crassifolia* for 2 weeks and in animals treated with Nimbecidin together with *S. crassifolia* (10% and 25%) for 4 weeks (Table 2).

### 3.3.2. Total protein:

Results exist in (Table 3) revealed that there was a significant ( $p < 0.05$ ) decrease in the total protein content in muscles of animals treated with Nimbecidin as compared with control (group 1). Animals given Nimbecidin and *S. crassifolia* (10% and 25%) for 2 & 4 weeks showed significant increase in the total protein compared with Nimbecidin treated group.

**Table 1:** Effect of *S. crassifolia* and / or Nimbecidin on the activity of reduced glutathione ( $\mu$ mol/wet g tissue) in the muscles of fish at two time periods.

Groups	Time periods( weeks)	
	2weeks	4weeks
Group 1	171.18 $\pm$ 6.07 <sup>a</sup>	171.18 $\pm$ 6.07 <sup>a</sup>
Group 2	172.86 $\pm$ 9.1 <sup>a</sup>	176.91 $\pm$ 1.01 <sup>ac</sup>
Group 3	176.82 $\pm$ 1.81 <sup>a</sup>	187.67 $\pm$ 10.09 <sup>c</sup>
Group 4	164.18 $\pm$ 6.27 <sup>b</sup>	163.26 $\pm$ 8.43 <sup>b</sup>
Group 5	210.36 $\pm$ 1.26 <sup>c</sup>	204.47 $\pm$ 14.66 <sup>d</sup>
Group 6	240.19 $\pm$ 13.39 <sup>d</sup>	212.89 $\pm$ 13.11 <sup>e</sup>

Data are represented as mean  $\pm$  SD of eight specimens.

Data with different superscript letters in a column are significant to each other at  $p < 0.05$

**Table 2:** Effect of *S. crassifolia* and / or Nimbecidin on the activity of catalase ( $\mu$ /mg protein) in the muscles of fish at two time periods.

Groups	Time periods( weeks)	
	2weeks	4weeks
Group 1	3.77 $\pm$ 0.61 <sup>a</sup>	3.71 $\pm$ 0.38 <sup>a</sup>
Group 2	3.98 $\pm$ 0.88 <sup>a</sup>	3.87 $\pm$ 0.56 <sup>a</sup>
Group 3	3.99 $\pm$ 0.78 <sup>a</sup>	3.99 $\pm$ 0.52 <sup>a</sup>
Group 4	2.84 $\pm$ 0.16 <sup>b</sup>	3.06 $\pm$ 0.01 <sup>b</sup>
Group 5	2.86 $\pm$ 0.23 <sup>bc</sup>	3.29 $\pm$ 0.08 <sup>a</sup>
Group 6	3.06 $\pm$ 0.19 <sup>c</sup>	3.77 $\pm$ 0.08 <sup>c</sup>

Data are represented as mean  $\pm$  SD of eight specimens.

Data with different superscript letters in a column are significant to each other at  $p < 0.05$ .

### 3.4. Histological results:

Control fish (Fig. 4a) and fish fed on 10% and 25% *S. crassifolia* supplementations (Figs 4b & c respectively) had normal muscle structure with no apparent tissue lesions.

In the muscles of animals treated with Nimbecidin for two weeks ( Fig. 4d) degeneration and necrosis processes, disorganized myofibers with chromophobic cytoplasm, haemocytic infiltration, oedema and inflammatory changes were noticed. Less proliferated

muscle lesions were recorded in animals exposed to Nimbecidin and fed on diet containing 10% and 25% *S. crassifolia* ( Figs. 4e & f respectively).

Exposure of fish to Nimbecidin for 4 weeks ( Fig. 5a) induced sever histopathological changes including necrosis and inflammatory changes, oedema and appearance of microhemorrhagic zones. These pathological changes were less pronounced in fishes fed diet supplied with 10% and 25% *S. crassifolia* (Figs 5b & c respectively).




Fig. 4: Sections in muscles of control Nile tilapia, *O. niloticus* and fish fed on ascending doses (0%, 10% and 25%) of *S. crassifolia* supplementations and subjected to 0.03 ppm Nimbecidin, or without it, for 2 weeks duration. (a) Control fish (Group 1), (b) Fish fed on supplementary 10% *S. crassifolia* (Group 2). (c) Fish fed on supplementary 25% *S. crassifolia* (Group 3) (d) Fish subjected to Nimbecidin (Group 4) showing degeneration and necrosis processes, disorganized myofibers with chromophobic cytoplasm, haemocytic infiltration, oedema and inflammatory changes. (e) Fish subjected to Nimbecidin+ supplementary 10% *S. crassifolia* (Group 5). Muscle lesions appeared less proliferated. (f) Fish subjected to Nimbecidin+ supplementary 25% *S. crassifolia* (Group 6). Muscle lesions appeared less proliferated. H&E-stained muscle sections. Bar= 100  $\mu$ m.




Fig. 5: Sections in muscles of Nile tilapia, *O. niloticus*, treated with 0.03 ppm Nimbecidin and ascending doses (0%, 10% and 25%) of *S. crassifolia* supplementation for 4 weeks duration. (a) Nimbecidin-subjected fish (Group 4) showing sever necrotic and inflammatory changes, oedema microhemorrhagic zones were observed. (b) Fish subjected to Nimbecidin+ supplementary 10% *S. crassifolia* (Group 5). Muscle lesions appeared less proliferated. (c) Fish subjected to Nimbecidin+ supplementary 25% *S. crassifolia* (Group 6) showing myofibrils with central nucleus and partially uniform cytoplasm and some degree of fat infiltration in the interstitial tissue. H&E-stained muscle sections. Bar= 100  $\mu$ m.

**Table 3:** Effect of *S. crassifolia* and / or Nimbecidin on the total protein content (mg/g tissue) in the muscles of fish at two time periods.

Groups	Time periods( weeks)	
	2weeks	4weeks
Group 1	0.64 $\pm$ 0.03 <sup>a</sup>	0.65 $\pm$ 0.04 <sup>a</sup>
Group 2	0.66 $\pm$ 0.06 <sup>ac</sup>	0.67 $\pm$ 0.06 <sup>ac</sup>
Group 3	0.7 $\pm$ 0.06 <sup>c</sup>	0.71 $\pm$ 0.06 <sup>c</sup>
Group 4	0.61 $\pm$ 0.02 <sup>b</sup>	0.60 $\pm$ 0.03 <sup>b</sup>
Group 5	0.65 $\pm$ 0.02 <sup>a</sup>	0.64 $\pm$ 0.03 <sup>a</sup>
Group 6	0.85 $\pm$ 0.02 <sup>d</sup>	0.73 $\pm$ 0.04 <sup>d</sup>

Data are represented as mean  $\pm$  SD of eight specimens.

Data with different superscript letters in a column are significant to each other at  $p < 0.05$ .

#### 4- Discussion

The nutritional value of algal species depends particularly on its biochemical composition. In the present study total protein given by *S. crassifolia* was 15.31mg/g dry weight and it is comparable to that reported by Al-Amoudi *et al.* (2009) for such species (16.7mg/g). Burtin (2003) reported that the protein content in brown seaweeds are generally ranging from 5 to 15% of dry weight, compared to red and green seaweeds (ranging from 10 to 30%). Chemical

analysis of the seaweeds collected from the Persian Gulf of Iran showed that the brown seaweed *Gracilaria corticata* had the highest protein content (19.3%, dry weight basis) and *S. ilicifolium* had the lowest (8.9%) (Ghadikolaei *et al.*, 2011). The differences in chemical composition between seaweeds are taxonomically and environmentally related (Herbreteau *et al.*, 1997 and Lourenço *et al.*, 2002).

In the present study, lipid content of *S. crassifolia* was found to be 1.86 mg/g dry weight and it is within the range for most seaweeds (1–3%) (; Sánchez-Machado *et al.*, 2004 Matanjun *et al.*, 2009 and Ghadikolaei *et al.*, 2011). Matanjun *et al.* (2009) found that the total lipid content in *E. cottonii*, *C. lentillifera* and *S. polycystum* were 1.10%, 1.11% and 0.29%, respectively.

Ash was the most abundant component of dried seaweeds. The level of ash recorded currently for *S. crassifolia* was 35.34 mg/g dry weight and it is comparable to those previously reported by Ortega-Calvo *et al.*, (1993), Rupérez, (2002) and Matanjun *et al.*, (2009).

Chemical and some plant pesticides may induce oxidative stress leading to generation of free radicals

and alteration in antioxidant or oxygen free-radical scavenging enzyme system and disturb membrane structure ( El-Shenawy *et al.*, 2010).

In fish, many environmental pollutants, including pesticides, are capable of inducing oxidative stress (Hincal *et al.*, 1995; Dorval *et al.*, 2003; Pandey *et al.*, 2003; Sayeed *et al.*, 2003; Monteiro *et al.*, 2006). The authors mentioned that this event results in the formation of highly reactive compounds such as free radicals or oxyradicals ( $O_2^-$ ,  $H_2O_2$  and  $\cdot OH$ ) that frequently react with cellular macromolecules, leading potentially to enzyme inactivation, lipid peroxidation, DNA damage and even cell death (Van der Oost *et al.*, 2003).

Glutathione-S-transferases (GST) are a group of enzymes that catalyze the conjugation of reduced glutathione (GSH) with a variety of electrophilic metabolites, and are involved in the detoxification of both reactive intermediates and oxygen radicals (Van der Oost *et al.*, 2003). In the present study exposure of Nile tilapia (*Oreochromis niloticus*) to sublethal dose of Nimbecidin induced significant reduction in the activity of glutathione (GSH). Similar findings were observed by Winkaler *et al.* (2007) who observed enhanced hepatic GST activity in freshwater fish *Prochilodus lineatus* following exposure to 5.0 g/L of neem extract and attributed it to the metabolism of organic compounds in the extract. Depletion of GSH level was also recorded in gills, muscles and intestine of *O. mossambicus* exposed to sub-lethal concentration (30  $\mu g/L$ ) of profenofos for 28 days (Kavitha and Rao, 2009) and in the freshwater fish, *Channa punctata*, in response to heavy metals (Pandey *et al.*, 2008).

Administration of *S. crassifolia* in combination with Nimbecidin induced significant elevation in GSH activity in comparison to that in animals exposed to Nimbecidin alone.

Elevation of GSH activity could be attributed to the antioxidant activity of *S. crassifolia* (Al-Amoudi *et al.*, 2009).

Catalase is one of the free radical scavenging enzymes which form the first line of defence against oxidative injury converting  $H_2O_2$  into  $H_2O$  and  $O_2$  (Dorval *et al.* 2003).

In the present study depletion in catalase values was found in muscles of fish treated with Nimbecidin. Bairy *et al.* (1996) declared that the depletion in level of catalase activity might be due to an increased production of the superoxide radical ( $O_2^-$ ), as an excess of this anion is known to inhibit catalase activity. Similarly, Winkaler *et al.* (2007) found that fish exposed to the sublethal concentrations of neem extract showed significant reduction in hepatic catalase activity which is likely to affect the capacity of liver cells to defend themselves and respond to contaminant-

induced oxidative stress. In tilapia, Avci *et al.* (2005) reported significant reduction in CAT activity in the muscle of *O. niloticus* obtained downstream of Kizilirmak River, Turkey, due to exposure to petrochemical industry contamination. Depletion was also recorded in various organs in fish in response to different stressors (Kavitha and Rao, 2009).

CAT activity in the muscles of *O. niloticus* was found to be ameliorated after treating with 10% and 25% of *S. crassifolia*. Atencio *et al.* (2009) demonstrated that low dose of Se supplementation has shown to protect CAT activity in the liver and kidney of cyanobacterial cells-exposed *O. niloticus*. Also, Karthikeyan *et al.* (2010 & 2011) reported that the diminishing in activity antioxidant enzymes including CAT in liver, blood and kidney in rats was restored significantly after an ameliorating period using the brown alga *P. boergesenii* at a concentration of 150 mg/kg body weight.

In the current study, administration of Nimbecidin induced reduction in the total protein content in the muscles of treated fish. Also, Hussein *et al.* (1996) observed a significant fall in the total plasma protein level in *O. niloticus* after exposure to the herbicide atrazine for 14 and 28 days. Jee *et al.* (2005) reported decreased blood proteins and albumin levels in *Sebastes schlegeli*, the Korean rockfish, exposed to a sublethal concentration of cypermethrin (pyrethroid insecticide) for 8 weeks. On the other hand, Winkaler *et al.* (2007) reported that the fish showed no variation in total plasma protein levels after exposure to sublethal concentrations of neem leaf extract. Administration of *S. crassifolia* has the ability to increase the total protein after 2 and 4 weeks of treatment. In tilapia, the defensive role of vitamin E on protein oxidation in liver and kidney of microcystins-exposed tilapias has been demonstrated (Prieto *et al.*, 2006).

Results obtained in the current work revealed that Nimbecidin caused many histological lesions which are progressively altered with time. These lesions included degenerative, oedemic disorganized myofibers with chromophobic cytoplasm, haemocytic infiltration, vacuolization, necrotic effects, and presence of microhemorrhagic zones. Similar damaging effects were observed in many target organs in fish after administration of toxic substances such as aluminum (Poleo *et al.*, 1994, lead (Yousif, 1998), cadmium (Hellstron *et al.*, 2001), copper (Mourad and Wahby, 1999, Takasuki *et al.*, 2004 and Sukaty, 2010), phenol (Mohammed, 1995), hexavalent chromium (Mishra, 2008) and after exposure to radiation (Ergenjeva and Shagava, 2006).

Treatment with *S. crassifolia* attenuated Nimbecidin-induced histopathology in fish muscles. The pathological alterations were less proliferated in

muscles of fish treated for 4 weeks with Nimbecidin combined with 25% supplementary *S. crassifolia*. This protective effect of *S. crassifolia* could be attributed to its antioxidant activity which is time and dose dependant. Atencio *et al.* (2009) investigated the antioxidant role of selenium (Se) and its capability to repair tissue alterations induced by exposure to cyanobacterial cells in the liver, kidney, heart and gastrointestinal tract in Nile tilapia, *O. niloticus*. Other studies were carried out on the protective effect of some antioxidants against toxins in fish such as vitamin E (Prieto *et al.*, 2008), and calcium (Abdel-Tawab, 2005 and Sukaty, 2010). Yen and Duh (1994) showed that antioxidant properties of some flavonoids are effective mainly via scavenging of superoxide anion radical which is responsible for damaging of the **membranes of cell organelles**.

In conclusion, the results of the present work indicated that the brown alga *Sargassum crassifolia* had ameliorative effect against muscle damage induced by Nimbecidin and this may be mediated by its potent antioxidant activities which had the ability to restore the membranal structure and function of the cell organelles in the muscles of *O. niloticus*

#### References

1. Abdel-Tawab, M.; Khattab, Y.A.E.; Ahmad, A.H. and Shalaby, A.M.E. (2005): Compensatory growth, feed utilization, whole-body composition and hematological changes in starved juvenile Nile tilapia (*Oreochromis niloticus*) (L). *J. App. Aquacul.*, 18(3): 13-17.
2. Aebi, H.; CAT, I.N. and Bergmeyer, H.U.(1974). Methods of enzymatic analysis. Chemic Academic Press, Inc, Verlag, NY. 2: 673-685.
3. Al-Amoudi, O.A.; Mutawie, H.H.; Patel, A.V. and Blunden, G. (2009). Chemical composition and antioxidant activities of Jeddah corniche algae, Saudi Arabia. *Saudi J. Biol. Sci.* 16: 23-29.
4. Ananthi, S.; Raghavendran, H.R.B.; Sunil, A.G.; Gayathri, V.; Ramakrishnan, G. and Vasanthi, H.R.(2010). In vitro antioxidant and in vivo anti-inflammatory potential of crude polysaccharide from *Turbinaria ornata* (Marine Brown Alga). *Food Chem. Toxicol.* 48: 187-192.
5. AOAC.(1995). Association of Official Analytical Methods. Official Methods of Analysis. 16<sup>th</sup> Edition. Anlington, Verginia, USA.
6. Atencio, L.; Moreno, I.; Jos, Á.; Prieto, A.I.; Moyano, R.; Blanco, A. and Cameán, A.M. (2009). Effects of dietary selenium on the oxidative stress and pathological changes in tilapia (*Oreochromis niloticus*) exposed to a microcystin-producing cyanobacterial water bloom. *Toxicon* 53: 269-282.
7. Athukorala, Y.; Kim, K. and Jeon, Y. (2006). Antiproliferative and antioxidant properties of an enzymatic hydrolysate from brown alga, *Ecklonia cava*. *Food Chem. Toxicol.* 44: 1065-1074.
8. Avci, A.; Kaçmaz, M. and Durak, I. (2005). Peroxidation in muscle and liver tissues from fish in a contaminated river due to a petroleum refinery industry. *Ecotoxicol. Environ. Safety* 60: 101-105.
9. Bainy, A.C.D.; Saito, E.; Carvalho, P.S.M. and Junqueira, V.B.C. (1996). Oxidative stress in gill, erythrocytes, liver and kidney of Nile tilapia (*Oreochromis niloticus*) from a polluted site. *Aquat. Toxicol.* 34: 151-162.
10. Bradford, M.(1976). A rapid and sensitive method for the quantification of the microgram quantities of protein utilizing the principle of protein-dye binding. *Ann. Biochem.* 7: 248-254.
11. Burtin, P. (2003). Nutritional value of seaweeds. *Elec. J. Environ. Agric. Food Chem.* 2:498-503.
12. Chitra, K.C.; Latchoumycandane, C. and Mathur, P.P. (1999). Chronic-effect of endosulfan on the testicular function of rat. *Asian J. Androl.* 1: 203-206.
13. Dar, A.; Baig, H.S.; Saifullah, S.M.; Ahmad, V.U.; Yasmeen, S. and Nizamuddin, M. (2007). Effect of seasonal variation on the anti-inflammatory activity of *Sargassum wightii* growing on the N. Arabian Sea coast of Pakistan. *J. Exp. Mar. Biol. Ecol.* 351: 1-9.
14. Devi, P.K.; Suganthi, N.; Kesika, P. and Pandian, S.K. (2008). Bioprotective properties of seaweeds: In vitro evaluation of antioxidant activity and antimicrobial activity against food borne bacteria in relation to polyphenolic content. *BMC Complementary and Alternative Medicine* 8: 38.
15. Dorval, J.; Leblond, V.S. and Hontela, A. (2003). Oxidative stress and loss of cortisol secretion in adrenocortical cells of rainbow trout (*Oncorhynchus mykiss*) exposed in vitro to endosulfan, an organochloride pesticide. *Aquat. Toxicol.* 63: 229-241.
16. Dunkel, F.V. and Ricilards, D.C. (1998). Effect of an azadirachtin formulation on six non-target aquatic macroinvertebrates. *Environ. Entomol.* 27: 667-673.
17. Ehresmann, D.W.; Deig, E.F.; Hatch, M.T. and Verdros, V.A. (1977). Antiviral substances from California marine algae. *Phycologia* 13: 37-40.
18. Ellman, G. (1959). Tissue sulphhydryl groups. *Aech. Biochem. Biophys.* 82: 70-77.

19. El-Shenawy, N.S. (2010) : Effects of insecticides fenitrothion, endosulfan and abemectin on antioxidant parameters of isolated rat hepatocytes. *Toxicol. In Vitro.* **24(4): 1148-57**
20. Ergenjeva, T.P. and Shagava, M. (2006): Muscular tissue of bong fish inhabiting water bodies located in the 30 kilometer zone around the Chernobyl Nuclear Power Plant. *Doklady Akadememii Nauk*, **411(4): 567-569.**
21. Gabryelak, T. and Klekot, J. (1985): The effect of paraquat on the peroxide metabolism enzymes in erythrocytes of freshwater fish species. *Comp. Biochem. Physiol. C*, **81: 415-418.**
22. Ghadikolaei, K.R.; Abdulalian, E. and Wing-Keong, N.g. (2011). Evaluation of the proximate, fatty acid and mineral composition of representative green, brown and red seaweeds from the Persian Gulf of Iran as potential food and feed resources. *J. Food Sci. Technol.* In Press.
23. Harikrishnan, R.; Rani, M.N. and Balasundaram, C. (2003). Hematological and biochemical parameters in common carp, *Cyprinus carpio*, following herbal treatment for *Aeromonas hydrophila* infection. *Aquacult.* **221: 41-50.**
24. Hellström, L.; Elinder, C.-G. and Dahlberg, B. (2001): Cadmium exposure and end-stage renal disease. *Am. J. Kidney Dis.*, **38: 1001-1008.**
25. Heo, S.J, Park, E.J, Lee, K.W and Jeon, Y.J (2005): Antioxidant activities of enzymatic extracts from brown seaweeds. *Bioresour Technol* **96: 1613-1623**
26. Hopher, B.; Liao, I.C.; Cheng, S.I.I. and Hsieh, C.S. (1983). Food utilization by red tilapia. Effect of diet composition, feeding level and temperature on utilization efficiencies for maintenance and growth. *Aquacult.* **3(2): 255-275.**
27. Herbretau, F.; Coffard, L.J.M.; Derrien, A. and De Roeck-Holzharuer, Y. (1997). The fatty acid composition of five species of macroalgae. *Bot. Mar.* **40: 25-27.**
28. Hincal, F.; Gurbay, A. and Giray, B. (1995): Induction of lipid peroxidation and alteration of glutathione redox status by endosulfan. *Biol. Trace Elem. Res.*, **47: 321-326.**
29. Hussein, S.Y.; El-Nasser, A. and Ahmed, S.M. (1996). Comparative studies on the effects of herbicide atrazine on freshwater fish *Oreochromis niloticus* and *Chrysichthyes auratus* at Assiut Egypt. *Bull. Environ. Contam. Toxicol.* **57: 503-510.**
30. Iwai, K. (2008). Antidiabetic and antioxidant effects of polyphenols in brown alga *Ecklonia stolonifera* in Genetically Diabetic KK-A<sup>y</sup> Mice. *Plant Foods for Human Nutrition* **63: 163-169.**
31. Jee, J.H.; Masroor, F. and Kang, J.C. (2005). Responses of cypermethrin-induced stress in haematological parameters of Korean rockfish *Sebastes schlegeli* (Hildendorf). *Aquacult. Res.* **36: 898-905.**
32. Kang, J.Y.; Khan, M.N.A.; Park, N.H.; Cho, J.Y.; Lee, M.C.; Fujii, H. and Hong, Y.K. (2008). Antipyretic, analgesic, and anti-inflammatory activities of the seaweed *Sargassum fulvellum* and *Sargassum thunbergii* in mice. *J. Ethnopharm.* **116: 187-190.**
33. Karthikeyan, R.; Manivasagam, T.; Anantharaman, P.; Balasubramanian, T. and Somasundaram, S.T. (2011). Chemopreventive effect of *Padina boergeseni* extracts on ferric nitrilotriacetate (Fe-NTA)-induced oxidative damage in Wistar rats. *J. App. Phycol.* **23:257-263.**
34. Karthikeyan, R.; Somasundaram, S.T.; Manivasagam, T.; Balasubramanian, T. and Anantharaman, P. (2010). Hepatoprotective activity of brown alga *Padina boergeseni* against CCl<sub>4</sub> induced oxidative damage in Wistar rats. *Asian Pacific J. Tropical Med.* **3: 696-701.**
35. Kavitha, P. and Rao, J.V. (2009). Sub-lethal effects of profenofos on tissue-specific antioxidative responses in a Euryhaline fish, *Oreochromis mossambicus*. *Ecotoxicol. Environ. Safety* **72: 1727-1733.**
36. Kim, Y.C.; An, R.B.; Yoon, N.Y.; Nam, T.J. and Choi, J.S. (2005). Hepatoprotective constituents of the edible brown alga *Ecklonia stolonifera* on tacrine-induced cytotoxicity in hep G2 cells. *Arch. Pharm. Res.* **28: 1376-1380.**
37. Kukovinets, O.S. and Kislitsyn, M.I. (2006). Natural Arylterpenes and their biological activity. *Chem. Nat. Comp.* **42(1).**
38. Lourenço, S.O.; Barbarino, E.B.; De-Paula, J.C.; Pereira, L.Q.S. and Marquez, U.M.L. (2002). Amino acid composition, protein content and calculation of nitrogen-to-protein conversion factors for 19 tropical seaweeds. *Phycol. Res.* **50: 233-241.**
39. Martinez, C.B.R. and Souza, M.M. (2002). Acute effects of nitrite on ion regulation in two neotropical fish species. *Comp. Biochem. Physiol.* **133A: 151-160.**
40. Matanjun, P.; Mohamed, S.; Mustapha, N.M.; Muhammad, K. and Ming, C.H. (2008). Antioxidant activities and phenolics content of eight species of seaweeds from north Borneo. *J. App. Phycol.* **20: 367-373.**
41. Matanjun, P.; Mohamed, S.; Mustapha, N.M. and Muhammad, K. (2009). Nutrient content of tropical edible seaweeds, *Eucheuma cottonii*,



- Caulerpa lentillifera and Sargassum polycystum. J. App. Phycol. 21: 75-80.
42. Mishra, A.K. and Mohanty, B. (2008): Acute toxicity impacts of hexavalent chromium on behavior and histopathology of gill, kidney and liver of fresh water fish. (*Channa punctatus* Bloch). Environ. Toxicol. Pharmacol., 2: 136-141.
  43. Mohamed, F.A.B. (1995): Studies on some toxicological character and compound pathological symptoms of phenol on *Tilapia nilotica* and *Tilapia zillii*. M.Sc. Thesis, Depart of Zoology. Fac. Science, Cairo Univ
  44. Monteiro, D.A.; Almeida, J.A.; Rantin, F.T. and Kalinin, A.L. (2006). Oxidative stress biomarkers in the freshwater characid fish, *Brycon cephalus*, exposed to organophosphorus insecticide Folisuper 600 (methyl parathion). Comp. Biochem. Physiol. 143C: 141-149.
  45. Mourad, M. and Wahby, O. (1999): Physiological and histological changes in *Tilapia zillii* (Gerv) exposed to sublethal concentration of effluent of the Egyptian copper works Egypt. 5. Acta Ichthyolo. Piscatoria. 31: 73-79.
  46. Mousa, M.A.A.; El-Ashram, A.M.M. and Hamed, M. (2008): Effect of neem leaf extract on freshwater fishes and zooplankton community. 8<sup>th</sup> International Symposium on Tilapia in Aquaculture.
  47. Okhawa, H.; Ohishi, N. and Yagi, N. (1979): Assay for lipid peroxide in animal tissues by thiobarbituric acid reaction. Ann. Biochem., 5: 351-358.
  48. Omoregie, E. and Okpanachi, M.A. (1992): Growth of *Tilapia zilli* exposed to sublethal concentrations of crude extracts of *Azadirachta indica*. Acta Hydrobiol., 34: 281-286.
  49. Omoregie, E. and Okpanachi, M.A. (1997). Acute toxicity of water extracts of bark of the Neem plant, *Azadirachta indica* (Lodd) to the cichlid *Tilapia zilli* (Gervais). Acta Hydrobiol. 39: 47-51.
  50. Ortega-Calvo, J.J.; Mazuelos, C.; Hermosin, B. and Sáz-Jiménez, C. (1993). Chemical composition of *Spirulina* and eukaryotic algae food products marketed in Spain. J. Appl. Phycol. 5:425-435.
  51. Pandey, S.; Parvez, S.; Sayeed, I.; Haque, R.; Bin-Hafeez, B. and Raisuddin, S. (2003): Biomarkers of oxidative stress: a comparative study of river Yamuna fish *Wallago attu* (Bl. and Schn.). Sci. Total Environ., 309: 105-115.
  52. Pandey, S.; Parvez, S.; Ansari, R.A.; Ali, M.; Kaur, M.; Hayat, F.; Ahmad, F. and Raisuddin, S. (2008). Effects of exposure to multiple trace metals on biochemical, histological and ultrastructural features of gills of a freshwater fish, *Channa punctata* Bloch. Chemico-Biological Interactions 174: 183-192.
  53. Plaza, M.; Cifuentes, A. and Ibañez, E. (2008). In the search of new functional food ingredients from algae. Trends in Food Sci. Technol. 19: 31-39.
  54. Poleo, A.B.S.; Lydydersen, B.O.; Krooglund, F. and Salbo, B. (1994): Chemistry of mixing zones between limed streams and acidic tributaries. Water Air Soil Pollut., 75: 3339-3351.
  55. Prieto, A.I.; Jos, A.; Pichardo, S.; Moreno I. and Cameán, A.M. (2008). Protective role of vitamin E on the microcystin-induced oxidative stress in tilapia fish (*Oreochromis niloticus*). Environ. Toxicol. Chem. 27: 1152-1159.
  56. Prieto, A.I.; Jos, A.; Pichardo, S.; Moreno, I. and Cameán, A.M. (2006). Differential oxidative stress responses to microcystins LR and RR in intraperitoneally exposed tilapia fish (*Oreochromis* sp.). Aquat. Toxicol. 77: 314-321.
  57. Rábago-Castro, J.L.; Sanchez, J.G.; Pérez-Castañeda, R. and González-González, A. (2006). Effects of the prophylactic use of Romet®-30 and copper sulfate on growth, condition and feeding indices in Channel catfish (*Ictalurus punctatus*). Aquacult. 253: 343-349.
  58. Ruperez, P. (2002). Mineral content of edible marine seaweeds. Food Chem. 79(1): 23-26.
  59. Sayeed, I.; Parvez, S.; Pandey, S.; Bin-Hafeez, B.; Haque, R. and Raisuddin, S. (2003): Oxidative stress biomarkers of exposure to deltamethrin in freshwater fish, *Channa punctatus* Bloch. Ecotoxicol. Environ. Saf., 56: 295-301.
  60. Sánchez-Machado, D.I.; López-Hernández, J. and Paseiro-Losada, P. (2004). Fatty acids, total lipid, protein and ash contents of processed edible seaweeds. Food Chem. 85: 439-444.
  61. Sukaty, K.A. (2010). Biological and histological studies on the influence of water hardness on copper toxicity on Nile tilapia fish (*Oreochromis niloticus*): M.Sc. Thesis, Biology department. Fac. Appl. Science, Um Al-Qura Univ.
  62. Takasusuki, J.; Araujo, M.R.R. and Fernandes, M.N. (2004): Effect of water pH on copper toxicity in the neotropical fish, *Prochilodus scrofa* (Prochilodontidae). Bull. Environ. Contam. Toxicol., 72: 1075-1082.
  63. Van der Oost, R.; Beyer, J. and Vermeulen, N.P.E. (2003). Fish bioaccumulation and biomarkers in environmental risk assessment: a review. Environ. Toxicol. Pharm. 13: 149-157.
  64. Winkaler, E.U.; Santos, T.R.M.; Machado-Neto, J.G. and Martinez, C.B.R. (2007). Acute lethal and sublethal effects of neem leaf extract on the neotropical freshwater fish *Prochilodus*

- lineatus*. Comp. Biochem. Physiol. 145C: 236–244.
65. Yen, G.C. and Duh, P.D. (1994). Scavenging effect of methanolic extract of peanut hulls on free radical and active oxygen species. J. Agricult. Food Chem. 42: 629–632.
66. Yossif, G.A.R. (1998): A study of the biological changes of *Oreochromis niloticus* due to lead toxicity applications. M.Sc. Thesis Depart. Zoology. Fac. Science. Cairo Univ

11/25/2011