

## Weather in Oceania a in 2024

Gangadhara Rao Irlapati

H. No.5-30-4/1, Saibabanagar, Jeedimetla, Hyderabad - 500 055, Telangana, India.  
Email: gangadhar19582058@gmail.com

**Abstract:** The continent of Oceania, which has a globally textual history, is home to various climate change-induced natural disasters. Australia's weather and climate has continued to change, the State of the Climate 2024 report has found. The report found an increase in extreme heat events, longer fire seasons, more intense heavy rainfall, and sea level rise. Heavy rainfall events are becoming more intense. This paper examines the early warning systems for all Oceania and suggests how to study such climate changes and natural disasters, suggesting ways to anticipate them. Geological hazards such as earthquakes can be studied by developing the Geoscope system, By developing the Monsoon Time Scales, metrological hazards such as heavy rains and floods and droughts and famines can be studied. Plans can be made accordingly. The study of Cosmology can explore the inextricable links between planetary movements and disasters and how the planets orbiting in space are affecting the disasters that occur on earth. So Oceanian scientists can develop the Geoscope and Monsoon Time Scales as outlined below and protect people from climate changes and natural calamities.

[Gangadhara Rao Irlapati. **Weather in Oceania a in 2024.** *J Am Sci* 2025;21(7):38-111]. ISSN 1545-1003 (print); ISSN 2375-7264 (online). <http://www.jofamericanscience.org>. 04 doi:[10.7537/marsjas210725.04](https://doi.org/10.7537/marsjas210725.04)

**Keywords:** Bioforecast (1965-70); A New Hypothetical Model of Cosmology (1970-77); Inquisition (1977-79); Basics of Geoscope (1980-87); Basics of Monsoon Time Scales (1987-91); Indian Monsoon Time Scale (1991); Researches on Earth and space related issues (1991-2000); Numerical Weather Periodic Tables (2000-10); Designs of Geoscope projects (2010-20); Designs of Global Monsoon Time Scales (2020-)

### Introduction:

The following is a list of weather events that occurred on Earth in the year 2024. The several weather events which had a significant impact were blizzards, cold waves, droughts, heat waves, wildfires, floods, tornadoes, and tropical cyclones.

In 2024, the world experienced many natural disasters, including:

**Tropical cyclones:** The 2024 tropical cyclone season was marked by severe storms, including Hurricane Helene, which killed hundreds of people in the United States. The storms were fueled by high sea surface temperatures, which is a sign of climate change.

**Flooding:** There were floods in Southeast Asia, where at least 30 people died in Thailand and Malaysia, West and Central Africa, Europe, including Germany and Switzerland, South Asia, including Bangladesh, India, Nepal, The United States.

**Wildfires:** There were wildfires in Brazil, North America, including the United States, Mexico, and Canada

**Tornadoes:** There were tornado outbreaks in May 6–10, May 19–27.

There were other natural disasters also included a super typhoon in the Philippines, a super typhoon in Taiwan, a landslide in Enga, a cyclone in Remal, a heat wave in Pakistan.

The World Risk Report 2024 ranked India third among 193 countries most vulnerable to disasters

Earth-hazards can occur unexpected and cause damage to health, homes, and livelihoods. Hazards associated with earth processes such as avalanche, coastal flooding, cold waves, drought, earthquakes, hail, heat wave, tropical cyclones, ice storm, lightning, riverine flooding, strong wind, tornado, tsunami, volcanic activity, wildfire, winter weather, mudslides, landslides, rock slides and rock falls, soil creep, subsidence, floods, frost heave, coastal hazards, volcanic eruptions, earthquakes, glacial surges and outburst floods, tsunamis, and other land collapses and so many. I have done a lot of research on the Earth's disasters and proposed the Geoscope to study and predict geological hazards and Monsoon Time Scales to predict and study the meteorological hazards. All these disasters can be studied and predicted early through these two inventions. They are explained in detail in this paper. Along with these two, my A New Model of Cosmology is also explained below Because the planets have an inextricable connection with the calamities that occur on earth. Scientists have done further research and development on these and study and predict the climate changes and natural calamities in advance.

Scientists should note a few things here. Geoscope can be constructed either in the simple method or in the elaborate method. Similarly one should develop a complete understanding of the universe and the planetary movement. Due to this we can fully study

climate changes and natural calamities. Because the effect of planetary movements including sun and moon has a lot on the climate changes and natural calamities on the Earth. Along with the rest of the cosmological theories, my cosmology is also provides an understanding of the universe. But one thing should be noted regarding the construction of the Monsoon Time Scales. It's better a country establish its own Monsoon Time Scale to get 100% successful results. If not, it can establish the its regional monsoons time scale, as it has also reflects climate changes over the country. All these not possible to establish, then they may take up the Indian Monsoon Time Scale, which is successfully proved out in practice, and study the climate changes of the country. Because the Indian Monsoon Time Scale, far away, reflecting the climate changes of all world countries. Scientists should decide which of the above instruments can analyze their country's climate and develop it.

**A New Model of Cosmology:**

Planets have an inextricable connection with the climate changes and natural calamities ties that occur on earth. The sun, earth, and moon are held together by gravity, and they interact in lots of ways. The moon orbits the earth because of the pull of the earth. And the earth orbits the sun because of the pull of the Sun. The Sun, Moon, Earth, these three rotations around each other cause or create many climate changes and natural calamities on the Earth.

An example of planets influencing earthquakes is that several recent studies, however, have found a correlation between earth tides (caused by the position of the moon relative to the earth) and some types of earthquakes. One study, for example, concludes that during times of higher earth and ocean tides, such as during times of full or new moon, earthquakes are more likely on shallow thrust faults near the edges of continents and in (underwater) subduction zones.

An example of how asteroids affect Earth's climate is that the ITCZ oscillations on either side of the equators due to earth's revolution determine the hemisphere's seasons (mainly winter and summer), it is clear that earth's revolution plays a crucial role in the seasonal reversal of the prevailing surface winds observed in the regions where monsoons occur.

There are many such examples of planets being associated with calamities. So scientists should also have some knowledge of cosmology. Scientists study my cosmology as well as other theories.

According to the A New Hypothetical Model of Cosmology, the cosmos is made up of universes in infinite number, having similar universal external and internal and structure and properties, embedded one in each other and extended in ascending and descending

order. To explain and justify this model, there are three universes so far known to us (a) Geo-universe (b) Atomic-universe (c) Photon-universe. These three are having similar universal external and internal structure and properties, embedded one in each other and extended in ascending and descending order. Of these three, we known some extent about the internal structure and properties of the Geo-Universe but we do not known its external structure. We know some extent about the external structure and properties of the Photon-universe but we do not know its internal structure. Between of these three universes, we came to know a large extent about the internal & external structure and properties of the Atomic-universe. Hence, I have taken the similarities of external structure & properties between the Geo-universe & Atomic-universe to propose that all the universes in ascending and descending order of the creation are having similar universal internal structure and properties. The similarities of external structure & properties between the Atomic-universe and Photon-universe are taken to propose that all the universe in ascending and descending order of creation are having similar external structure and properties. And the manner in which of these three universes i.e., embedded one in each other, extended in ascending and descending order to propose that all the universes in ascending and descending order of the creation are embedded one in each other and extended in ascending and descending order. This doesn't mean that these photon, atom, universe etc. are arranged one on another as cycles separately. The cosmos enormous mixed compound of photons, atoms, universes etc. that are extended in ascending and descending order, embedded one in each other in the form of super matter or super fluid or super fluid matter.

## **2. Similar universal structure & properties:**

Of these three, we known some extent about the internal structure and properties of the Geo-universe but we do not know its external structure and properties. We know some extent about the external structure and properties of the Photon-universe but we do not know its internal structure and properties. Between of these three universes, we came to know a large extent about the internal and external, structure and properties of the Atomic-universe. So, I have taken the similarities of internal structure & properties between the Geo-universe & Atomic-universe to propose that all universes in ascending and descending order of the cosmos are having similar universal internal structure and properties. The similarities of external structure & properties between the Atomic-universe and Photon-universe are taken to propose that all the universes in ascending and descending order of

cosmos are having similar universal external structure and properties.

**Similar External Structure & Properties**

According to the model, all the universes in ascending and descending order of the creation are having similar external structure and properties. All the universes in either ascending or descending order of creation have the similar external structure and properties. So, we have imagine the external structure and properties of the atom compare with the external structure and properties of the photon. In the same way,

imagine the photon external structure and properties compare with the external structure and properties of the atom. Because, according to my cosmological principle all the universes in the ascending and descending order of creation must have similar external and internal structure and properties. To explain and justify this, I have taken many similarities between the atom and photon.

To justify this, I have taken many similarities between the atom and photon.

For example:-

Atomic-Universe	Photon-Universe
1) The atom appearing in several forms such as Hydrogen to uranium etc., being due to the Internal structure having different atomic particles at various numbers	2) The particle “Photon” related to energy appearing in several forms such as radio waves, gamma rays, violet rays etc being may be probably due to the internal structure having different particles at various numbers.
2)The atom exhibiting several physical and chemical Properties such as weight, colour, taste, hardness etc being due to the internal structure having different particles at various number.	)The particle “photon” related to energy exhibiting properties such as wave length colour, temperature etc being may be Probably due to the internal structure having different particles at various number.

**Similar Internal Structures & Properties**

According to the model, all the universes in ascending and descending order of the creation are having similar internal structure and properties. All the universes in either ascending or descending order of creation have the similar internal structure and properties. So, we have imagine the internal structure and properties of the atom compare with the internal structure and properties of the Geo-universe that's the universe seen around our earth. . In the same way, imagine the internal structure and properties of the Geo-universe, compare with the compare with the internal structure and properties of the atom. Because, according to my cosmological principle all the universes in the ascending and descending order of creation must have the similar external and internal and structure and properties. To explain and justify this, I have taken the many similarities between the atomic-universe and Geo-Universe.

Atomic-Universe	Geo-Universe
1)Various atomic particles at different sizes in several numbers are present in the atom	1) Various astronomical objects at different sizes in several numbers are present in the Geo- Universe
2) These atomic particles having three types of charges at negative, positive and neutral states are present in the atom	2) These astronomical objects having three type of charges at positive, negative and neutral states are present in the Geo-Universe
3) Positively charged protons are present in the nucleus	3) Stars built by atoms having positive charged nucleus are present in centre of the Geo-Universe
4) Neutrons at neutral state are present in the Nucleus.	4) Planets at neutral state are present in Centre of the Geo – Universe
5) Negatively charged electrons are present at large distance of the atomic nucleus in the	5) Here is a concept that anti-matter cosmic bodies built by atoms having negatively charged nucleus are present at large distance of the Geo-Universe.

atom	
6) Additional neutrons called isotopes are present.	6) Additional planets called satellites around the planets are present
7) Radiation emitting from the atom.	7) Cosmic rays emitting from the Geo- Universe.
8) There is a property of nuclear fission is in the atom.	8) There is a property of super Nova is in the Geo -Universe.

**Descending order of creation:**

The Geo-universe that means the Universe seen around our earth is having magnificent structure and properties such as galaxies, stars and planets and some planets such as earth having continents, countries, oceans, trees, animals. cyclones, human beings etc. Such Geo-universe being built by Universes of its descending order of creation that means atoms.

Atomic-universe that means the atom present in several forms from hydrogen to uranium etc is another gigantic universe, having magnificent structure and properties such as electrons, protons, neutrons, etc., and continents, countries, oceans, cyclones, trees, animals, human beings may be present on some neutrons having suitable conditions exactly similar to the earth planet resembling to the Geo-universe. Such Atomic universe being built by universes of its descending order of creation that means energy particle ‘photons’.

The Photon-universe that means the particle “photon” related to energy present in several forms of electromagnetic radiation is also another gigantic universe having magnificent structure and properties resembling to Geo-universe and atom. Such Photon-universe may also being built by universes of its descending order of creation that is not yet known to us.

Thus the descending order of creation continuous infinitely.

**Ascending order of creation:**

The Photon-universe that means the particle related to energy “photon” having magnificent structure and properties is being as a primary syntactic unit in the universe of its ascending order of creation that means atom. All components in the atom are built by these “photons” in infinite number. Such each and every energy particle “photon” in the Atomic-universe is basis to an infinite descending order of creation.

The Atomic—universe that means the “Atom” having magnificent structure and properties is being as a primary syntactic unit in the universe of its ascending order of creation that means in our Geo-universe. All components in the Geo-universe such as stars, planets etc., are built by these atoms in infinite number. Such each and every atom in the Geo-universe is basis to an infinite descending order of creation.

The Geo-Universe that means the “Universe” seen around our earth is a gigantic universe that is known

to us, having magnificent structure and properties is being as a primary syntactic unit in the universe of its ascending order of creation that is not yet known to us. All components in that universe are built by these Geo-universes in infinite number. Such each and every Geo-universe in that ascending creation is basis to an infinite descending order of creation.

Thus the ascending order of creation continuous infinite.

**Other justifications:**

The cosmological principle is a fundamental principle and assumption of cosmology stating that, on a large scale, the universe is both homogeneous and isotopic, in the words, the cosmological principle posits a relatively uniform universe.

The perfect cosmological principle is an extension of the cosmological principle, and states that the universe is homogeneous and isotropic in space and time. In this view the universe looks the same everywhere (on the large scale), the same way as it everywhere (on the large scale), the same as it always has and always will.

According to the universality of physical laws, all parts of the universe are subject to the same simple laws of nature that we find here on the earth, planets, stars, and galaxies move according to the same laws of gravity that governs the flight of a baseball. Light from distant galaxies reveals the same atomic and nuclear physics that we observe in our laboratories.

**Results and analysis:**

**Universal similarities:** According my theory, there are three universes so far known to us (a) Geo-Universe (b) Atomic-Universe (c) Photon-Univrs. These three are having similar structure and properties. Of these three, we known some extent about the internal structure and properties of the geo-niverse but we do not known its external structure. We know some extent about the external structure and properties of the photon-universe but we do not know its internal structure. Between of these three universes, we came to know a large extent about the internal & external structure and properties of the atomic-universe. Hence, I have taken the similarities of external structure & properties between the photon-universe & atomic-universe to propose that all the universes in ascending and descending order of the creation are having similar external structure and properties. The similarities of internal structure & properties between the atomic-universe and geo-universe are taken to propose that all

the universe in ascending and descending order of creation are having similar internal structure and properties.

***Uniform comparisons between atom and photon:***

The similarities of external structure & properties between the atom and photon are taken to propose that all these two are having similar internal structure and properties.

***Structure:*** The Atom appearing in several forms such as hydrogen to uranium etc., being due to the internal structure having different atomic particles at various number. In the same manner the “photon” related to energy appearing in several forms such as radio waves, gamma rays, violet rays etc being may be probably due to the internal structure having different particles at various numbers.

***Properties:*** The atom exhibiting several physical and chemical properties such as weight, colour, taste, hardness etc being due to the internal structure having different particles at various number. The “photon” related to energy exhibiting properties such as wave length colour, temperature etc being may be probably due to the internal structure having different particles at various number.

Various atomic particles at different sizes in several numbers are present in the atom Various astronomical objects at different sizes in several numbers are present in the Geo- Universe.

***Uniform comparisons between Atom and Geo-universe:*** The similarities of intertural structure & properties between the atom and geo-universe are taken to propose that all these two are having similar internal structure and properties.

1. Various atomic particles at different sizes in several numbers are present in the atom. In the similar way various astronomical objects at different sizes in several numbers are present in the geo- universe. 2. These atomic particles having three types of charges at negative, positive and neutral states are present in the atom. In the similar way, these astronomical objects having three type of charges at positive, negative and neutral states are present in the geo-universe. 3. Positively charged protons are present in the nucleus. In the similar way, Stars built by atoms having positive charged nucleus are present in centre of the Neutrons at neutral state are present in the nucleus. In the similar way, planets at neutral state are present in centre of the geo-universe. 5. Negatively charged electrons are present at large distance of the atomic nucleus in the atom. In the similar way, there is a concept that anti-matter cosmic bodies built by atoms having negatively charged nucleus are present at large distance of the geo-universe. 6. Additional neutrons called isotopes are present. In the similar way, additional planets called satellites around the planets are present. 7.

Radiation emitting from the atom. In the similar way, cosmic rays emitting from the geo-universe. 8. There is a property of nuclear fission in the atom. In the similar way, there is a property of super Nova in the geo-universe.

**Study and discussion:**

The Cosmology is one of the most creative and bizarre areas of science, concerned with the studies of origin, structure, nature and evolution of the universe. There are two main theories, steady state theory and the big bang theory, that explain the structure of the universe. For example, The big bang theory has faced many criticisms by many scientists as being inadequate to explain the relativity and complexity of the universe. Therefore, it not sufficient to correctly model the origins of the universe.

According to Bud Rapanault (quora); ‘The Big Bang Theory is essentially unscientific because the physical model it presents does not resemble the cosmos we observe in any of its particulars. None of the distinguishing features of the Big Bang Theory are part of the cosmological landscape that lies before us. The Big Bang Theory itself and the *ad hoc* inflationary epoch are unobservable by terms of the model.

Curved and expanding space time cannot be directly detected but are integral to the model.

The Big Bang Theory model requires that 95% of the universe consist of some dark matter and dark energy neither of which can be empirically detected and both of which are simply additional adhoc patches necessary to make the model predictions conform to physical reality.

In addition, the Big Bang Theory rests on two assumptions, one simplistic and naïve, the other dubious

The cosmos is a unified, coherent, and simultaneous entity.

The cosmological redshift is a recessional velocity.

According to George Yool(quora); current evidence like the cosmological principle, hubble ultra deep field and alternatives like quantum relativity suggest a universe has no beginning or end in which big bangs are galactic processes we can observe empirically. There are many esteemed critics such as; NASA WMAP beyond big bang theory; Einstein evolving universe.

Hoyle The big bang theory got its name from a man who thought the theory was total nonsense.

Plus 34 more famous scientists around the world in an open joint letter to the scientific community has been criticized the Big Bang Theory ( Big Bang Theory Busted By 33 Top Scientists)Rense.com

**Doctrines:**

After many researches and studies on the origin, structure, nature and evolution of the cosmos, I proposed many doctrines.

**Geo-universe:** Geo-universe, that means the universe seen around our earth is a gigantic universe, having magnificent structure and properties such as galaxies, stars, planets etc. There are continents, oceans, countries and living beings on some planets in our geo-universe. Such gigantic geo-universe is a little atom in its ascending world. That means our universe is in that ascending world just as atom is in our universe. We do not know whether our geo-universe is in the air, water or elsewhere in that ascending world. In my opinion, our Geo-universe is in an intelligent creature in that ascending world. But what is not known to that creature in that ascending world is that there are galaxies, stars, planets and human worlds in their atoms. In their view it is thought that there are tiny particles in their atoms. But they do not know that there are galaxies, stars, planets and human worlds in their atoms just as we think that there are electrons, protons, neutrons in our atom. But we do not realize that there are galaxies, stars, planets and human worlds in the atom. So do they think they have little particles. Such geo-universe being built by universes of its descending order of creation and the same time being act as a primary syntactic unit in its ascending order of creation that is not yet known to us.

**Atomic universe :** Atomic-universe, that mean the atom present from hydrogen to uranium etc. is a gigantic universe, having similar universal structure and properties such as galaxies, stars, planets in the form of electrons, protons and neutrons exactly similar to our universe. Just as there are continents, oceans, countries and human beings on some earth-like planets in our Geo-universe, continents, oceans, countries and living beings can be present on some neutrons in the atom. That is the atom is a vast universe as our universe. But we do not recognize them. For the creatures in the atom, in their view the atom is a great universe but they do not know that another gigantic universe exists above them, and that their atom in which they resides is acts as a tiny atom in our universe just as we do not know that our gigantic universe is a small particle in the universe above us. There may be humans on neutrons who have civilizations like countries, governments, planes, trains, ships, motors etc. just there are humans who have civilizations on our planet. Such atoms being built by universes of its descending order of creation that are photons and the same atom being act as a primary syntactic unit in its ascending world that is our universe..

**Photon-universe:** Photon-universe, that mean photon, the particle related to energy existing in several forms of electromagnetic radiation etc. is a gigantic universe,

having similar universal structure and properties similar to our universe that have galaxies, stars, planets and similar to atom that have electrons, protons and neutrons. Just as there are continents, oceans, countries and human beings on some planets in our Geo-universe, continents, oceans, countries and living beings can be present on some planets in the photons. That is, the photon is a vast universe as our universe and as the atom. But we do not recognize them. For the creatures in the photon, in their view their photon is a great universe but they do not know that another there is a gigantic universe in their ascending world that is atom, and there is another world above the atom that is our universe. There may be humans in the photon who have civilizations like countries, governments, planes, trains, ships, motors etc. just there are humans who have civilizations on our planet. Such photon is being built by universes of its descending order of creation that are not yet known to us and the same photon is being act as a primary syntactic unit in its ascending world that is atom.

#### **Basics of Geoscope projects:**

Many researches and studies were conducted by me between 1980-1987 and Basics of Geoscope & its projects were proposed and designed by me in 1987 for all world regions and countries in 1987 with many good eminence intentions and ambitions intended to study and research the earth's underground and surface matters for public purposes with many proposals i.e to take and keep the entire underground to be under the control of National Geoscope System/National Geoscope Projects to study the underground mysteries; explore the underground resources; increasing artificial underground waters by attracting the sea waters to the areas of deserts through layers by electro-ionization; create artificial rains by attracting vaporized sea waters to the desert plains through the sky by geo-magnetizing atmosphere when the weather is surrounded by water molecules during the trough or low pressure areas, create artificial storms and making them our control by moving desert planes and pour rains; restore and recreate people in past by images that are preserved in the earth's magnetic field by new technology Geo-Machine and study geological resources by constantly studying the National Geoscope System/National Geoscope Projects. This is not what Buckminster had proposed in 1962 and many similar other architectures in the name of Geoscope. My invention is completely different and proposed with good eminence intentions as mentioned above.

There is nowhere on Earth that's immune from quakes but a few places are far less likely to have one. Qatar is one such country and there are a few others,

including Norway, Finland and Sweden. These Nordic countries rarely have quakes. Of all the continents, Antarctica has faced the least earthquakes. Though no place is completely safe from earthquakes, Qatar is considered to be the country with earthquakes. The Arabian plate, which includes Saudi Arabia, is an entirely separate plate. And Saudi Arabia does not even collide with any other fault lines. Because it does not coincide with any of the other plates or even separated from some of the earth's fault lines, Saudi Arabia is left largely untouched by the earthquake.

#### **Construction:**

Geoscope means- a mechanical architecture established in between the underground and observatory with the help of bore-well proposed for conducting geological studies to know the earthquakes, ores and water currents etc.

A borehole having suitable width and depth has to be dug in the earthquake prone areas. An observatory having research & analysis facilities has to be constructed on the borehole. Apparatus & sensors to recognize the geo-physical and geo-chemical changes generated in the underground such as foreshocks, chemical changes, electrogeopulses, micro-vibrations, pressure, geomagnetic forces etc should be inserted into the underground and linked with the concerned analysis sections of the observatory that is above the ground to study the changes taking place in the underground.

That means-relative results of geological & geographical researches & developments of past, present and future should be interposed, coordinated and constantly developed. The apparatus related to the geology and geography such as Richter scale etc also should be set in the observatories of the Geoscope. we can make many more modern ideas & modifications thus bringing many more improvements & developments in the Geoscope.

Many kinds of super high remote sensing technology in the area of sensor physics, signal processing used specially image processing, electromagnetic detection technology etc should be used in the Geoscope. Geophysical deep underground detectors and mineral exploration equipments, natural gas sensors etc should be used in the Geoscope. Electromagnetic sensors may also be used in the

#### **Materials and Methods:**

A borehole having suitable width and depth has to be dug in the earthquake prone area. An observatory having the most modern high-tech research facilities has to be constructed on that bore-well. Most modern mechanical systems like electronic, physical and chemical sensors and apparatus to recognize the underground physical and chemical conditions such as the underground mineral resources,

rise and fall of the underground water levels, micro-vibrations and waves generated in the underground, differences in pressure, temperature and other seismic activities in the underground should be inserted into the underground and linked with the concerned research and study departments of the observatory that is above the bore-well to research and study the conditions and changes taking place in the underground. The results of researches of the geophysical and geological sciences just like Richter scale etc., also should be setup in the Geo-scope. Many kinds of super high remote sensing technology in the area of sensor physics, signal processing used specially image processing, electromagnetic detection technology etc should be used in the Geo-scope. Geophysical deep underground detectors and mineral exploration equipments, natural gas sensors etc should be used in the Geo-scope. Electromagnetic sensors may also be used in the Geo-scope project.etc. That means relative results of geological & geophysical researches & developments of past, present and future should be interposed, coordinated and constantly developed. We can make many more modern ideas & modifications thus bringing many more improvements & developments in the Geo-scope.

#### **Types of geoscopes:**

Geoscope can be built in many types and various forms just like Simple Geoscope Model, Home-Made Geoscope Model and Modern Geoscope Model. Simple Geoscope Model is having simple construction involving no expenditure that is a deep well having suitable width and depth has to be dug. Construct a room over the well. Wash the inner walls of the room with white lime. Fix an ordinary electric bulb in the room. That is enough. Home-made Geoscope is also very simple and easy construction involves no expenditure moreover even students, children's and science enthusiasts can make the Home-made Geoscope and detect the earth-quakes 24 to 28 hrs in advance. By making certain changes and alterations, a house having a well can be converted into a Geoscope i.e., wash the inner walls of that house with white lime. Fix ordinary electric bulbs in the room. The Home-made Geoscope is complete. Both these two are very easy methods. Besides these two methods, Micro-Geoscope is an elaborate construction. It is a modern technology system consisting of surface laboratory and underground research facilities. For this model a deep bore-well having suitable width and depth has to be dug. A surface laboratory having the most modern high-tech underground research facilities has to be constructed on that bore-well to study, analyze and recognize the underground conditions. Underground research apparatus should be inserted into the underground and

linked with the concerned research and study departments of the laboratory that is above the bore-well to research and study the conditions and changes taking place in the underground.

**Simple geoscope method:** This is a simple construction involving no expenditure. A deep well having suitable width and depth has to be dug in the earthquake prone area. Construct a room over the well. Wash the inner walls of the room with white Lime. Fix an ordinary electric bulb in the room.

Observe the colour of the room lighting daily. When the bulb glows, the light in room generally appears white in colour, but before occurrence of an earthquake, the room lighting turns blue in colour. The onset of earth-quake can be guessed by this “Seismic luminescence Emission”

**Principle:** Due to stress of continental plates and some other reasons on a place where there are favourable chances for earth-quake to occur, the pressure is induced in the underground. As a result, there is a steady rise in the pressure around the focus centre. Because of the large disparity in the magnitude of energies involved, gas anomalies such as (a) Helium emission (b) Chemico-seismic anomalies such as sulphur, calcium, nitrogen etc., chemical compounds (c) Seismic atomic radiations of radioactive mineral compounds such as radon show up much earlier even at large distance from the epic-centre which enter the well through the underground springs. These gas anomalies occupy the room in this manner; emit radiation which gives ultraviolet blue colour (sometimes red) to the room.

**Home-made geoscope method:** This construction involves no expenditure. Even students, children’s and science enthusiasts can make the Home-Made Geoscope and detect the earth-quakes 24 to 28 hrs in advance. By making certain changes and alterations, the houses in the earthquake prone area having a well can be converted into a Geoscope i.e., wash the inner walls of the house with white Lime fix ordinary electric bulbs in the room.

Observe the colour of the room lighting in the house daily. When the bulb glows, the light in room generally appears white in colour, but before occurrence of an earthquake, the room lighting turns blue in colour. The onset of earth-quake can be guessed by this “Seismic luminescence Emission”

**Principle:** Due to stress of continental plates and some other reasons on a place where there are favourable chances for earth-quake to occur, the pressure is induced in the underground. As a result, there is a steady rise in the pressure around the focus centre. Because of the large disparity in the magnitude of energies involved, gas anomalies such as (a) Helium emission (b) Chemico-seismic anomalies such as

sulphur, calcium, nitrogen etc., chemical compounds (c) Seismic atomic radiations of radioactive mineral compounds such as radon show up much earlier even at large distance from the epic-centre which enter the well through the underground springs. These gas anomalies occupy the room in this manner; emit radiation which gives ultraviolet blue colour (sometimes red) to the room.

**Modern geoscope method:** A borehole having suitable width and depth has to be dug into the underground in the above earthquake prone area. A surface laboratory having the most modern high-tech underground research facilities has to be constructed on that bore-well to research and study the conditions and changes taking place in the underground. Electronic, physical and chemical sensors and apparatus, super high remote sensing technology in the area of sensor physics, signal processing used specially image processing ,electromagnetic detection technology, deep underground detectors and mineral exploration equipments, natural gas sensors, electromagnetic sensors etc to recognize the underground physical and chemical conditions such as the underground mineral resources, rise and fall of the underground water levels, micro-vibrations and waves generated in the underground, differences in pressure, temperature and other seismic activities in the underground etc should be inserted into the underground and linked with the concerned research and analyze departments of the above surface underground research laboratory that is above the bore-well to analyze the conditions and changes taking place in the underground. That means researches & developments of past, present and future should be interposed, coordinated and constantly developed. We can make many more modern ideas& modifications thus bringing many more improvements & developments in the Geoscope.

**Management:** Observe the geophysical & geochemical changes such as foreshocks, chemical changes, ground water levels, strain in rocks, thermal anomalies, seismic-luminescence gas anomalies, electrogeopulses, micro-vibrations, pressure, geomagnetic forces, etc taking place in the underground. The onset of earthquakes can be guessed by analyzing the aforesaid studies in the concerned analysis sections of the laboratory that is above the well.

#### **Central data processing center:**

In this Geoscope system, there should be established Local Geoscope centers and Central Data Processing Centre in the above earthquake prone area for managing the system in a coordinated manner.

One or more required number of Geoscopes should be established in the above earthquake prone area. The

observation personnel in the respective Geoscope centers should watch the onset of earthquakes day and night.

There should be established a Central Data Processing Centre to co-ordinate and codify the information supplied by the Local Geoscope Centres of the earthquake prone area in a coordinated manner.

Whenever any Local Geoscope Centre sends warning about the onset of earthquakes, the observation personnel should immediately send the information to its central data processing centre. The central data processing centre analyze the information supplied by the local geoscope centre and estimates the epicentre, time, area to be affected urban places etc., details of the impending earthquake and send to the authorities, and media and warnings in advance to take precautions.

#### **Results and analysis:**

Many investigations were carried out and successfully proved out in practice. The risk of earthquakes in Andhra Pradesh is less but the source is greater in north India and other regions in the world including the earthquake prone area the establishment of the Geoscope is very useful to study and predict the earthquakes. Among them, electrogeogram test is one that's thought to be the heartbeat of the underground. Similarly, the study of the luminescent phenomena, electromagnetic emission and light radiation, thermo-luminescence and fracto-mechanoluminescence are others. Several researches and studies have been conducted as described above and obtained many key results.

**Seismicluminescence study:** Gas anomalies emission: Over the centuries, there have been many reports of earthquake lights, both before and while the ground is shaking.

Most rock contain small amounts of gases that can be isotopically distinguished from the normal atmospheric gases. There are reports of spikes in the concentrations of such gases prior to a major earthquake; this has been attributed to release due to pre-seismic stress or fracturing of the rock. One of these gases is radon, produced by radioactive decay of the trace amounts of uranium present in most rock. Radon is useful as a potential earthquake predictor because it is radioactive and thus easily detected, and its short-half life makes radon levels sensitive to short-term fluctuations. The earthquakes with which these changes are supposedly linked were up to a thousand kilometers away, months later, and not at a magnitudes. In some cases the anomalies were observed at a distant site, but not at closer sites.

And, the lights are caused by electrical properties of certain rocks. The earthquake lights can take many different shapes, forms, and colors. Common forms of

earthquake lights include bluish flames that appear to come out of the ground at ankle height; orbs of light called ball lightning that float in the air for tens of seconds or even minutes; and quick flashes of bright light that resemble regular lightning strikes, except they come out of the ground instead of the sky and can stretch up to 200 meters. When nature stresses certain rocks, electric charges are activated. The lights can occur hours to days before major earthquakes and also during actual shaking. They have been recorded at distance of up to 160 kilometers from the epicenter. Earthquake lights are likely to be very helpful with earthquake prediction. To study seismic luminescence Geoscope can be built in many forms just like Simple geoscope model, Home-made geoscope model and Modern geoscope model etc.

Construct the simple geoscope should be placed in the earthquake prone area described above to study the seismic luminescence as follows. This is a simple model involving no expenditure. A well having suitable width and depth has to be dug. Construct a room over the well. Wash the inner walls of the room with white Lime. Fix an ordinary electric bulb in the room.

Construct home-made geoscope should be placed in the earthquake prone area described above to study the seismic luminescence as follows. This is also very simple and easy model involves no expenditure. Even students, children's and science enthusiasts can make the Home-Made Geoscope and detect the earth-quakes 24 to 28 hrs in advance. By making certain changes and alterations, a house having a well can be converted into a Geoscope i.e., wash the inner walls of that house with white Lime. Fix ordinary electric bulbs in the room.

The two Geoscope structures described above are easy to construct, easy to use and easy to analyze the Seismic luminescence study. Observe the colour of the room lighting daily. When the bulb glows, the light in room generally appears white in colour, but before occurrence of an earth-quake, the room lighting turns ultra violet blue in colour. The onset of earth-quake can be guessed by this "Seismic luminescence emission"

In modern methods to analyze the seismic luminescence, a deep bore-well having suitable width and depth has to be dug in the earthquake prone areas. A laboratory having most modern high-technological research and analysis facilities including a mechanical system to analyze the seismic luminescence and gas anomalies emerging from underground has to be constructed on that well. All types of modern sensors and apparatus including a mechanical system to catching/grabbing/absorbing the seismic luminescence or gas anomalies emerging from the

underground to recognize the seismic luminescence and other seismic activities should be inserted into the underground and linked with the concerned research analyzing sections of the laboratory that is above the well to observe, study, research and analyze the seismic luminescence and seismic changes existing and taking place in the underground. By that earthquakes can be warned by analyzing the luminescence as given the above.

Observe the fracto luminescence gas anomalies existing and taking place in the underground. The onset of earthquakes can be guessed by analyzing the aforesaid seismic luminescence studies in the concerned analysis sections of the laboratory that is above the well.

Due to stress of continental plates and some other reasons on a place where there are favourable chances for earth-quake to occur, the pressure is induced in the underground. As a result, there is a steady rise in the pressure around the focus centre. Because of the large disparity in the magnitude of energies involved, gas anomalies such as shown below show up much earlier even at large distance from the epicentre which enter the well through the underground springs.

(a) Emission of Helium, Hydrogen etc

(b) Emission of chemico-seismic evaporation anomalies such as sulphur, calcium, nitrogen etc., ,

(c) Emission of seismic atomic radiations such as radon from radioactive mineral compounds etc

These gas anomalies occupy the room in this manner; emit radiation which gives blue colour (sometimes red) to the room.

Collect and analyze the above mentioned gas anomalies and seismic luminescence in the concerned section established in laboratory that is above the well. Study the gas anomalies and seismic luminescence in the research and analysis sections of the Geoscope daily 24 hours 365 days. When the gas anomalies or seismic luminescence are released the earthquakes can be considered.

Here is a very important is to be grasped. Before occurring of an earthquake, gas anomalies as stated above such as radon, helium, hydrogen and chemico-mineral evaporations such as sulphur, calcium, nitrogen and other fracto-luminescence radiations show up earlier even at large distances from the epicentre due to stress, disturbances, shock waves and fluctuations in the underground forces. These gas anomalies & fracto luminescence radiations and other chemical evaporations enter into the well through the underground springs. When these anomalies occupy the simple Geoscope rooms or Home-made Geoscope rooms above the well, the room lighting turns violet in colour. The light in the room scattered in the presence of these gas anomalies, fracto-luminescence radiations

and other chemico-mineral evaporations the ultra violet radiation is emitted more and the room lighting turns in violet colour. Our eye catches these variations in the radiation of the lighting in the room easily since- The violet rays having smaller wave length

The violet rays having property of extending greatly

The light becoming weak in the violet region

The eyes having greater sensitivity to violet radiation

Due to all these reasons, the room may appear violet in colour then we can predict the impending earth quakes 12 hours in advance. This principle is also applies to the section built in modern research and analysis methods that is above the well

**Electrogeogram Test:** This is also easy study to recognize the impending earth quake. A borehole having suitable width and depth has to be dug in the earthquake prone area.

An earth wire or rod should be inserted into the underground by the borehole and linked with the concerned analysis section having apparatus to detect, compare measure of the electric currents of the electric circuit of the earth systems. Otherwise by observing the home electric fans.etc. We can also study the electrogeopulses studies to predict the impending earth quake.

Observe the changes in the electric currents of the earth system 24 hours, 365 days. From a power station, the electricity is distributed to the far-off places. Normally the circuit of the power supply being completed through the earth system. Whenever if the disturbances occurs in the layers of the earth's underground, the fluctuation rate will be more due to the earth quake obstructions such as pressure, faults, vibrations, water currents etc., of the earth's underground. So we can forecast the impending earth quake by observing the obstruction of electric currents of circuit of the earth system in the observatory of the Geoscope and also by the obstruction sounds in the electric fans etc.

#### **Study and discussion:**

Many studies and experiments have been carried out on the Geoscope project and all were successfully proved out in practice. And also several designs have been proposed to study and explore the underground. The risk of earthquakes in Andhra Pradesh is less but the source is greater in North India and other regions in the world including the earthquake prone area where the establishment of the Geoscope is very useful.

#### **Applications:**

Geoscope is to detect natural calamities such as earthquakes etc. as well as underground resources. Along with these, I have also made some proposals just like artificial rains to another new earth in the space based on the Geoscope. Their details are given below.

By setting up the National Geoscope Project in and maintain, that country can be predicted the impending earthquakes, volcanic hazards (and storm surges, tsunamis etc consequence secondary hazards due to the earthquakes occur in the womb that means underground of the sea or ocean if the country have the chances of occurring of these disasters) in advance. And also the country can be predicted mineral and underground resources by inserting many kinds of super high remote sensing technology in the area of sensor physics, signal processing used specially image processing electromagnetic detection technology and geophysical deep underground detectors and mineral exploration equipments, natural gas sensors etc in the underground through the Geoscope. Setting up the National Geoscope Project and maintain will also be useful in emerging industries such as geothermal and geo-sequestration etc.

Geoscopes should be designed in the possible coastal areas where tsunamis are likely to occur. A tsunami or tidal wave, also known as a seismic sea wave, is a series of enormous waves in displacement of a large volume of water body caused by the earthquakes, underground landslides, volcanic eruptions, asteroids generally in an ocean or a large lake. Tsunamis can travel 20-30 miles per hour with waves 10-100 feet high. The effects of tsunamis are devastating. Tsunami damage is first caused by the immense force of the tidal wave hitting the shoreline. I conducted some studies on the tsunamis. Some studies have been conducted by me on the tsunamis to study and predict the tsunamis and designed the Geoscope in 1987 to keeping the tsunamis. Geoscope should be designed in the coastal areas of the sea and earthquakes and its consequent secondary hazards such as tidal forces, rogue waves, tsunami can be predicted by virtue of performing studies as described above. Geoscope is very useful in studying, predicting and mitigating the tsunamis and its dangers.

Geoscopes should be designed in the possible areas where landslides are likely to occur and the earthquakes and its secondary consequent hazards such as landslides mud slides, mass movements, sink holes, coastal erosion, lahars, mud flows, etc can be estimated by virtue of performing studies as described above.

Geoscopes should be designed in the volcano areas and volcanic activities such as volcanic gases, and steam generated eruptions, explosive eruption of high – silica lava, effusive eruption of low-silica lava, debris flow and carbon dioxide emission etc can be predicted by virtue of performing studies as described above. Let's discuss about some of the key studies.

By setting up the National Geoscope projects and maintain, a country can be predicted the impending

earthquakes, volcanic hazards (and storm surges, tsunamis etc consequence secondary hazards due to the earthquakes occur in the womb that means underground of the sea or ocean if the country have the chances of occurring of these disasters) in advance. And a country can be predicted mineral and underground resources by inserting many kinds of super high remote sensing technology in the area of sensor physics, signal processing used specially image processing electromagnetic detection technology and geophysical deep underground detectors and mineral exploration equipments, natural gas sensors etc in the underground by using the Geoscope.

Setting up the National Geoscope Project and maintain will also be useful in emerging industries such as geothermal and geo-sequestration etc.

Geoscope projects can be built where the earthquakes are likely to occur and study the earthquakes.

Build Geoscope in the seismic areas and earthquakes can be predicted by virtue of performing studies as described above.

#### **Basics of Monsoon Time Scale:**

There are many mysteries and unsolved issues in the monsoonal climate and Weather systems that cannot explain and solve. According to the researches and studies on the Monsoon Time Scales, it is known that there will be major global climate changes in the coming years "i.e" heavy rains, floods and storms etc. will occur until about 2075 and there will be droughts and famines etc. until about 2150. Through the establishment of Monsoon Time Scales, we can know the future consequences of the climate changes. Plans can be made accordingly. I call on world scientists to design and establish the Monsoon Time Scale following the Basics of Monsoon Time Scales outlined below, based on the India Monsoon Time Scale which is successfully proved out in practice and break down the mysteries of the Indian monsoon.

Each region of the world can establish monsoon time scales for their respective regions. Accurate results can only be obtained if the monsoon time scale belonging to their regions are obtained. For example, it is better if the Canada country establish its Canada Monsoon Time Scale. If not, countries can set up regional Monsoon Time scales belonging to their respective regions. For example, countries in the North American continent can establish the North American Time Scale. If these are not possible to establish, then they can set up the Indian Monsoon Time Scale and study the climate changes of their countries. Because the Indian Monsoon Time Scale, far away, reflects climate changes in distant all world regions.

By establishing the Monsoon Time Scale and maintain, a country can be estimated the impending weather conditions and natural calamities such as monsoon

movements, rains, floods, landslides, avalanches, blizzard, droughts, famines extreme winter conditions, heavy rainfall, mudflows, extreme weather, storms, cloud burst, sand storms, hails and winds etc all climate, meteorological and weather related conditions & natural calamities in advance. Surface water resources can also still be found. We can make separate monsoon time scales per each and every individual country. As a part of this, I have proposed and designed Basics of Monsoon Time Scales for all countries separately.

After much research, I have proposed some basics regarding method and design to prepare a country's Monsoon Time Scale as outlined below.

**Method and Design:**

**Design:** Prepare a Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of a country's Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed into a square graphic scale. This scale should be designed in three ways i.e Basic scale, Filled scale, Analyzed scale;

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** This is the second scale that is filled with data and explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is scientifically analyzed the filled scale by data, it explains monsoon patterns weather conditions of the scale.

**Method:** There are two methods in formation and process of the Monsoon Time Scales. The first one is in the single form and next one is designed in four parts.

**Single & Full length Scale:** Prepare the Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of a country's Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed in a single and full length type square graphic scale. It can be formed on a Paper or a Wall or a Table.

**Parts & paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is beginning from 1<sup>st</sup> April to July 12<sup>th</sup>.  
The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.  
And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

These separate scales can be pasted into one scale as explained below.

Cut along the edges of dates on the right side of the first part and paste it to along the edges of date of 13<sup>th</sup> July on left side of the second part.

Cut along the edges of dates on the right side of the second part and paste it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and paste it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part .

When paste this manner, we get long full-length Monsoon Time Scale.

**Computerization:**

Monsoon Time Scales can also be computerized. Besides rather than in manual type scale, if we are able to create a computer model scale which to be the most obvious.

**Material and Data:**

Construction of the Monsoon Time Scales requires enormous data of low pressure systems, depressions tropical cyclones/storms, snowfall and sand storms etc. that formed over and affecting a region should be taken as data to prepare the Monsoon Time Scale. An accurate scale is available if we can collect and analyze the exact climate data.

What should the data be taken?

For example, countries where monsoon occur should taken low pressure systems as data.

Countries where storms occur can be taken storms as data.

European countries can taken the Westerlies as data.

Snowy countries of polar climate can take snowfall, snowy rains, graupel, snowpellets as data.

Desert or hot climate countries can take sand or dust storm incidents as data.

Scientists can also be taken yearly climate changes as a key data as every year occurs routinely in their countries.

**Management:**

The main weather events such as monsoon pulses in the form of low pressure systems if any of a monsoon region formed over a region or country have been entering on the scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds should be entered on the Monsoon Time Scale as per date and month of each and every year. If we can managing the scale in this manner continuously, we can study the past, present and future movements of monsoons of a region or country. I took the numbers to analysis the

variations in data. Researchers have to decide what kind of data to take and how to analyze the data.

**Researches and results:**

The research and study should be done in the same way as described below in the Indian Monsoon Time Scale and the results should be obtained.

**Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

**Indian Monsoon Time Scale:**

I have undertaken the Indian Monsoon Time Scale as the model scale following all the rules of Basics of Monsoon Time Scales. The reason I took the Indian Monsoon Time Scale as the model research was because I was in the Indian monsoon region. I know the information about Indian monsoon very well.

The Indian Monsoon Time Scale is a chronological sequence of events arranged in between time and weather with the help of a scale for studying past's, present and future movements of the monsoon of India and its relationship with rainfall and other weather problems and natural calamities. From where to wherever to be taken the time and weather data to analyze, the researcher can decide on his discretion according to available weather data.

**Method and design:**

**Design:** For this, I took a period of 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of India's as the time and the data of monsoonal low pressure systems, depressions and storms of 139 years from 1880 to 2027 that were formed over the Indian region taken as the climate, on the whole comprising of a large time and climate took and framed into a square graphic scale. I designed this scale in three ways i.e Basic scale, Filled scale, Analyzed scale as described below.

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** The second one is filled by data scale, it explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is filled and analyzed by data, it explains monsoon patterns of the scale.

**Method:** There are three methods used to design this scale. The first one is the single and full length scale and second one is parts & past scale. The last one is computer model made entirely by computer system.

**Single & Full length Scale:** I prepared the Indian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or

according to the chronology of India's time and climate) of 139 year from 1880 to 2027 or a required period, comprising of a large time and climate was taken and framed in a single and full length type square graphic scale. It can be formed on a paper, board, wall or table.

**Parts & Paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is from 1<sup>st</sup> April to July 12<sup>th</sup>.

The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.

And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

These separate scales are pasted into one scale as described below below.

Cut along the edges of dates on the right side of the first part and pasted it to along the edges of date of 13th July on left side of the second part.

Cut along the edges of dates on the right side of the second part and pasted it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and pasted it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part.

When pasted in this manner, we get long full length Indian Monsoon Time Scale

**Computer model scale:**

Besides this above two manual scales, I have prepared a computer Indian Monsoon Time Scale generated by the computer system from the year 1888 to 1983 for the period of 1<sup>st</sup> June to September 30<sup>th</sup>. If we are able to create a computer model scale which to be the most obvious.

**Material & data:** The monsoon pulses in the form of low pressure systems over the Indian region have been taken as the data to the construction of this scale. For this, a lot of enormous data of low pressure systems, depressions and cyclones that formed over the Indian region were taken as the climate from many resources just like Mooley DA, Shukla J(1987); characteristics of the west ward-moving summer monsoon low pressure systems over the Indian region and their relationship with the monsoon rainfall. Centre for Ocean-land Atmospheric interactions, University of Maryland, college park, MD., and from many other resources and from many other resources just like The world's 7 Tropical Cyclone seasons around the world etc.

**Management:**

The monsoon pulses in the form of low pressure systems over the Indian region are taken and

entered on the scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds pertaining to the date and month of the each and every year. How the Indian monsoons have been travelling for the last 140 years since 1880 onwards are recorded on the Indian Monsoon Time Scale. I took the numerical/statistical method to analysis the variations in data. If we have been managing the scale in this manner continuously, we can study the past, present and future movements of monsoon of India. Researchers have to decide what kind of data to take and how to analyze the data.

### **Results&analysis:**

I did comprehensive researches on the Indian Monsoon Time Scale and analyzed many key mysteries related to the monsoonal system. The Indian Monsoon Time Scale reveals many secrets and mysteries of the Indian monsoon and its relationship with movement of axis of the Earth around the Sun in the universe & its influences on the Earth's atmosphere. Let's study the mystery of the Indian monsoon and discuss the rest of other features of the Indian Monsoon Time Scale later.

When examine the scale, I noticed that several passages or path-ways of monsoon pulses it have been some cut-edge paths and splits passing through its systematic zigzag cycles in a systematic manner in parallel and stacked next to each other in ascending and descending order clearly seen on the Indian Monsoon Time Scale. If the thin arrows along the passages identified on the Indian Monsoon Time Scale are drawn from 1880 to the current year, then the monsoon paths appears. Many other methods can analyze the Indian Monsoon Time Scale. In my researches I have noticed that depending on the incidence of heavy rains & floods in some years and droughts & famines in another years were happened according to the travel of monsoon path. The path of monsoon when travelling over four months from June to September good rainfall or heavy rains and floods were occurred. And the path when travelling over last months i.e July or August or September, low rainfall and droughts were occurred. Particularly, there are two main passages. The first one is main path or passage of the Indian monsoon(Southwest monsoon) and the second one is path or passage of the north-east monsoon. The first one is on the left side over the months of June, July, August, September(southwest monsoon) and another path on the right side over the months of October, November, December are visible in the Indian Monsoon Time Scale

### **Pre-path of Indian monsoon:**

Keep track the Indian Monsoon Time Scale carefully. When we look at the Indian Monsoon Time

Scale, several paths appears. Two of these are important. These can be called main path of the Indian monsoon and pre-path of the main passage of the Indian monsoon. The main path appears clear and its pre-path appears unclear. Due to unavailability of data, it is not known how the pre-path of the Indian monsoon traveled before 1888. But according the studies-

Between 1727-1751 years, it traveled in the shaped of concave direction for about 24 years and caused low rainfall and droughts in many years.

Between 1752-1811 years, it traveled in the shape of convex direction for about 60 years and caused good rainfall and floods in many years.

Between 1812-1835 years, it traveled in the shape of concave direction for about 25 years and caused low rainfall and droughts in many years.

Between 1836-1895 years, it traveled in the shaped of convex direction for about 60 years and caused good rainfall and floods in many years.

Between 1896-1919 years, it traveled in the shape of concave direction for about 24 years and caused low rainfall and droughts in many years.

Between 1920-1981 years, it traveled in the shape of convex direction for about 62 years and caused good rainfall and floods in many years.

Between 1982-2009 years, it traveled in the shape of concave direction for about 27 years and caused low rainfall and droughts in many years.

From 2010, it is going to travel upwards in the shape of convex direction for 56 years that's until 2066 and will be resulting good rainfall and floods in the coming years.

### **Main-path of Indian monsoon:**

Keep track the Indian Monsoon Time Scale carefully. During the 1865-1895's, the main path-way of the Indian monsoon was rising over June, July, August. During 1896-1920's, it was falling over August, September. During 1920-1965's, it was rising again over July, August, September. During 1965-2020s, it was falling over September. From 2020, it is now rising upwards and estimated traveling over the months of June, July, August by the 2060.

Due to unavailability of data, it is not known how the main path of the Indian monsoon traveled before 1888. But according the studies, it is known that it traveled in the shape of convex direction for 56 years between 1865-1897 and caused good rainfall in many years. During this 4 months period of (June, July, August, September) of Indian monsoon season, the line of path of the monsoon was travelled over all these four months. As a result, there were heavy rains and floods in most years.

From 1898 to 1920, the line of path of the Indian monsoon was travelled over the months of

August and September in the shape of concave direction. In this 4 months monsoon season, the line was travelled just over two months only. As a result, it rained only two months instead of four months monsoon season and caused low rainfall in many years,

From 1920 to 1964, the line of path of the Indian monsoon was travelled over the months of July, August and September in the shape of convex direction. In this 4 months monsoon season, the line was travelled over three months. As a result, it rained only three months instead of four months monsoon season and resulted good rainfall in more years.

From 1965 to 2020, the passage of the Indian monsoon was travelled over the months of August to mid-august in the shape of deep sloping direction, In this 4 months monsoon season, the line was travelled just over two months for a short period only. As a result it rained only two months instead of four months monsoon season. and caused low rainfall and droughts in many yearcavF

From 2020, the line of path of the Indian monsoon seems likely rising over the months of July and to June in future in the shape of upper ascending direction and will be resulting heavy rains & floods in coming years during 2020-2066. This is an assessment based on the study of situations from 1888. As per new analysis-

Between 1727-1751 years, it traveled in the shaped of concave direction for about 24 years and caused low rainfall and droughts in many years.

Between 1752-1811 years, it traveled in the shape of convex direction for about 60 years and caused good rainfall and floods in many years.

Between 1812-1835 years, it traveled in the shape of concave direction for about 25 years and caused low rainfall and droughts in many years.

Between 1836-1895 years, it traveled in the shaped of convex direction for about 60 years and caused good rainfall and floods in many years.

Between 1896-1919 years, it traveled in the shape of concave direction for about 24 years and caused low rainfall and droughts in many years.

Between 1920-1981 years, it traveled in the shape of convex direction for about 62 years and caused good rainfall and floods in many years.

Between 1982-2009 years, it traveled in the shape of concave direction for about 27 years and caused low rainfall and droughts in many years.

From 2010, it is going to travel upwards in the shape of convex direction for 56 years that's until 2056 and will be resulting good rainfall and floods in the coming years.

#### **Study&discussion:**

The results obtained as above are studied and discussed as follows.

The Indian Monsoon Time Scale reveals many other secrets of the monsoon & its relationship with rainfall & other weather problems and natural calamities. Some bands, clusters and paths of low pressure systems clearly seen in the Indian Monsoon Time Scale, it have been some cut-edge paths passing through its systematic zigzag cycles in ascending and ascending orders which causes heavy rains & floods in some years and droughts & famines in another years according to their travel. And also we can find out many more secrets of the Indian monsoon such as droughts, famines, cyclones, heavy rains, floods, onset & withdrawal of monsoon etc. by keen study of the Indian Monsoon Time Scale. The passages clearly seen in the Indian Monsoon Time Scale are sources of monsoon pulses. The tracking date of main path & other various paths of monsoon etc., of the Indian Monsoon denotes the onset of the monsoon, monsoon pulses or low pressure systems. These observations can mean that pulses of the monsoon are repeatedly determined by the number of repeats.

Furthermore example, the main passage of line of monsoon travel from June to September and September to June are also signs to impending weather conditions of a country. For example, during 1865-1895's, ConAAe main path-way of the Indian monsoon was rising over June, July, August. During 1896-1920's, it was falling over August, September. During 1920-1965's, it was rising again over July, August, September. During 1965-2020s, it was falling over September. From 2020, it is now rising upwards and estimated traveling over the months of June, July, August by the 2066.

(There may be a difference of 5 to 10 or more years between those periods. This is because currently it can not be estimated with certainty that the respective period will start or end in the ruling period.)

The tracking date of main path & other various paths of the Indian Monsoon denotes the onset of the monsoon, monsoon pulses or low pressure systems, storms and its consequent secondary hazards and storms etc.. And also we can find out many more secrets of the Indian monsoon such as droughts, famines, cyclones, heavy rains, floods, real images of the Indian Monsoon, and onset & withdrawals of the monsoon etc. by keen study of the Indian Monsoon Time Scale.

For example, the date of tracking ridge of path is the sign to the impending cyclone and its secondary consequent hazard floods, storm surges etc.,

Another example, the thin and thick markers on the upper border line of the Indian Monsoon Time Scale are the signs to the impending heavy rains & floods

and droughts & floods. The thick marking of clusters of low pressure systems on the Indian Monsoon Time Scale is the sign to the impending heavy rains and floods and the thin marking of clusters of low pressure systems on the Indian monsoon time scale is the sign to the impending droughts and famines.

These are just some studies of the Indian monsoon. There are many more secrets in the Indian monsoon. Indian scientists should get rid of them. We can find out many more secrets of weather conditions by keen study of the Indian Monsoon Time Scale.

#### **Basics of Australian Monsoon Time Scale:**

The Australian summer Monsoon is traditionally referred to as the wet season in Northern Australia when over three Quarters of the annual rainfall occurs. The Australian summer Monsoon is just a portion of the greater Australian-Indonesian Monsoon that extends from the equator to about 15°S and Westward from 100°E to about 155°E the greater. The rainfall season occurs from September to February and it is a major source of energy for the Hadley circulation during boreal winter, This is also known as Indo-Australian Monsoon and the Australian Monsoon may be considered to be the same system, the Indo-Australian Monsoon.

Also known as the Indo-Australian Monsoon. The rainy season occurs from September to February and it is a major source of energy for the Hadley circulation during boreal winter. The *Maritime Continent Monsoon* and the *Australian Monsoon* may be considered to be the same system, the Indo-Australian Monsoon.

It is associated with the development of the Siberian High and the movement of the heating maxima from the Northern Hemisphere to the Southern Hemisphere. North-easterly winds flow down Southeast Asia, are turned north-westerly/westerly by Borneo topography towards Australia. This forms a cyclonic circulation vortex over Borneo, which together with descending cold surges of winter air from higher latitudes, cause significant weather phenomena in the region.

The onset of the monsoon over the Maritime Continent tends to follow the heating maxima down Vietnam and the Malay Peninsula (September), o Sumatra, Borneo and the Philippines (October), to Java, Sulawesi (November), Irian Jaya and Northern Australia (December, January). However, the monsoon is not a simple response to heating but a more complex interaction of topography, wind and sea, as demonstrated by its abrupt rather than gradual withdrawal from the region. The Australian monsoon (the "Wet") occurs in the southern summer when the monsoon trough develops over Northern Australia. Over three-quarters of annual rainfall in Northern Australia falls during this time.

The Australian Monsoon Time Scales is a chronological sequences of events arranged in between Time and Climate with the help of a scale for studying the past's, present and future movements of the Australian monsoon regions and its relationship with rainfall and other weather problem and natural calamities.

Prepare the Australian Monsoon Time Scale having 365 horizontal days from March 21<sup>st</sup> to next year March 20<sup>th</sup> or a required period comprising of a large time and climate have been taken and framed into a square graphic scale.

The main weather events if any of the Australian monsoon region such as low pressure systems, depressions and storms/cyclones etc have been entering on the Australian Monsoon Time Scale as per date and month of each and every year.

If we have been managing the Australian Monsoon Time Scale in this manner continuously, we can see the image and its past's, present's and future movements of the Australian monsoon and study it's originals, climatic changes and futuristic dimensions. By establishing the Australian Monsoon Time Scales which can help to study the movements of the the Australian monsoon.

#### **Method and Design:**

**Design:** Prepare a Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Australian Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed into a square graphic scale.

This scale should be designed in three ways i.e Basic scale, Filled scale, Analyzed scale;

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** This is the second scale that is filled with data and explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is scientifically analyzed the filled scale by data, it explains monsoon patterns weather conditions of the scale.

**Method:** There are two methods in formation and process of the Australian Monsoon Time Scales. The first one is in the single form and next one is designed in four parts.

**Single & Full length Scale:** Prepare the Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup>

or according to the chronology of Australian's Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed in a single and full length type square graphic scale. It can be formed on a paper, board, wall or a Table.

**Parts & paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is beginning from 1<sup>st</sup> April to July 12<sup>th</sup>.<sup>5</sup>

The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.

And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

These separate scales can be pasted into one scale as explained below.

Cut along the edges of dates on the right side of the first part and paste it to along the edges of date of 13<sup>th</sup> July on left side of the second part.

Cut along the edges of dates on the right side of the second part and paste it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and paste it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part.

When paste this manner, we get long full-scape Australian monsoon Time Scale.

#### **Computer Model:**

Australian Monsoon Time Scales can also be established as a computer model. Besides rather than in manual type scale, If we are able to create a computer model scale which to be the most obvious.

#### **Material and Data:**

Construction of the Australian Monsoon Time Scales requires enormous data of low pressure systems, depressions, tropical storms, sand storms etc that affecting a region and formed over a region should be taken as data to prepare the Australian Monsoon Time Scale. An accurate scale is available if we can collect and analyze the exact climate data.

**Management:** The main weather events if any of Australian monsoon such as monsoon pulses in the form of low pressure systems if any of a monsoon region formed over the Australian monsoon have been entering on the Australian Monsoon Time Scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds as per date and month of each and every year. If we can managing the scale in this manner continuously, we can study the past, present and future movements of Australian monsoon. I took

the numbers to analysis the variations in data. Researchers have to decide what kind of data to take and how to analyze the data.

#### **Researches&results:**

The study should be done in the same way as described in the Indian Monsoon Time Scale and the results should be obtained.

#### **Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

#### **Basics of Western North Pacific Monsoon Time Scale:**

The Western North Pacific Monsoon Time Scales is a chronological sequences of events arranged in between the Time and Climate with the help of a scale for studying the past's, present and future movement of the Western North Pacific monsoon regions and its relationship with rainfall and other weather problem and natural calamities.

Prepare the Western North Pacific Monsoon Time Scale having 365 horizontal days from March 21<sup>st</sup> to next year March 20<sup>th</sup> or a required period comprising of a large time and climate have been taken and framed into a square graphic scale.

The main weather events if any of the Western North Pacific monsoon region such as low pressure systems, depressions and storms/cyclones etc have been entering on the Western North Pacific Monsoon Time Scale as per date and month of each and every year.

If we have been managing the Western North Pacific Monsoon Time Scale in this manner continuously, we can see the image and its past's, present's and future movements of the Western North Pacific monsoon and study it's originals, climatic changes and futuristic dimensions.

By establishing the Western North Pacific Monsoon Time Scale which can help to study the movements of the the Western North Pacific monsoon.

#### **Method and Design:**

**Design:** Prepare a Western North Pacific Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Western North Pacific Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed into a square graphic scale.

This scale should be designed in three ways i.e Basic scale, Filled scale, Analyzed scale;

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** This is the second scale that is filled with data and explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is scientifically analyzed the filled scale by data, it explains monsoon patterns weather conditions of the scale.

**Method:** There are two methods in formation and process of the Western North Pacific Monsoon Time Scales. The first one is in the single form and next one is designed in four parts.

**Single & Full length Scale:** Prepare the Western North Pacific Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Western North Pacific's Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed in a single and full length type square graphic scale. It can be formed on a paper, board, wall or a Table.

**Parts & paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is beginning from 1<sup>st</sup> April to July 12<sup>th</sup>.

The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.

And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

These separate scales can be pasted into one scale as explained below.

Cut along the edges of dates on the right side of the first part and paste it to along the edges of date of 13<sup>th</sup> July on left side of the second part.

Cut along the edges of dates on the right side of the second part and paste it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and paste it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part.

When paste this manner, we get long full-scape Western North Pacific Monsoon Time Scale.

#### **Computer Model:**

Western North Pacific Monsoon Time Scale can also be established as a computer model. Besides rather than in manual type scale, If we are able to create a computer model scale which to be the most obvious.

#### **Material and Data:**

Construction of the Western North Pacific Monsoon Time Scale requires enormous data of low pressure systems, depressions, tropical storms, sand storms etc that affecting a region and formed over a region should be taken as data to prepare the Western North Pacific Monsoon Time Scale. An accurate scale is available if we can collect and analyze the exact climate data.

**Management:** The main weather events if any of Western North Pacific monsoon such as monsoon

pulses in the form of low pressure systems if any of a monsoon region formed over the Western North Pacific monsoon have been entering on the Western North Pacific Monsoon Time Scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds as per date and month of each and every year. If we can managing the scale in this manner continuously, we can study the past, present and future movements of Western North Pacific monsoon. I took the numbers to analysis the variations in data. Researchers have to decide what kind of data to take and how to analyze the data.

#### **Researches & results:**

The study should be done in the same way as described in the Indian Monsoon Time Scale and the results should be obtained.

#### **Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

#### **Basics of North-Australian Monsoon Time Scale:**

The North-Australian Monsoon Time Scales is a chronological sequences of events arranged in between Time and Climate with the help of a scale for studying the past's, present and future movements of the North-Australian monsoon regions and its relationship with rainfall and other weather problem and natural calamities.

Prepare the North-Australian Monsoon Time Scale having 365 horizontal days from March 21<sup>st</sup> to next year March 20<sup>th</sup> or a required period comprising of a large time and climate have been taken and framed into a square graphic scale.

The main weather events if any of the North-Australian monsoon region such as low pressure systems, depressions and storms/cyclones etc have been entering on the North-Australian Monsoon Time Scale as per date and month of each and every year.

If we have been managing the North-Australian Monsoon Time Scale in this manner continuously, we can see the image and its past's, present's and future movements of the North-Australian monsoon and study it's originals, climatic changes and futuristic dimensions.

By establishing the North-Australian Monsoon Time Scales which can help to study the movements of the the North-Australian monsoon.

#### **Method and Design:**

**Design:** Prepare a North-Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of North-Australian Time and Climate) of 139 year from 1880 to 2027

comprising of a large Time and Climate should be taken and framed into a square graphic scale.

This scale should be designed in three ways i.e Basic scale, Filled scale, Analyzed scale;

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** This is the second scale that is filled with data and explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is scientifically analyzed the filled scale by data, it explains monsoon patterns weather conditions of the scale.

**Method:** There are two methods in formation and process of the North-Australian Monsoon Time Scales. The first one is in the single form and next one is designed in four parts.

**Single & Full length Scale:** Prepare the North-Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of North-Australian's Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed in a single and full length type square graphic scale. It can be formed on a paper, board, wall or a Table.

**Parts & paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is beginning from 1<sup>st</sup> April to July 12<sup>th</sup>.

The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.

And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

These separate scales can be pasted into one scale as explained below.

Cut along the edges of dates on the right side of the first part and paste it to along the edges of date of 13th July on left side of the second part.

Cut along the edges of dates on the right side of the second part and paste it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and paste it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part .

When paste this manner, we get long full-scape North-Australian Monsoon Time Scale.

**Computer Model:**

North-Australian Monsoon Time Scales can also be established as a computer model. Besides rather than in manual type scale, If we are able to create a computer model scale which to be the most obvious.

**Material and Data:**

Construction of the North-Australian Monsoon Time Scales requires enormous data of low pressure systems, depressions, tropical storms, sand storms etc that affecting a region and formed over a region should be taken as data to prepare the North-Australian Monsoon Time Scale. An accurate scale is available if we can collect and analyze the exact climate data.

**Management:** The main weather events if any of North-Australian monsoon such as monsoon pulses in the form of low pressure systems if any of a monsoon region formed over the North-Australian monsoon have been entering on the North-Australian Monsoon Time Scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds as per date and month of each and every year. If we can managing the scale in this manner continuously, we can study the past, present and future movements of North-Australian monsoon. I took the numbers to analysis the variations in data. Researchers have to decide what kind of data to take and how to analyze the data.

**Researches & results:**

The study should be done in the same way as described in the Indian Monsoon Time Scale and the results should be obtained.

**Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

**Basics of Indo-Australian Monsoon Time Scale:**

The Indo-Australian Monsoon Time Scales is a chronological sequences of events arranged in between Time and Climate with the help of a scale for studying the past's, present and future movements of the Indo-Australian monsoon regions and its relationship with rainfall and other weather problem and natural calamities.

Prepare the Indo-Australian Monsoon Time Scale having 365 horizontal days from March 21<sup>st</sup> to next year March 20<sup>th</sup> or a required period comprising of a large time and climate have been taken and framed into a square graphic scale.

The main weather events if any of the Indo-Australian monsoon region such as low pressure systems, depressions and storms/cyclones etc have been entering on the Indo-Australian Monsoon Time Scale as per date and month of each and every year.

If we have been managing the Indo-Australian Monsoon Time Scale in this manner continuously, we can see the image and its past's, present's and future

movements of the Indo-Australian monsoon and study its originals, climatic changes and futuristic dimensions.

By establishing the Indo-Australian Monsoon Time Scales which can help to study the movements of the the Indo-Australian monsoon.

#### **Method and Design:**

**Design:** Prepare a Indo-Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Indo-Australian Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed into a square graphic scale.

This scale should be designed in three ways i.e Basic scale, Filled scale, Analyzed scale;

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** This is the second scale that is filled with data and explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is scientifically analyzed the filled scale by data, it explains monsoon patterns weather conditions of the scale.

**Method:** There are two methods in formation and process of the Indo-Australian Monsoon Time Scales. The first one is in the single form and next one is designed in four parts.

**Single & Full length Scale:** Prepare the Indo-Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Indo-Australian's Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed in a single and full length type square graphic scale. It can be formed on a paper, board, wall or a Table.

**Parts & paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is beginning from 1<sup>st</sup> April to July 12<sup>th</sup>.

The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.

And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

These separate scales can be pasted into one scale as explained below.

Cut along the edges of dates on the right side of the first part and paste it to along the edges of date of 13<sup>th</sup> July on left side of the second part.

Cut along the edges of dates on the right side of the second part and paste it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and paste it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part .

When paste this manner, we get long full-scape Indo-Australian Monsoon Time Scale.

#### **Computer Model:**

Indo-Australian Monsoon Time Scales can also be established as a computer model. Besides rather than in manual type scale, If we are able to create a computer model scale which to be the most obvious.

#### **Material and Data:**

Construction of the Indo-Australian Monsoon Time Scales requires enormous data of low pressure systems, depressions, tropical storms, sand storms etc that affecting a region and formed over a region should be taken as data to prepare the Indo-Australian Monsoon Time Scale. An accurate scale is available if we can collect and analyze the exact climate data.

**Management:** The main weather events if any of Indo-Australian monsoon such as monsoon pulses in the form of low pressure systems if any of a monsoon region formed over the Indo-Australian monsoon have been entering on the Indo-Australian Monsoon Time Scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds as per date and month of each and every year. If we can managing the scale in this manner continuously, we can study the past, present and future movements of Indo-Australian monsoon. I took the numbers to analysis the variations in data. Researchers have to decide what kind of data to take and how to analyze the data.

#### **Researches&results:**

The study should be done in the same way as described in the Indian Monsoon Time Scale and the results should be obtained.

#### **Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

#### **Basics of Borneo-Australian Monsoon Time Scale:**

The Borneo-Australian Monsoon Time Scales is a chronological sequences of events arranged in between Time and Climate with the help of a scale for studying the past's, present and future movements of the Borneo-Australian monsoon regions and its

relationship with rainfall and other weather problem and natural calamities.

Prepare the Borneo-Australian Monsoon Time Scale having 365 horizontal days from March 21<sup>st</sup> to next year March 20<sup>th</sup> or a required period comprising of a large time and climate have been taken and framed into a square graphic scale.

The main weather events if any of the Borneo-Australian monsoon region such as low pressure systems, depressions and storms/cyclones etc have been entering on the Borneo-Australian Monsoon Time Scale as per date and month of each and every year.

If we have been managing the Borneo-Australian Monsoon Time Scale in this manner continuously, we can see the image and its past's, present's and future movements of the Borneo-Australian monsoon and study it's originals, climatic changes and futuristic dimensions.

By establishing the Borneo-Australian Monsoon Time Scales which can help to study the movements of the the Borneo-Australian monsoon.

#### **Method and Design:**

**Design:** Prepare a Borneo-Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Borneo-Australian Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed into a square graphic scale.

This scale should be designed in three ways i.e Basic scale, Filled scale, Analyzed scale;

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** This is the second scale that is filled with data and explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is scientifically analyzed the filled scale by data, it explains monsoon patterns weather conditions of the scale.

**Method:** There are two methods in formation and process of the Borneo-Australian Monsoon Time Scales. The first one is in the single form and next one is designed in four parts.

**Single & Full length Scale:** Prepare the Borneo-Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Borneo-Australian's Time and Climate) of 139 year from 1880 to 2027

comprising of a large Time and Climate should be taken and framed in a single and full length type square graphic scale. It can be formed on a paper, board, wall or a Table.

**Parts & paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is beginning from 1<sup>st</sup> April to July 12<sup>th</sup>.

The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.

And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

**These separate scales can be pasted into one scale as explained below.**

Cut along the edges of dates on the right side of the first part and paste it to along the edges of date of 13<sup>th</sup> July on left side of the second part.

Cut along the edges of dates on the right side of the second part and paste it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and paste it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part .

When paste this manner, we get long full-scape Asian Australian Monsoon Time Scale.

#### **Computer Model:**

Borneo-Australian Monsoon Time Scales can also be established as a computer model. Besides rather than in manual type scale, If we are able to create a computer model scale which to be the most obvious.

#### **Material and Data:**

Construction of the Borneo-Australian Monsoon Time Scales requires enormous data of low pressure systems, depressions, tropical storms, sand storms etc that affecting a region and formed over a region should be taken as data to prepare the Borneo-Australian Monsoon Time Scale. An accurate scale is available if we can collect and analyze the exact climate data.

**Management:** The main weather events if any of Borneo-Australian monsoon such as monsoon pulses in the form of low pressure systems if any of a monsoon region formed over the Borneo- Australian monsoon have been entering on the Borneo-Australian Monsoon Time Scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds as per date and month of each and every year. If we can managing the scale in this manner continuously, we can study the past, present and future movements of Borneo-Australian monsoon. I took the numbers to analysis the variations in data. Researchers have to

decide what kind of data to take and how to analyze the data.

#### **Researches&results:**

The study should be done in the same way as described in the Indian Monsoon Time Scale and the results should be obtained.

#### **Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

#### **Basics of Australian Indonesian Monsoon Time Scale:**

The Australian Indonesian Monsoon Time Scales is a chronological sequences of events arranged in between Time and Climate with the help of a scale for studying the past's, present and future movements of the Australian Indonesian monsoon regions and its relationship with rainfall and other weather problem and natural calamities.

Prepare the Australian Indonesian Monsoon Time Scale having 365 horizontal days from March 21<sup>st</sup> to next year March 20<sup>th</sup> or a required period comprising of a large time and climate have been taken and framed into a square graphic scale.

The main weather events if any of the Australian Indonesian monsoon region such as low pressure systems, depressions and storms/cyclones etc have been entering on the Australian Indonesian Monsoon Time Scale as per date and month of each and every year.

If we have been managing the Australian Indonesian Monsoon Time Scale in this manner continuously, we can see the image and its past's, present's and future movements of the Australian Indonesian monsoon and study it's originals, climatic changes and futuristic dimensions.

By establishing the Australian Indonesian Monsoon Time Scales which can help to study the movements of the the Australian Indonesian monsoon.

#### **Method and Design:**

**Design:** Prepare a Australian Indonesian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Australian Indonesian Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed into a square graphic scale.

This scale should be designed in three ways i.e Basic scale, Filled scale, Analyzed scale;

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** This is the second scale that is filled with data and explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is scientifically analyzed the filled scale by data, it explains monsoon patterns weather conditions of the scale.

**Method:** There are two methods in formation and process of the Australian Indonesian Monsoon Time Scales. The first one is in the single form and next one is designed in four parts.

**Single& Full length Scale:** Prepare the Australian Indonesian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Australian Indonesian's Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed in a single and full length type square graphic scale. It can be formed on a paper, board, wall or a Table.

**Parts & paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is beginning from 1<sup>st</sup> April to July 12<sup>th</sup>.

The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.

And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

These separate scales can be pasted into one scale as explained below.

Cut along the edges of dates on the right side of the first part and paste it to along the edges of date of 13th July on left side of the second part.

Cut along the edges of dates on the right side of the second part and paste it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and paste it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part .

When paste this manner, we get long full-scape Australian Indonesian Monsoon Time Scale.

#### **Computer Model:**

Australian Indonesian Monsoon Time Scales can also be established as a computer model. Besides rather than in manual type scale, If we are able to create a computer model scale which to be the most obvious.

#### **Material and Data:**

Construction of the Australian Indonesian Monsoon Time Scales requires enormous data of low pressure

systems, depressions, tropical storms, sand storms etc that affecting a region and formed over a region should be taken as data to prepare the Australian Indonesian Monsoon Time Scale. An accurate scale is available if we can collect and analyze the exact climate data.

**Management:** The main weather events if any of Australian Indonesian monsoon such as monsoon pulses in the form of low pressure systems if any of a monsoon region formed over the Australian Indonesian monsoon have been entering on the Australian Indonesian Monsoon Time Scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds as per date and month of each and every year. If we can managing the scale in this manner continuously, we can study the past, present and future movements of Australian Indonesian monsoon. I took the numbers to analysis the variations in data. Researchers have to decide what kind of data to take and how to analyze the data.

#### **Researches&results:**

The study should be done in the same way as described in the Indian Monsoon Time Scale and the results should be obtained.

#### **Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

#### **Basics of Asian Australian Monsoon Time Scale:**

The Asian Australian Monsoon Time Scales is a chronological sequences of events arranged in between Time and Climate with the help of a scale for studying the past's, present and future movements of the Asian Australian monsoon regions and its relationship with rainfall and other weather problem and natural calamities.

Prepare the Asian Australian Monsoon Time Scale having 365 horizontal days from March 21<sup>st</sup> to next year March 20<sup>th</sup> or a required period comprising of a large time and climate have been taken and framed into a square graphic scale.

The main weather events if any of the Asian Australian monsoon region such as low pressure systems, depressions and storms/cyclones etc have been entering on the Asian Australian Monsoon Time Scale as per date and month of each and every year.

If we have been managing the Asian Australian Monsoon Time Scale in this manner continuously, we can see the image and its past's, present's and future movements of the Asian Australian monsoon and study it's originals, climatic changes and futuristic dimensions.

By establishing the Asian Australian Monsoon Time Scales which can help to study the movements of the the Asian Australian monsoon.

#### **Method and Design:**

**Design:** Prepare a Asian Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Asian Australian Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed into a square graphic scale.

This scale should be designed in three ways i.e Basic scale, Filled scale, Analyzed scale;

**Basic Scale:** The first one is preliminary basic scale, it explains the structure of the scale.

**Filled Scale:** This is the second scale that is filled with data and explains how to fill or manage the scale.

**Analyzed Scale:** And the third one is scientifically analyzed the filled scale by data, it explains monsoon patterns weather conditions of the scale.

**Method:** There are two methods in formation and process of the Asian Australian Monsoon Time Scales. The first one is in the single form and next one is designed in four parts.

**Single& Full length Scale:** Prepare the Asian Australian Monsoon Time Scale having 365 horizontal days from April 1<sup>st</sup> to next year March 31<sup>st</sup> (or January 1<sup>st</sup> to December 31<sup>st</sup> or March 21<sup>st</sup> to next year March 20<sup>th</sup> or according to the chronology of Asian Australian's Time and Climate) of 139 year from 1880 to 2027 comprising of a large Time and Climate should be taken and framed in a single and full length type square graphic scale. It can be formed on a paper, board, wall or a Table.

**Parts & paste Scale:** The single and full length square graphic scale is to be long. So that it is divided into four parts easy to carry and keep and suitable for publication. I designed to make it into 4 parts and then pasted it into one scale.

The first part is beginning from 1<sup>st</sup> April to July 12<sup>th</sup>.

The second part is from 13 July to October 23<sup>rd</sup>.

The third part is from 24<sup>th</sup> October to February 3<sup>rd</sup>.

And the fourth part is 4<sup>th</sup> February to March 31<sup>st</sup> ending.

These separate scales can be pasted into one scale as explained below.

Cut along the edges of dates on the right side of the first part and paste it to along the edges of date of 13<sup>th</sup> July on left side of the second part.

Cut along the edges of dates on the right side of the second part and paste it to along the edges of date of 24<sup>th</sup> October on left side of the third part.

Cut along the edges of dates on the right side of the third part and paste it to along the edges of date of 4<sup>th</sup> February on left side of the fourth part .

When paste this manner, we get long full-scape Asian Australian Monsoon Time Scale.

#### **Computer Model:**

Asian Australian Monsoon Time Scales can also be established as a computer model. Besides rather than in manual type scale, If we are able to create a computer model scale which to be the most obvious.

#### **Material and Data:**

Construction of the Asian Australian Monsoon Time Scales requires enormous data of low pressure systems, depressions, tropical storms, sand storms etc that affecting a region and formed over a region should be taken as data to prepare the Asian Australian Monsoon Time Scale. An accurate scale is available if we can collect and analyze the exact climate data.

**Management:** The main weather events if any of Asian Australian monsoon such as monsoon pulses in the form of low pressure systems if any of a monsoon region formed over the Asian Australian monsoon have been entering on the Asian Australian Monsoon Time Scale in stages by 1 for low, 2 for depression, 3 for storm, 4 for severe storm and 5 for severe storm with core of hurricane winds as per date and month of each and every year. If we can managing the scale in this manner continuously, we can study the past, present and future movements of Asian Australian monsoon. I took the numbers to analysis the variations in data. Researchers have to decide what kind of data to take and how to analyze the data.

#### **Researches&results:**

The study should be done in the same way as described in the Indian Monsoon Time Scale and the results should be obtained.

#### **Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

#### **Basics of Malasian Australian Monsoon Time Scale:**

Malaysian-Australian Monsoon, the monsoon system affecting southeast Asia and Australia. It is characterized by winds that blow from the southeast during cooler months of the northwest during the warmer months of the year. Southeast Asia and northern Australia are Combined in one monsoonal system that differs from others because of the peculiar and somewhat symmetrical distribution of land masses on both sides of the equator.

In this respect, the northwest monsoon of Australia is unique. The substantial masses of water

between Asia and Australia have a moderating effect on tropospheric temperatures, weakening the summer monsoon. The many islands (e.g., Philippines and Indonesia) provide an infinite variety of topographic effects. Typhoons that develop within the monsoonal air bring additional complications. It would be possible to exclude North China, Korea, and Japan from the monsoonal domain because their seasonal rhythm follows the normal midlatitude pattern—a predominant outflow of cold continental air in winter and frontal depressions and rain alternating with fine, dry anticyclonic weather in the warm season. On the other hand, the seasonal reversal of wind direction in this area is almost as persistent as that in India. The winter winds of northeastern Asia are much stronger because of the relative proximity of the Siberian anticyclone. The tropical ridge of high pressure is the natural boundary between these non-monsoonal areas and the monsoonal lands farther south.

The northern limit of the typical monsoon may be set at about 25° N latitude. Farther north the summer monsoon is not strong enough to overcome the effect of the traveling anticyclones normally typical of the subtropics. As a result, monsoonal rains occur in June and also in late August and September, separated by a mild anticyclonic drought in July. In South China and the Philippines the trade winds prevail in the October–April (winter) period, strengthened by the regional, often gusty outflow of air from the stationary Siberian anticyclone. Their disappearance and replacement by opposite (southwesterly) winds in the May–September (summer) period is the essence of the monsoon. In any case, these monsoonal streams are quite shallow, about 1,500 metres (4,900 feet) in winter and 2,000 metres (about 6,600 feet) in summer. They bring rain only when subject to considerable cooling, such as anywhere along the steep windward slopes of the Philippines and Taiwan. On the larger islands there are contrasting effects: the slopes facing west receive most of their rainfall from May to October and experience drought from December to April, whereas the slopes facing east receive orographic rains (those produced when moist air is forced to rise by topography) from September to April and mainly convectional rains from May to October.

In Vietnam and Thailand the summer monsoon is more strongly developed because of the wider expanses of overheated land. The southwesterly stream flows from May to October, reaching a thickness of 4 to 5 km (about 2.5 to 3 miles); it brings plentiful but not extraordinary rainfall. The period from November through February is the cool dry season, and the period from March through April is the

hot dry one; in the far south the coolness is but relative. Along the east coast and on the eastward slopes, more rain is brought by the winter monsoon. In the summer, somewhere between Thailand and Cambodia in the interior, there may be a faint line of convergence between the southwesterly Indian-Myanmar monsoon and the southeasterly Malaysian monsoon.

Monsoonal winds are weak over Indonesia because of the expanses of water and the low latitude, but their seasonal reversal is definite. From April to October the Australian southeasterly air flows, whereas north of the Equator the flow becomes a southwesterly. The Malaysian-Australian monsoon generally maintains its dryness over the islands closer to Australia, but farther north it carries increasing amounts of moisture. The northeasterly flow from Asia, which becomes northwesterly south of the Equator, is laden with moisture when it reaches Indonesia, bringing cloudy and rainy weather between November and May. The wettest months are December in most of Sumatra and January elsewhere, but rainfall patterns are highly localized. In Java, for instance, at sea level alone there are two major regions: an “equatorial” west with no dry season and a “monsoonal” east with extreme drought in August and September.

Because of its relatively small size and compact shape, Australia shows relatively simple monsoonal patterns. The north shore is subject to a clear-cut wind reversal between summer (November–April, northwesterly flow) and winter (May–September, southeasterly flow) but with two definite limitations: first, the northwesterly, rain-bearing monsoonal wind is often held offshore and is most likely to override the land to any depth during January and February; second, even in summer there often are prolonged spells of southeasterly trade winds issuing from traveling anticyclones, separating the brief monsoonal incursions. The Australian summer monsoon is thus typical in direction and weather type but quite imperfect in frequency and persistence. Its thickness is usually less than 1,500 metres (4,900 feet) over the sea and 2,000–2,500 metres (6,600–8,200 feet) over the land.

Much less typical are the marginal monsoonal manifestations. On the northwest coast there frequently is a northwesterly airflow in the summer (December–March), as opposed to the winter southeasterlies, but this stream is very shallow and does not bring any rain; that is, its weather is not monsoonal even though its direction is so. On the northeast coast the onshore air is humid and brings rain, but its direction is only partly modified in summer. Most of the summer winds that arrive there occur as a

northeasterly flow, although at other times the flow can be mostly southeasterly.

**Construction:** Keeping in view of study of the aforesaid Keeping in view of study of the aforesaid Malasian Australian Monsoon thoroughly, I have prepared the Malasian Australian Monsoon Time Scale Malasian Australian Monsoon Time Scale is a Chronological sequence of events arranged in between time and weather with the help of a scale for studying the past’s, present and future movements of Malasian Australian monsoon and its relationship with rainfall and other weather problem and natural calamities. Prepare the Malasian Australian Monsoon Time Scale having 365 horizontal days from March 21<sup>st</sup> to next year March 20<sup>th</sup> of a required period comprising of a large time and weather have been taken and framed into a square graphic scale.

**Data:** The monsoon pulses in the form of low pressure systems main weather events if any of the Malasian Australian monsoon region have been taken as the data to prepare this scale.

**Management:** The main weather events if any of the Malasian Australian monsoon have been entering on the Indian Monsoon Time Scale as per date and month of the each and every year. If we have been managing this scale in this manner continuously, we can study the past, present and future movements of Malasian Australian monsoon.

**Researches&results:**

The study should be done in the same way as described in the Indian Monsoon Time Scale and the results should be obtained.

**Study & discussion:**

The obtained results should be studied and analyzed in the same way as described below in the Indian Monsoon Time Scale.

**Evidences that strengthened the Global Monsoon Time Scales:**

**1. Historical evidences that strengthened the Global Monsoon Time Scales:**

Many historical texts in the scriptures such as the Bible and the Quran’s also reinforce the Global Monsoon Time Scales. For example, the text in the Genesis, chapter 41 similar to that on the Global Monsoon Time Scales it was reported that in the past centuries, the monsoons have been going up and down (Rise and Fall )in ordinary English “ there comes seven years of great heavy rains and floods throughout the land of Egypt. And there shall arise after them seven years droughts and famines “. These scriptures reinforce the basic principle of Global Monsoon Time Scales.

## ***2.The IIT'S Study of 100 years of Indian monsoon that strengthened the Global Monsoon Time Scales:***

Deficient rainfall led to the collapse of the Mansabdari system, started by Mughal emperor Akbar, in the late 17<sup>th</sup> century. Similarly, drought interspersed with violent monsoon rains sounded the death knell for the Khmer empire of south-east Asia in the 15<sup>th</sup> century. A recent study by researchers at Indian Institute of Technology, Kharagpur(IIT-KGP) has revealed that abrupt changes in the Indian monsoon strengthened during last 900 years and their linkages to socio-economic conditions in the Indian subcontinent by Nil K. Gupta, Professor at the geology and geophysics, Department of IIT-KGP, highlights that decline of Indian dynasties was linked to weak monsoon and reduced food production.

Rise and fall: Several dynasties, such as the Sena in Bengal, Solanki in Gujarat in the mid-13<sup>th</sup> century and Paramara and Yadav in the early to mid-14<sup>th</sup> century- all of which flourished during the dry phases of Indian summer monsoon suggesting role of the climate in the sociopolitical crisis, the study revealed. The paper published in international journal PALEO 3 highlights three phases in the 900 years stretch-Medieval climate from 950 CE to 1350 CE, Little Ice Age from 1350 CE to 1800 CE and Current Warm Period and phases from 1800 CE till today. The paper highlights strong monsoon during Medieval Climate Anomaly and Current Warm Period and phases of weak. There can be no doubting the profound impact of the abrupt shifts of rainfall on human history-a fact we need to constantly remind ourselves in this day and age of ir retrievable climate change. Abrupt shifts in the ISM precipitation has similarly impacted history in India, Prof.Gupta said.

For the study on long-term spatio temporal variability of the ISM, a group of researchers, which also included experts from Wadia Institute of Himalayan Geology, looked at palaeoclimatic records using oxygen isotope proxy record from speleothems(a structure formed in a cave by deposition of minerals from water) at the Wah Shikar cave Meghalaya.

We took samples from every half millimeter or sometimes even one-third of a mm, and we dated using uranium-thorium time series. Such fine sampling of less time interval means we were covering data at two-three years' interval while most researches collect data 20-30 years' interval. We even captured the drought events of last few centuries, Prof Gupta said. The results showed abrupt shifts in the ISM, he added.

For more recent phases of human history the study suggests that from the beginning of the 19 century, the changes in the ISM became more abrupt with a rise in

atmospheric temperature that coincides with the dawn of the Industrial Revolution.

An increase in the frequency of abrupt shifts in the ISM during the last centuries, coincidental with a rise in atmospheric temperature, suggests occurrence of more climatic surprises in future consequent to future rise in the global temperature and subsequently more precipitation in the form of rain at higher altitudes."the paper said.

Prof.Gupta said that they were doing similar work extending their palaeoclimatic study to 6000 years ago to see the impact of climatic change on Indus Valley civilization and on population migrations.

## ***3.Studies of the Indian Institute of Tropical Meteorology, Pune that strengthened the Global Monsoon Time Scales:***

Studies of long time series of the Index of All India area-weighted mean summer monsoon rainfall anomalies during the period 1871-2017 based on IITM Homogeneous Indian Monthly Rainfall Data Set have revealed the several interesting aspects of the inter-annual and decadal-scale variations in the monsoon that strengthened the Global Monsoon Time Scales.

**FLOOD YEARS:** During the period of 1871-2015, there were 19 major flood years:1874, 1878, 1892, 1893, 1894, 1910, 1916, 1917, 1933, 1942, 1947, 1956, 1959, 1961, 1970, 1975, 1983, 1988, 1994.

**DROUGHT YEARS:** And in the same period of 1871-2015, there were 26 major drought years: 1873, 1877, 1899, 1901, 1904, 1905, 1911, 1918, 1920, 1941, 1951, 1965, 1966, 1968, 1972, 1974, 1979, 1982, 1985, 1986, 1987, 2002, 2004, 2009, 2014, 2015.

Depending on the data mentioned above, it is interesting to note that there have been alternating periods extending to 3-4 decades with less and more frequent weak monsoons over India.

For example, the 44-year period 1921-64 witnessed just three drought years and happened good rainfall in many years. This is the reason that when looking at the Indian Monsoon Time Scale you may note that during 1920-1965's, the passage of the Indian monsoon had been rising over July, August, September in the shape of concave direction and resulting good rainfall in more years..

During the other periods like that of 1965-87 which had as many as 10 drought years out of 23, This is the reason that when looking at the Indian Monsoon Time Scale you may note that during 1965-2004's the path of the Indian monsoon had been falling over the September in the shape of convex direction and causing low rainfall and droughts in many year.

**4. Studies by the Massachusetts Institute of Technology, Cambridge, National Research Foundation, Singapore, Singapore-MIT Alliance for Research and Technology (SMART) that strengthened the Global Monsoon Time Scales:**

A study of the Massachusetts Institute of Technology, Cambridge supported and in part by the National Science Foundation, the National Research Foundation of Singapore, and the Singapore-MIT Alliance for Research and Technology (SMART) founds that the Indian monsoons, which bring rainfall to the country each year between June and September, have strengthened since 2002. Between 1950 and 2002, they found that north central India experienced a decrease in daily rainfall during the monsoon season. To their surprise, they discovered that since 2002, precipitation in the region has revived, increasing daily rainfall. That heightened monsoon activity has reversed a 50-year drying period during which the monsoon season brought relatively little rain to northern and central India. Since 2002, the researchers have found, this drying trend has given way to a much wetter pattern, with stronger monsoons supplying much-needed rain, along with powerful, damaging floods, to the populous north central region of India. A shift in Indian Monsoon Time Scale may explain this increase in monsoon. Consistent with the studies of the above research institutions, this is the reason that when looking at the Indian Monsoon Time Scale you may note that between 1950-2002, the path of the Indian monsoon had been falling over the July and August in the shape of convex direction and decreasing rainfall and since 2002, the Indian monsoon has been rising over July, August, September in the shape of concave direction and precipitation in the region has revived, increasing daily rainfall.

**5. Global Monsoon Time Scales strengthens global researches such as Milankovitch cycles etc that Earth spin on it's axis around the Sun is the root cause of variations in monsoons, seasons and other climate changes:**

Earth has seasons because its axis of rotation is tilted at an angle of 23.5 degrees relative to our orbital plane—the plane of **Earth's orbit around the sun**. The collective effects of changes in the Earth's rotation around its axis and revolution around the Sun such as axial tilt etc may be influenced climatic patterns on the earth. When examining the Global Monsoon Time Scales/ Indian Monsoon Time Scale closely from 1880 to the present, there are many ups and downs in the monsoon cycles. This is the reason for the ups and downs with the monsoons is that the climate changes on the earth forms along the Earth's spin on its axial tilts around the sun. When the Global Monsoon Time Scales/ Indian Monsoon Time Scale is being examined

it is known that there are many unknown mysteries in the Earth's spin on its axial tilts around the Sun. Astrophysicists discover the mysteries of the Earth's spin on its axial tilts around the Sun based on the Global Monsoon Time Scales/ Indian Monsoon Time Scale. Global researches around the world such as **Milankovitch cycles** etc strengthened that the Earth's spin on its axis around the Sun is the root cause of the variations in the monsoons.

**Applications:**

**An overview of current position of monsoons:**

Before explaining the current monsoon and climate conditions, let's take a overview of monsoon pattern since 1880.

Keep track the Indian Monsoon Time Scale carefully. When we look at the Indian Monsoon Time Scale, several paths appears. Two of these are important. These can be called main path of the Indian monsoon(second one-right side) and pre-path of the main passage of the Indian monsoon(first one-left side).

**Pre-path of the Indian monsoon:**

Due to unavailability of data, it is not known how these passages of the Indian monsoon traveled before 1888. But according to the study of records of droughts, famines and floods it is guessed that-

Between 1727-1751 years, it traveled in the shaped of concave direction for about 24 years and caused low rainfall and droughts in many years.

Between 1752-1811 years, it traveled in the shape of convex direction for about 60 years and caused good rainfall and floods in many years.

Between 1812-1835 years, it traveled in the shape of concave direction for about 25 years and caused low rainfall and droughts in many years.

Low pressures, depressions, storms, rainfall, heavy rains, floods and droughts etc. data available since 1880 sufficiently. So since 1880, the path and movements of the monsoons and climate have been scientifically proven and confirmed with certainty as follows.

Between 1836-1895 years, it traveled in the shaped of convex direction for about 60 years and caused good rainfall and floods in many years.

Between 1896-1919 years, it traveled in the shape of concave direction for about 24 years and caused low rainfall and droughts in many years.

Between 1920-1981 years, it traveled in the shape of convex direction for about 62 years and caused good rainfall and floods in many years.

Between 1982-2009 years, it traveled in the shape of concave direction for about 27 years and caused low rainfall and droughts in many years.

From 2010, it is going to travel upwards in the shape of convex direction for 56 years that's until

2056 and will be resulting good rainfall and floods in the coming years.

#### **Main-path of Indian monsoon:**

Due to unavailability of data, it is not known how these passages of the Indian monsoon traveled before 1888. But according to the study of records of droughts, famines and floods it is guessed that-

Between 1797-1836 years, it traveled in the shaped of concave direction and caused low rainfall and droughts in many years.

Between 1837-1860 years, it traveled in the shape of convex direction and caused good rainfall and floods in many years.

Between 1861-1882 years, it traveled in the shape of concave direction and caused low rainfall and droughts in many years.

Low pressures, depressions, storms, rainfall, heavy rains, floods and droughts etc. data available since 1880 sufficiently. So since 1880, the path and movements of the monsoons and climate have been scientifically proven and confirmed with certainty as follows.

Between 1883-1901 years, it traveled in the shape of convex direction and caused good rainfall and floods in many years.

Between 1902-1928 years, it traveled in the shape of concave direction and caused low rainfall and droughts in many years.

Between 1929-1950 years, it traveled in the shape of convex direction and caused good rainfall and floods in many years.

Between 1950-1965 years, it traveled in the shape of concave direction and caused low rainfall and droughts in many years.

Between 1965-1981 years, it traveled in the shape of convex direction and caused good rainfall and floods in many years.

Between 1982-2020 years, it traveled in the shape of concave direction and caused low rainfall and droughts in many years.

From 2020, it is going to travel upwards in the shape of convex direction for 56 years that's until 2056 to 2075 and will be resulting good rainfall and floods in the coming years.

#### **Current weather condition:**

While examining the Indian Monsoon Time Scale, it appears that the summer Monsoon is traveling in the upper direction.

For example, the pre-path of monsoon was at its lowest point on July 25th, 2000 slowly moved up and reached July 11th, 2010 after 10 years. And the main-path of the monsoon was at its lowest point on August 17th, 2000 slowly moved parallel to the pre-path with a difference of about 30 days and reached August 12, 2010 after 10 years.

When the same monsoon is seen after 10 years, the pre-path of monsoon was at July 11th, 2010 slowly moved further up and reached July 4th, 2020 after 10 years. And the main-path of the monsoon was at on August 12th, 2010 slowly moved parallel to the pre-path with a difference of about 30 days and reached August 02, 2020 after 10 years.

In the current year 2022, the pre-path of Indian summer monsoon was traveling upwards and reached to the 29th June. Beside this, the main-path of Indian summer monsoon also traveled upwards parallel to the pre-path of Indian summer monsoon with a difference of about 30 days and reached to the 29th July, As it moves further up, changes in the climate are likely increasing and there are more chances of heavy rains and floods in the coming years

Although these reports were revealed by the Indian Monsoon Time Scale, they reflect the upcoming global climate changes. However, if we set up separate Monsoon Time Scales for the respective monsoon systems & countries and analyze the data of their monsoon systems and countries, accurate results will be obtained for the respective country and monsoon.

#### **Future:**

As discussed above, the convex period of pre-path which traveled between 1918-1981 will be traveled between 2010-2060 and the convex period of the main-path which traveled between 1926-1981 will be traveled between 2020-2075.

As result, heavy rains and floods are going to occur all over the world countries including above country in the coming seasons. Rain is a major component of the water cycle and is responsible for depositing most of the fresh water. It provides water for hydroelectric power plants, crop irrigation, drinking water and suitable conditions for many type of ecosystems.

Widespread heavy rainfall from a active monsoon or cyclone has several benefits as it is usually spread over a number of days. Increased rainfall helps the ground to hold more moisture, which in turn means that future crops have major benefit with more moisture being made available for a longer time. Heavy rains can cause pooling, overflowing rivers and runoffs, and flooding. These events may result in evequations, power outages, supply shortages, traffic obstructions and road closures, infrastructure damage and debris.

And also future climate changes are expected to include a warmer atmosphere, a warmer and more acidic ocean, higher sea levels, flooding, storms and more large change in precipitation patterns.

Therefore, precipitation including heavy rains, snow, floods will occur. People who live in the water catchment areas may be trapped in floods as the water flow into the towns and villages in their former way.

As a result massive loss of life and property is going on. So the scientists establish the Monsoon Time Scale. Many cities, Islands and villages situated on the shore of rivers and seas will get absorbed in the water. Heavy rains, floods, cyclones can lead to disease spread and damage to ecosystems and infrastructures. Human health issues can increase mortality etc.

According to an estimate, rivers, lakes, reservoirs, barrages and dams etc. may full with waters in the coming years. Through this research proposal, we can know the future consequences of rivers, lakes, reservoirs, barrages and dams etc. Plans can be made accordingly. So, scientists can establish the Indian Monsoon Time Scale for rivers, lakes, reservoirs, barrages and dams etc. and predict what is going to happen in the rivers, lakes, reservoirs, barrages and dams etc. basin catchment areas in the coming years roughly.

Water generally collects in a rivers, lakes, reservoirs, barrages and dams etc. from precipitation and other sources such as groundwater recharges, springs, natural ice snow packs. In the recent decades, monsoon or climate is weakening and rains are shrinking. Rivers, reservoirs, barrages, ponds are falling and drying. Some rivers, lakes, reservoirs, barrages and dams etc. are extinct. Some rivers, lakes, reservoirs, barrages and dams etc. may have dried up or water flowing in the river may have reduced. Climate changes, heavy rains, droughts etc. affect the rivers. Due to these climate changes, monsoon failures and drought conditions, water catchment areas are becoming villages and towns as people made houses with a feeling that the rains do not come and the rivers, lakes, reservoirs, barrages and dams etc. are not inundated with waters. However, governments should consider one important thing. Perhaps sometime in the coming years and decades, the monsoon repeats as early as previous years and decades, there heavy rains and floods are going to happen in the coming years. The rivers, lakes, barrages, reservoirs and ponds will be filled with waters. People who live in those water catchment areas are trapped in the heavy rains and floods as the rivers, lakes, reservoirs, barrages and dams etc. flow into the towns and villages in their former way. Or the rivers, lakes, reservoirs, barrages and dams etc. that are still flowing in abundance will cause even more abundant floods in the future. Due to all of these, some advantages and disadvantages are going to happen in future. As a result massive loss of life and property is going on. It is known that during the next 50 years there will be changes in the monsoon climate and heavy rains will flood the rivers, lakes, reservoirs, dams in the coming years. It is possible to predict what climate conditions will be like in rivers, lakes, reservoirs, barrages and dams etc. basin areas in

the next 50 years roughly by Indian Monsoon Time Scale. Indian Monsoon Time Scale will be used to study the past, present and future movements of climate and monsoon and its rainfall conditions and assess & evaluate the upcoming conditions of rivers, lakes, reservoirs, barrages and dams etc. and taking necessary precautions on the basis of those parameters. So, scientists need to develop Indian Monsoon Time Scales to analyze the climate changes affecting the rivers, lakes, reservoirs, barrages and dams etc. Through them, the climate changes and flow of the rivers, lakes, reservoirs, barrages and dams etc. can be predicted about 50 years in advance and measures can be taken accordingly.

Here is an important point to be grasped that the Indian Monsoon Time Scale's analysis is concerned with the Indian monsoon region but it reflects and informs the climate changes of all the countries of the world. In that case the aforesaid Monsoon Time Scale must reflect the climate changes of the country which is close to the aforesaid monsoon. Monsoon Time Scale gives accurate results if it is related to the climate of the country.

#### **Scientific theorem:**

This is a phenomenon of Earth and space sciences and effect of astronomical bodies and forces on the earth's geophysical atmosphere. The cause is unknown however the year to year change of movement of axis of the earth inclined at  $23\frac{1}{2}$  degrees from vertical to its path around the sun does play a significant role in formation of the monsoon.

Everything in the universe just like oceans, solid earth, biological, atmosphere, geomagnetism, global and regional geophysical systems and sun, moon, planetary, solar-terrestrial astrophysical systems have many different types of interactions with each other. Many combinations of these simple interactions can lead to surprising emergent phenomena and play a key role in creation of monsoons and other weather changes and natural calamities on the earth.

Monsoon is traditionally defined as a seasonal reversing winds. The primary cause of monsoons is the difference between annual temperature trends over land and sea. In winter the land is colder than the sea. Most of the time during the summer the land is warmer than the ocean. This causes air to rise over the land and air to blow in from the ocean to fill the void left by the air that rose. However, the physical factors of these monsoon are mainly influenced by the rotations and revolutions of the earth around the sun.

Earth rotates or spins on its axis and it also orbits or revolves west to eastward around the sun. Rotation and revolution are two motions of the Earth. Rotation of the Earth is its turning on its axis. Revolution of

Earth is the movement of the Earth around the sun. The Earth rotates about an imaginary line that passes through the North and South poles of the planet. This line is called axis of rotation. Earth rotates about this axis once each day approximately 24 hours. The earth's axis of rotation is tilted by 23.5 degrees from the plane of it's orbit around the sun. The cause is unknown but the year to year change of movement of axis of the earth inclined at 23½ degrees from vertical to its path around the sun does play a significant role in formation of clusters, bands & paths of the Indian Monsoon and stimulates the Indian weather. The inter-tropical convergence zone at the equator follows the movement of the sun and shifts north of the equator merges with the heat low pressure zone created by the rising heat of the sub-continent due to direct and converging rays of the summer sun on the India Sub-Continent and develops into the monsoon trough and maintain monsoon circulation.

#### **Conclusion:**

We can make many more modifications, thus bringing many more developments in the Monsoon Time Scales, Geoscope projects and can examine the possibilities of using them according to the climate conditions and natural calamities of the country.

**Appeal:** However, much efforts and sacrifice did tho, I could not get government recognition and social support. My researches were ignored and darkened. I am a victim of racism and discrimination, negligence and jealousy. Throughout my life, I have experienced hardships all my life. I was abused, humiliated and beaten when I asked to provide research opportunities. I was pushed out of the gate, when I asked to provide research opportunities. I was insulted by my race. I was tied to a pole and beaten. My thoughts and researches were subjected to the wrath of racists, casteists and fanatics as well as fellow scientists and resulted into oppression on me. My lab was invaded. Illegal cases were framed and foisted against me. I faced trials, handcuffed and led through streets police enquiries and court trials/hearings, and imprisoned. Political recommendations and officials support, cash and caste, region and religion may play a key role in giving support and opportunities, awards and rewards, respect and recognition to depressed communities. But I have no of them. I am now making my life's last journey due to disregard, despair and serious illness, severe poverty.

Kindly find out my researches in all social networking websites or can obtain by sending your email to me. These findings are very helpful for research institutions, universities researches. And also these findings can be very helpful for Ph.D students, Postdocs, professors, seniors, scientists and science

enthusiasts who want to innovate. I will send them the valuable information I have.

For example, those who want to design Monsoon Time Scales for their regional or country' Monsoons and conduct weather predictions have trouble in making the Monsoon Time Scales, kindly contact me at my email id [gangadhar19582058@gmail.com](mailto:gangadhar19582058@gmail.com) and take my suggestions and assistance. I will send you complete details of the Monsoon time scales. Further if you want, I will create a manual Monsoon Time Scale and send the same to you for study and research. However for this, data of list of monsoon pulses in the form of monsoonal low pressure systems, depressions and storms formed over their monsoon region or country last 100 and above years since 1880 as cited in the Reference-1 (i.e Mooley DA, Shukla J(1987); Characteristics of the west ward-moving summer monsoon low pressure systems over the Indian region and their relationship with the monsoon rainfall. centre for ocean-land atmospheric interactions, university of Maryland, college park, MD.). I will make and send it to you. So, researchers send Monsoon data of their region or country, I will make and send Monsoon Time Scales for their region or country. These monsoon time scales are very helpful for research institutions, universities researches and also these can be very helpful for Ph.D students, Postdocs, professors, seniors, scientists and science enthusiasts who want to conducting researches and studies on climate changes there. Because, through these Monsoon Time Scales it is known in advance that what kind of climate changes have occurred in your country in the past 100 years and what kind of climate changes are going to happen in the coming 100 years.

**I am now making my life's last journey in serious illness and poverty. Illness weakening the health and mind slows down and forgetfulness is coming. It is not known how long I will live and when I will die, but I know my time is near. Hence, I humbly request that if world scientists have invented any technology in future that re-create humans of past, kindly remember and re-create me to complete my uncompleted researches as an attendant in your research laboratory.**

**GANGADHARA RAO IRLAPATI**

**Corresponding Author:**

**Gangadhara Rao Irlapati**

**H.No.5-30-4/1,**

**Saibabanagar, Jeedimetla(IDA)**

**Hyderabad-500055,**

**Telangana State, INDIA**


**Email: [gangadhar19582058@gmail.com](mailto:gangadhar19582058@gmail.com)**

**Googlepay/PhonepeA/cNo.+91 6305571833**

**Kotak Bank A/C No. 8447 502 446**



Simple Geoscope Model:




Home-Made Geoscope Model:



Seismic luminescence study:





Electrogeogram Test:




GEOSCOPE




















THE ITCZ MOVES NORTH OVER THE INDIAN REGION. THE ITCZ PASSING OVER THE ANDAMAN PROVINCE.






















మహారాజశ్రీ రెవెన్యూ డివిజనల్ ఆఫీసరు  
వారి దీక్షాసముఖమునకు,  
అమలాపురం.

తూర్పుగోదావరి జిల్లా, కొత్తవేలి లాలాకా మెర్సెసాలెం గ్రామకాపురముడు ఇర్లపాటి  
పుల్లయ్య కుమారుడు ఇర్లపాటి గంగాధరరావు అను నేను మిక్కిలి విదేయతో నమస్కరించి  
దాఖలు చేసుకొను విన్నపములు.

అయ్యో,

నేను శాస్త్ర పరిశోధనలు చేసి దేశానికి నేవలు చేయాలనే ఆశయమును కలిగిన  
శాస్త్రపరిశోధకును. ఇంటి వద్దనే చిన్న పరిశోధనాలయమును వెట్టుకొని ప్రయోగాలు చేసు  
కొంటున్నాను. సూక్ష్మ అపిర్మావము, నిర్మాణము, ధర్మాలు, పరిణామము మానవసూక్ష్మ మతము  
దైవము మొదలగు విషయాలను విశదీకరిస్తూ, వాదాలను ప్రతిపాదించాను. ఇట్టికాకుండా  
ప్రజలను తుఘ్నాలు, కర్మకాటకాలు, నరదలవంచి ప్రకృతివైపరీత్యాలనుండి కాపాడబానిసగాను  
కొన్ని సేలలును వడ్డతులను దీర్చునోపు వంటి తరీకరాలను రూపొందిస్తున్నాను. ఇంకా  
అనేక శాస్త్రీయ ప్రచురణలు ప్రచారము ద్వారా నేవచేస్తున్నాను. అయితే మాగ్రామ కరణంగారు,  
మునసబుగారు, ఆత్రేయపురం రెవెన్యూ ఇన్స్పెక్టరుగారు, కొత్తవేలి తహసిల్దారు గారు ఇతరులు  
మాధనముకాల్తో నా సిద్ధాంతాలను విమర్శిస్తూ, వాగాభవము చేస్తున్నారు. నా పరిశోధనలకు  
అడ్డంకులు కలిగిస్తున్నారు. నాకు కులధర్మపత్రమున్న సంతకము వెట్టుకుండా బాదిస్తున్నారు.  
దయతో ఈ విషయమే విచారించి నాకు రక్షణ కల్పించమని న్యాయము చేయమని వేడుకొనుచున్నాను.

ఇటు, తమ విశ్వాసనీయుడు,  
9. Gangadhara Reddy  
6-7-77  
:ఇర్లపాటి గంగాధరరావు:

మెర్సెసాలెం,  
తే. 6-7-1977

11-27-

Received a tipped report Taluk Magistrate Kotta Peta with the following:-  
 Ref: A.S. 5973/77 dt 21.7.77 Taluk office Kotta Peta

From: Sri P. Subbarao, Esq. Taluk Magistrate Kotta Peta To: The Station House Officer Ravalapalem.

Sir: Subj: Forgery Signature - Srig. Jalapati gangadhara Rao of Ravalapalem V. Report of the Revenue Inspector, Amalapuram.

Ref: Report of the Revenue Inspector, Amalapuram dt 21.7.77.

The Rev. Inspector Amalapuram, enquired and reported that Smt. Relangi Rathamma wife of Musalaiah of Nerlapalem Village applied for grant of a tree (Jewinid) situated on the north-west portion of her house for which house site Patta was granted. On the above petition the signatures of Village Headman, Nerlapalem and the Rev. Inspector, Amalapuram were forged.

The Rev. Inspector, Amalapuram further reported that Smt. Relangi Rathamma in her statement deposed that the son of Sri Jalapati Pullaiah forged the signatures. As such the Rev. Inspector, Amalapuram has called for the individual and owned in to the matter and reported that he failed Intermediate and left hand writer. He accepted that he forged signatures and the signatures of the Village Headman, Nerlapalem and the Rev. Inspector, Amalapuram. He is a very dangerous boy and is up to any thing.

In the above case, Sri Jalapati gangadhara Rao of Pullaiah of Nerlapalem Village, the offender in the instant case may be dealt with according to law. Please intimate the action taken in the matter.

1. The following records are enclosed here with duty officer's file and enclosed.

2. Slip containing forged Signature.

3. Statement recorded from Sri Jalapati gangadhara Rao of Pullaiah of Nerlapalem Village.

4. Statement of Smt. Relangi Rathamma wife of Musalaiah of Nerlapalem Village.

5. Report of the Rev. Inspector, Amalapuram dated 21.7.77.

6. The offender is produced before you through the Rev. Inspector Amalapuram for taking in to custody.

Enclos: - As stated above.  
 (sd) P. Ramasubbingi Head clerk.

yours faithfully,  
 sd, P. Subbarao  
 Taluk - Magistrate  
 Kotta Peta.

Copy Submitted to the collector, Kakinaada.  
 Copy Submitted Superior Sub-division of Police, Kakinaada,  
 Copy to the Rev. Divl. Officer - Amalapuram.  
 Copy to the Circle Inspector of Police - Amalapuram.

To the }  
Jahesildan }  
Kotte Peta }

-26-

Sir I registered the above as C. no 53/47 U/S 420,  
467, and 471 J.P.C and copies of F.I.R. submitted to all  
concerned officers and original F.I.R were sent to J.P.C Magistrate  
Kotte Peta.

Sd/- K.N. Hemanth He. 1635-  
S/O 21. 7. 77  
Kavalapalau.

" True copy "

Sd/-  
He. 247  
S/O Kavalapalau

IN THE COURT OF THE JUDICIAL MAGISTRATE OF THE I CLASS KOTHAPETA.  
 PRESENT: SRI D. VENKATAMARAYANA, B.COM., LL.B., Judicial Magistrate  
 of the I Class.

TUESDAY, the 27th day of November, 1979.

C.C.No. 13/79.

Between:

The State of Andhra Pradesh, through

The State Inspector of Police, Razole  
 Cr.No.53/79 of Ravulapalem P.S.

.. Complainant.

and

Irlapati Gangadhara Rao,  
 s/o Fullayya, Aged 19 yrs.  
 Merlapalem.

.. Accused.

This case coming on 20.11.79 for hearing before me in the presence of the State Complainant and the accused appearing in person and having stood over for consideration till this day, the court delivered the following:-

JUDGMENT

The Inspector of Police, Razole has laid the charge sheet in Cr.No.53/79 of Ravulapalem Police Station Under Sections 420, and 471 IPC against the accused herein.

2. The case of the prosecution is that P.W.1 is resident of Merlapalem village and she is living in a house constructed in R.S.No.129 in Merlapalem village which was given to her by the Revenue Department. There is a tamarind tree in the said house site near her house. The branches of the said tree were over-hanging on her house endangering safety to her house. She was advised to apply for patta of the said tamarind tree. The accused who had come to know about it approached P.W.1 two weeks prior to 21.7.77 and offered his services to get the said tree or patta for her and he induces her to affix her thumb impression on the application written by him and wanted her to get the recommendations of the Village Munsif and Revenue Inspector, Atreyapuram. When she expressed her inability to secure their signatures he resorted to forging of the signatures of Village Munsif, Merlapalem and Revenue Inspector (P.W.4). Completing the application and the recommendations he presented the application in the Taluk Office,

... for verification and enquiry on 21.7.77, contacted P.W. 1 to ... also questioned the accused at the village chavidi of Ryali before whom the accused admitted the offence and P.W.4 recorded the statements of P.W.1 and the accused. The accused was produced before the Tahsildar, Kothapeta who forwarded the accused to the Police Station, Ravulapalem along with Exs.p1 to p4. The police, Ravulapalem registered Cr.No.63/77 u/s. 420,467 and 471 IPC. Therefore, the accused is liable for punishment under sec. 420,467 and 471 IPC.

3. The case was taken on file against the accused under sec. 420, 467 and 471 IPC. When the accused appeared before this court, copies of documents contemplated under sec. 207 Cr.P.C. were furnished to him and he was examined on the contents of the documents. He denied the offence. On consideration of the documents, a charge under sec. 420, 467 and 471 IPC were framed, read over, interpreted and explained to the accused in Telugu to which he pleaded not guilty and claimed to be tried.

4. The prosecution, in support of its case, examined P.W.1, who wanted to apply patta of the tamarind tree, P.W.2 the village Munsif, Ryali, P.W.3, village Kanna of Ryali, P.W.4 the Revenue Inspector in whose presence the accused is alleged to have confessed the offence, P.W.5 the Head Constable who registered the crime. P.W.6 the Investigating Officer, P.W.7, the Tahsildar who forwarded the accused and report of P.W.4 to Ravulapalem P.S. and got marked Ex.p1 to p6. The accused did not adduce any oral or documentary evidence.

5. After closure of the prosecution evidence, the accused was examined u/s. 213 Cr.P.C. regarding the incriminating circumstances appearing in the evidence of the prosecution against the accused. The plea of the accused is total denial of the offence. He stated that P.W.4 is superstitious and fanatic and that when P.W.4 was talking about God once he told him that human being was ... therefore, P.W.4 gave wild in that connection

is that he was beaten by P.W.4 and others and he was forced to put his signature on Ex.P3 and also Ex.P2. Further, the plea of the accused is that there was altercation between him and P.W.4 with regard to the existence of God and also with regard to obtaining of signature of P.W.4 on the caste certificate. Except, the confession statement of the accused Ex.P3 before P.Ws. 2 to 4, there is no direct evidence to connect the accused with the offences charged against him. P.W.4 is an illiterate. She does not know on which paper the accused obtained her thumb impression. Even for a moment sake, it is presumed that it is the accused who obtained the signature of P.W.1, on Ex.P1, Ex.P1 itself is completely in torn condition and the Tahsildar, Kothapeta who is competent authority to grant patta of the tamarind tree, would not have acted upon the petition Ex.P1. Moreover, the prosecution failed to explain the reason why the accused forged the signature of P.W.4 and the Village Munsif, Merlapalem on Ex.P1 and by forging the signature what is the wrongful gain the accused wanted to obtain. There is no evidence to show that it is the accused who filed Ex.P1 petition and other enclosures in the Tehsil Office, Kothapeta. Further, there is a typed petition filed in this case which contains the recommendation of the Village Munsif and the recommendation of Revenue Inspector-P.W.4. It is not marked by prosecution. To support a conviction U/s. 467 IPC, there must be evidence that the document is a false document, within the meaning of section 464 IPC and that it was forged by the accused with some intent mentioned in sec. 463 IPC. It is not sufficient that some possible intent may be inferred from the facts, it is necessary such intent should be established by evidence, which is lacking in this case. Under Sec. 420 IPC, there must be evidence that the person deceived delivered to someone, or consented that some person shall retain certain property, that the person deceived was induced by the accused to do as above, that such person acted upon such inducement in consequence of his having been deceived by the accused, that the accused acted fraudulently.

and that subsequently when he approached P.W.4 to sign on the caste certificate, he demanded Rs. 10/- from him and that subsequently he reported the matter to the Revenue Divisional Officer, Amalapuram about the demanding of illegal gratification of P.W.4. The R.D.O. Amalapuram has promised to enquire into the matter. Therefore, this case is falsely foisted against him. When he was coming from Ravulapalem the Village servant took him before P.W.4. Thereafter he was ~~kept~~ taken to village chavidi where P.Ws. 1 to 4 were present and they beat him and obtained his signature on Ex.P3 and subsequently he was taken to the Pansilgar, Kothapeta from there he was sent to Police Station, Ravulapalem and that he is innocent and he did not commit any offence.

6. The point for consideration is whether the prosecution has been able to establish its case against the accused, beyond all reasonable doubt?

7. The case of the prosecution is that the accused forged the signature of P.S.4 the Revenue Inspector and Village Munsif, Merlapalem (who is no more alive). Ex.P1 is the petition which contains the alleged forged signatures of Village Munsif, Merlapalem and Revenue Inspector (P.W.4). Ex.P1 is in torn condition. The alleged signature of Village Munsif, Merlapalem is completely torn and the signature of P.W.4 is also torn completely except some portion. It also contains the thumb impression alleged to have been affixed by P.W.1. The prosecution to establish that it is the accused who is responsible for the alleged forgery of signatures of P.W.4 and Village Munsif, Merlapalem relies on Ex.P1 petition and Ex.P2 the slip which is also alleged to have been signed by the accused in the presence of P.Ws. 2 to 4. There is no direct evidence available, in this case, who witnessed the forging of the signatures of P.W.4 and Village Munsif, Merlapalem. Even the alleged signatures are in torn condition. Regarding the statement of the accused recorded by P.W.4 in the presence

dishonestly when so inducing that person, that the accused so induced that person intentionally, that such act of the accused was likely to cause damage or harm to that person in property. There must also evidence of fraudulent or dishonest intention at the time of the omission of the act in respect of which the cheating is alleged. Since the main part of the alleged signatures of P.W.4 and Village Munsif, Merlapalem (who is no more) are completely torn and Ex.P1 is in such a condition that the Tahsildar, Kothapeta would not have been acted upon it in granting patta of the tamarind tree to the petitioner i.e., P.W.1. Therefore the question of commission of offences of cheating and thereby dishonestly inducing delivery of property, forgery of a valuable security or authority to make transfer any valuable security and using a genuine a forged document which is known to be forged are not proved against the accused, beyond all reasonable doubt.

In the result, the accused is given the benefit of doubt. The accused is found not guilty of the offences punishable Under sections 420, 467 and 471 IPC. and he is acquitted Under sec. 248(1) Cr.P.C.

Dictated to the Shorthand-writer, transcribed by him, Corrected by me and pronounced in Open Court on this the 27th day of November, 1979 in the presence of the accused.

Sd.D.Venkata Narayana, 27.11.79  
Judicial Magistrate of the  
1st Class, Kothapeta.

Appendix of evidence.  
Witnesses examined for.

Prosecution:

P.W.1: Relangi Rattamma  
P.W.2: Pericherla Satyanarayanaraju.  
P.W.3: T.V.Sriramachandra Murty.  
P.W.4: Malladi Panduranga Vithal,  
RI, Atreyapuram.  
P.W.5: K.M.Meera Sahe,  
HC 1625, Ravulapalem P.S.  
P.W.6: T.B.Pundarikakshudu,  
Inspector of Police,  
Ravulapalem.  
P.W.7: P.Subba Rao,  
Tahsildar, Kothapeta.

Defence:

None.

Documents marked:

- Ex.P1: Forged petition, dt. 10.7.77 of P.W.1
- Ex.P2: Slip
- Ex.P3: Statement of accused. Nil.
- Ex.P4: Statement of P.W.1
- Ex.P5: F.I.R. in Cr.No. 53/77.
- Ex.P6: Petition forwarded by the Tahsildar, Kothapeta to the S.H.O. Ravulapalem.

M.Os marked:

Nil.

Sd. D. Venkatanarayana  
27.11.79  
Judicial Magistrate of I Class  
Kothapeta.

-/true copy/-

*J. Venkatesh*  
J. P.C. MAGISTRATE  
KOTHAPETA.

*63*  
*25/11/79*

CALENDAR AND JUDGMENT  
IN THE COURT OF THE JUDICIAL MAGISTRATE OF THE I CLASS  
KOTHAPETA.  
C.C.No. 13/79.

Date of:  
Offence: 2 weeks prior to 21.7.77  
Complaint: 1.2.79  
Apprn. of accused: 13.2.79.  
Release on bail: 13.2.79.

Commencement of trial: 2.4.79  
Close of trial: 20.11.79.  
Sentence/Order: 27.11.79  
The presiding officer is on CL from 22.11.79 to 24.11.79 and is on permission on 25.11.79).

Explanation for the delay and remarks: The delay is due to non-production of witnesses by the complainant.  
Complainant: The S.H.O. Ravulapalem Cr.No.53/79.

-----  
Name of accused. Father's name. Age. Religion. Calling Village Taluk  
-----

Irlapati Gangadha-  
ra Rao. Pullayya 19 Hindu Mazdoor Merla- Kotha-  
palem. peta

-----

Offence: Under Sec. 420, 467 and 471 IPC.  
Finding: Not guilty.  
Sentence/Order: The accused is acquitted U/s 248(1) Cr.P.C. of the offence Under Sec. 420, 467 and 471 IPC.


Sd.D.Venkata Narayana  
27.11.1979  
Judl.Magistrate of the 1st class  
Kothapeta.

-/true copy/-

J. F. C. MAGISTRATE -  
KOTHAPETA.

28/11/79.







401  
राज्य मन्त्री  
विज्ञान और प्रौद्योगिकी, परमाणु ऊर्जा,  
अन्तरिक्ष, इलेक्ट्रॉनिक्स एवं महासागर विकास  
भारत सरकार, नई दिल्ली

MINISTER OF STATE  
SCIENCE & TECHNOLOGY, ATOMIC ENERGY,  
SPACE, ELECTRONICS & OCEAN DEVELOPMENT  
GOVERNMENT OF INDIA

9th December, 1988.

Dear Shri Rao,

I have your letter dated 15th November, 1988,  
enclosing a petition from Shri Gangadhara Rao  
Irlapati.

2. I will try to help.

Yours sincerely,

( K.R. NARAYANAN )

Shri A.J.V.B. Maheswara Rao,  
Member of Parliament (LS),  
43, North Avenue,  
New Delhi.

Hyderabad,  
Date:03-06-1989

To

The Director General,  
Council of Scientific and Industrial Research,  
Rafi Marg, New Delhi-I.

Sir,

Sub: Invention of Geoscope - Requested for further  
research and development at the National Geophysical  
Research Instituted - Reg.

- Ref: 1) Letter dated: 03-12-1987 of A.J.V.B.M. Rao,  
Member of Parliament (LS), Amalapuram.  
2) Letter No.401/VIP/MOS/88 Dated:8th December,1988  
of Sri K.R.Narayanan, Minister of State Science  
& Technology, New Delhi.

I am a poor scientist with an ideal to serve the Country  
through Scientific research. I have invented and built a  
small Geoscope at my house which can help to study the  
underground.

Geoscope is a simple and wonderful invention. A borehole  
having suitable width and depth has to be dug. An  
Observatory having research and analysis facilities has to be  
constructed on the borehole various ~~xxxxxxxxxxxxxxxx~~  
sensing apparatus to recognize the geophysical and geochemical  
changes generated in the underground should be inserted into  
the underground through the borehole and linked with the  
concerned analysis departments of the observatory that is  
above the ground to study the changes taking place in the  
underground.

Kindly provide research facilities to carryout further  
researches on the Geoscope project at N.G.R.I. Hyderabad.

Gangadhara Rao Irlapati  
C/o. R. Mohana Rao,  
Saibaba Nagar,  
Jeedimetla,  
Hyderabad, AP.

Yours faithfully,

*G. Gangadhara Rao*

In the High Court of Andhra Pradesh at Hyderabad.  
Special Original Jurisdiction

Wednesday the Sixth day of September  
One thousand nine hundred and eighty nine

Present

The Hon'ble Mr. Justice Lakshmana Rao

Writ Petition No. 12355 of 1989

Between:

Irlapati Gangadhara Rao.

..

Petitioner

And

1. Union of India, rep. by its Secretary,  
Ministry of Science & Technology, Anusardhana  
Bhavan, Rafi Marg, New Delhi-1.
2. Council of Scientific & Industrial Research,  
rep. by its Director General, Rafi Marg, New Delhi-1.
3. National Geophysical Research Institutes rep.  
by its Director, Taranaka, Hyderabad. .. Respondents.

Petition under Art. 226 of the Constitution of India praying that in the circumstances stated in the affidavit filed herein the High Court will be pleased to issue an appropriate writ or order or direction declaring

- i) that the inaction of the respondent authorities in not considering petitioner's representations for carrying out research and scientific investigations as arbitrary, unreasonable and illegal;
- ii) a direction may be issued to the respondents 2 & 3 to consider the petitioner's representations so as to enable him to carry out scientific investigations in respondent 3 institution, or any such other appropriate direction may be passed;
- iii) Costs be awarded to the petitioner;

For the Petitioner : Mr. K. Ramakrishna Reddi, Advocate

For the Respondents : Mr. S. Venkateswara Rao, S.C. for Central Govt.

The Court made the following: ORDER

Heard the learned counsel for the petitioner as well as the learned Standing counsel for the Central Govt. appearing on behalf of the respondents.

The relief sought for in this writ petition is a direction to the respondents to consider the respondent representations submitted by the petitioner to ~~xxxx~~ provide facilities to enable him to carry out scientific investigations in National Geophysical Research Institute, Hyderabad and pass appropriate orders thereon.

Having regard to the facts and circumstances of the case, it is directed that the respondents shall consider the representation dated 3-6-89 submitted by the petitioner and pass appropriate orders thereon as early as possible preferably within three months from the date of receipt of a copy of this order.

The writ petition is accordingly disposed of. No costs.

Sd/- S. R. Choudary  
Asst. Registrar

//true copy//

Asst. Registrar

To  
1. The Secretary, Union of India, Ministry of Science & Technology,  
Anusardhana Bhavan, Rafi Marg, NEW DELHI-1.  
2. The Director General, Council of Scientific & Industrial Research,  
Rafi Marg, NEW DELHI-1.

From:  
Gangadhar Rao Iratapati,  
Merlapalem Village  
Vubalanka Post - 532232,  
Atrypuram, V.G. District,  
Andhra Pradesh.

To - - -  
The Director of General of  
Meteorology,  
India Meteorological Department  
New Delhi.

Through : Shri G.M.C. Balayogi  
Member of Parliament (LS)  
Amalapuram.

Sir,

Sub: Global Monsoon Time-Scales - Indian Monsoon Time Scale -  
Requested for further research & development - Reg.,

I am a poor scientist with an ideal to serve the country  
research. I have built a small Lab at my house and conducting  
research on the Global Monsoon systems. As a part of this, I have  
invented the Indian Monsoon Time Scale which can help to study  
the past, present and future movements of the Indian Monsoon.

I am request you that kindly accept my Indian Monsoon  
Time Scale and Develop in the services of the country.

Merlapalem

15-08-1996.

Yours faithfully,

S. Gangadhar Rao  
15/8/96.



सं०  
 भारत सरकार  
 भारत मौसम विज्ञान विभाग  
 मौसम विज्ञान के महानिदेशक का कार्यालय  
 मौसम भवन, लोदी रोड  
 नई दिल्ली-११०००३  
 तार का पता :  
 महामौसम, नई दिल्ली

NO. NA-153  
 GOVERNMENT OF INDIA  
 INDIA METEOROLOGICAL DEPARTMENT  
 OFFICE OF THE  
 DIRECTOR GENERAL OF METEOROLOGY  
 MAUSAM BHAVAN, LODI ROAD,  
 NEW DELHI-110003  
 Telegraphic Address  
 DIRGENMET, NEW DELHI  
 दिनांक/Date. Oct. 2/ 1991.

To

Shri Gangadhara Rao Irlapati,  
 Merlapalem Village,  
 Vubalanka Post 533237  
 Atryapuram, E.C. Distt.,  
ANDHRA PRADESH

Sir,

Kindly refer to your letter dated 15.8.91 received through Shri G.M.C. Balayogi, M.P. regarding the invention of an instrument by you which can help to forecast cyclones, rains and earthquakes to days in advance. In order to examine your proposal further it is requested that you may kindly furnish the following details to this office:


(i) The scientific principles on which your instrument functions and the type of data obtained through it.

(ii) Method of analysis of data and the inference drawn from it to forecast cyclones, earthquakes and heavy rain claimed by you.

(iii) Specific samples of forecast on cyclones, earthquakes and heavy rain you claim to provide 18 days in advance.

(iv) Verification procedure with specific instances.

(v) *Scientific* Specification publication, if any, on your instrument. (Give detailed reference)

Yours faithfully,  
  
 (M.C. PANT) 17/10/91  
 Director  
 for Director General of Meteorology.

-37-

सं०  
भारत सरकार  
भारत मौसम विज्ञान विभाग  
मौसम विज्ञान के महानिदेशक का कार्यालय  
मौसम भवन, लोदी रोड  
नई दिल्ली-११०००३  
सार का प्रता :  
महामौसम, नई दिल्ली



NO. NA-153  
GOVERNMENT OF INDIA  
INDIA METEOROLOGICAL DEPARTMENT  
OFFICE OF THE  
DIRECTOR GENERAL OF METEOROLOGY  
MAUSAM BHAVAN, LODI ROAD,  
NEW DELHI-110003  
Telegraphic Address:  
DIRGENMET, NEW DELHI

दिनांक/Date..Nov.....1996

To

Shri Gangadhar Rao Irlapati,  
C/o K. Chiranjeevi,  
H.No. 28-3, Saibabanagar,  
Judimetta,  
Hyderabad.


Subject:- Request for forwarding the copies of  
representation to President of India and other  
VVIP.

Sir,

Kindly refer to your letter dated September 12, 1996  
addressed to the Secretary, Lok Sabha Secretariat, Parliament  
House, New Delhi on the subject quoted above.

In this connection, you are requested to kindly refer our  
earlier letters of even number dated 8.6.95 and 8.1.96 in which  
you were advised suitably for your weather prediction device and  
recruitment in the Central Government establishment as well.  
You may proceed accordingly in your future action.

Yours faithfully,

  
(S.C. GOYAL)  
Director

for Director General of Meteorology



अर्जा श्रीकांत, प्राई.आर.टी.एम.  
**ARJA SRI KANTH, IRTS**  
 Tel.: 23387260  
 Fax: 23389025

90-  
 2008/2008/175/1000  
 निजी सचिव  
 खात राज्य मंत्री  
 भारत सरकार  
 शास्त्री भवन, नई दिल्ली-110 001  
 PRIVATE SECRETARY TO  
 MINISTER OF STATE FOR MINES  
 GOVERNMENT OF INDIA  
 SHASTRI BHAWAN, NEW DELHI 110 001

24 March 2008

Dear Sh. Ajit Tyagi Ji

Dr.T.Subbarami Reddy, Hon'ble Union Minister of State for Mines directed me to forward a representation received from Sh. I Gangadhara Rao, Hyderabad requesting for considering his proposal of Indian Weather Time Scale. The merits of the proposal may be examined.

A line of action taken may be communicated to apprise Hon'ble Union Minister.

With regards,

Yours sincerely,

  
 (Arja Srikanth)

AVM Ajit Tyagi  
 Director General of Meteorology,  
 India Meteorological Department,  
 Mausam Bhavan, Lodi Road,  
 New Delhi  
 Fax:011-24699216

Copy to Sh.I.Gangadhara Rao, Asst Section Officer, AP Public Service Commission, Nampally, Hyderabad 500055.

- 53 -

No. F-12016/1/00-NA/100

भारत सरकार  
भारत मौसम विज्ञान विभाग  
मौसम विज्ञान के महाविदेशक का कार्यालय  
मौसम भवन, लोदी रोड, नई दिल्ली-110003  
तार का पता: महामौसम, नई दिल्ली  
दूरभाष: 24611068, 24631913



GOVERNMENT OF INDIA  
INDIA METEOROLOGICAL DEPARTMENT  
OFFICE OF THE  
DIRECTOR GENERAL OF METEOROLOGY  
MAUSAM BHAWAN, LODI ROAD, NEW DELHI-110003  
Telegraphic Address: DIRGENMET, NEW DELHI  
Tel. No. 24611068/ 24631913, Fax No. 24643128,

November, 2009.

1. December

✓  
Shri Gangadhara Rao Irapati  
A.S.O., A.P.P.S.C., Nampally,  
Beside Gandhi Bhawan,  
Hyderabad - 500 001, A.P.

Subject:- "Indian Weather Time Scale" - regarding.

Sir,

With reference to your letter addressed to Secretary, Ministry of Earth Sciences, regarding forecast relating to prediction of cyclone, monsoon, heavy rainfall etc., you may kindly refer this office letter No. O-49106/537 dated 25/26.7.2005.

However, your dedication and interest in the field of meteorology is highly appreciated.

Thanking you,

Yours faithfully,

*T. Kumar*  
1.12.09  
(Awadhesh Kumar)  
Scientist 'E'

for Director General of Meteorology

33

1/25

सं०  
भारत सरकार  
भारत मौसम विज्ञान विभाग  
मौसम विज्ञान के महानिदेशक का कार्यालय  
मौसम भवन, लोदी रोड,  
नई दिल्ली-110003  
सार का पता :  
महामौसम, नई दिल्ली



NO. 49106/537  
GOVERNMENT OF INDIA  
INDIA METEOROLOGICAL DEPARTMENT  
OFFICE OF THE  
DIRECTOR GENERAL OF METEOROLOGY  
MAUSAM BHAVAN, LODI ROAD  
NEW DELHI-110003  
Telegraphic Address :  
DIRGENMET, NEW DELHI

दिनांक/Date... 25/07/2005

To:

Shri Gangadhara Rao Irlapati,  
H.No.5-30-4/1,  
Saibaba Nagar,  
Jeedimetla,  
Hyderabad.  
Andhra Pradesh  
Pin.Code No. 500 055.

Sub:- Project proposal to forecast drought, monsoon and rainfall etc.

Sir,

Kindly refer to your letter, regarding the project proposal for forecast the droughts, monsoon positions and rainfall etc. with the help of scale of data. You are requested to submit the project to Deptt. of Science and Technology (DST) through proper channel for necessary action.

M. Satya Kumar  
(M. Satya Kumar)  
Director Aviation Service  
For Director General of Meteorology

✓

371

  
 भारत सरकार  
 GOVERNMENT OF INDIA

भारत मौसम विज्ञान विभाग  
 INDIA METEOROLOGICAL DEPARTMENT

टेलिफोन : 25535220, 25535223, 25535254  
 TELEPHONE : 25535211, 25535245  
 टेलिग्रा : 145 7792 OBSR IN (Electronic)  
 TELEX : 0145 7227 MPNA IN

FAX : 091 020 25533201  
 कार : पून उल्लेख, पुणे  
 TELEGRAM : Weather, Pune

E-mail : [edgmrpune@hotmail.com](mailto:edgmrpune@hotmail.com)  
 मौसम विज्ञान के अपरमहानिदेशक (अनुसंधान)  
 शिवाजीनगर, पुणे - 411 005  
 Additional Director General of Meteorology (Research)  
 Shivajinagar, Pune - 411 005

No. ....  
 Date .....

GT-021(MISC)/6675  
 Dt. 28.08.2008.  
 13th

TO,  
 Shri.I.Gangadhara Rao  
 Asst.Section Officer,  
 A.P.Public Service Commission,  
 Beside Gandhi Bhavan,  
 Nampally,Hyderabad-500055,  
 Andhra Pradesh.


Sub: Project Proposal, "Indian Weather Time Scale" requested for establishment at Met.Centre,  
 Hyderabad.  
 Ref: Your letter dated Nil

Sir,

Kindly refer to your letter on the subject cited above .

Your project proposal has been examined by this office and it has been found that the  
 proposal "Indian Weather Time Scale" is without adequate scientific details/ reason. Therefore,  
 this office is unable to evaluate your project.

Thanking you.

  
 (Dr. T.P. Singh)  
 Meteorologist.Gr.I  
 For Additional Director General of Meteorology (Research)  
 Shivajinagar,Pune-5

161

GOVERNMENT OF ANDHRA PRADESH  
REVENUE (DM.III) DEPARTMENT

Letter No.25241/DM.III(3)/2009

dated:08.07.2009

From  
Sri.G.Ravi Babu, IAS.,  
Addl. Commissioner for Disaster Management &  
E.O. Dy. Secretary to Government,  
Revenue (DM) Department,  
A.P. Secretariat,  
**HYDERABAD – 500 022.**

To  
Sri. Gangadhara Rao Irlapati,  
H.No.5-30-4/1, Saibaba Nagar,  
Jeedimetla, Hyderabad – 500 055.

Sir,

Sub:- Project proposal – Establishment of “Andhra Pradesh State  
Weather Time Scale” – Regarding.

Ref:- From Sri.I Gangadhar Rao, Saibaba Nagar, Jeedimetla,  
Hyderabad letter dated 11.06.2009.

\*\*\*\*

With reference to your letter cited, you are requested to attend personally in the chambers of Addl. Commissioner for Disaster Management, Revenue (DM) Dept., A.P. Secretariat, Hyderabad on 13.07.2009 at 4.00 p.m. to explain the function of the “Andhra Pradesh State Weather Time Scale” by which the monsoon movements and its weather problems and natural calamities such as heavy rains, floods, droughts, cyclones etc., can be estimated on the Screen of the scale in advance etc.,

Yours faithfully,

*M. J. Srinivasan*  
for Addl. Commissioner for Disaster Management &  
E.O. Dy. Secretary to Government

20

IN THE GRAM PANCHAYAT OF THE MERLAPALEM VILLAGES  
 CERTIFYING DECISION P. R. NO. 87  
 ON THE 13th DAY OF DECEMBER, 1988  
 PARTICULARS OF GANGADHARA RAO IRLAPATI

This is to certify that the particulars of Gangadhara Rao Irlapati which are given below:-

**FAMILY PARTICULARS**

Name: Gangadhara Rao  
 Sir name: Irlapati  
 Father's Name: Pullayya  
 Place of Birth: Merlapalem  
 Date of Birth: 25th May, 1958

**NATIVITY PARTICULARS**

Nativity of village: Merlapalem  
 Mandal: Atreyasapuram  
 District: East Godavari  
 State: Andhra Pradesh

**COMMUNITY PARTICULARS**

Caste: Scheduled Caste  
 Sub-Caste: Mala  
 Religion: Hindu  
 Nationality: Indian  
 Social Position: Poor  
 Social conduct: Good patriot

The above particulars are true and correct as per the enquiry, verification and written witness of senior adults of the Gram Panchayat.



Utho Desam Pradhani  
 20/12/88 (Signature)

**ACADEMICAL PARTICULARS**

Scientific Qualification: None, Natural Genius  
 General Education  
 Elementary School Study: 1 to 5 classes  
 Upper Primary School study: 6 to 7 classes  
 High School Study: 8 to 10 classes  
 Pre-University course: Intermediate  
 Graduation: B.A. (Arts)  
 Post-Graduation:  
 Technical: P.T. (Trysem)

**RESEARCH EXPERIENCE PARTICULARS**

Year of starting of researches: 1963  
 Year of continuing of researches: 1988  
 Name of the research: Theory of Univariate (1977)  
 Place of the research: Scientific studies  
 Results of research: He has awarded his life  
 Total Period of his service: He has worked to the country for 25 years


**PRESENT SITUATION PARTICULARS**

Occupation: Un-employed  
 Wealth: Poverty  
 Health: Illness

Signature: *(Signature)*  
 Designation: GRAM PANCHAYAT  
 MERLAPALEM

20/12/88

4



47

72 -

No. DST/SECY/2009  
भारत सरकार

विज्ञान और प्रौद्योगिकी मंत्रालय  
विज्ञान और प्रौद्योगिकी विभाग  
टेक्नोलॉजी भवन, नया महरौली मार्ग, नई दिल्ली-110 016

GOVERNMENT OF INDIA  
MINISTRY OF SCIENCE & TECHNOLOGY  
DEPARTMENT OF SCIENCE & TECHNOLOGY  
Technology Bhavan, New Mehrauli Road, New Delhi-110 016

डा.टी.रामसामी  
सचिव  
Dr. T. RAMASAMI  
SECRETARY

June 1, 2009

Dear Shri Irlapati Rao,

I receive your letter of 11<sup>th</sup> May, 2009. Thank you. You may be aware that IITM is currently under the administrative control of Ministry of Earth Sciences. However, I have written to the Director, IITM requesting him to do the feasible in consultation with their Secretary.

Kindest regards,

Yours sincerely,  
*T. Ramasami*  
(T. Ramasami)

Shri Gangadhara Rao Irlapati  
Asst. Section Officer  
A.P. Public Service Commission  
(Beside Gandhi Bhavan)  
Nampally, Hyderabad 500 001

Tel. : 0091-11-26510068 / 26511439 • Fax : 0091-11-26863847 / 26862418 • E-mail : dstsec@nic.in

48  
160  
FROM  
M.G.GOPAL, I.A.S.,  
SECRETARY.



TO  
THE COMMISSIONER FOR DISASTI  
MANAGEMENT, AND EX.OFFICIO  
PRINCIPAL SECRETARY TO  
GOVERNMENT,  
REVENUE (DM.III) DEPARTMENT,  
ANDHRA PRADESH,  
HYDERABAD.

LETTER NO:869/ADB/4/2009, DT:15.07.2009.

Sir,

Sub:- A.P.P.S.C. - Egtt., - Forwarding the A.P.  
State Wather time scale prepared by  
Sri I.Gangadhar Rao, A.S.O., A.P.P.S.C.,  
Hyderabad - Regarding.


Ref:- Representation of Sri I.Gangadhar Rao,  
along with A.P. Weather time scale.

\* \* \*

I am directed to forward herewith the representation  
of Sri I.Gangadhar Rao, Assistant Section Officer, O/o Andhra  
Pradesh Public Service Commission, Hyderabad along with his  
reported research work on Andhra Pradesh State Weather Report  
for your consideration and necessary action.

Yours faithfully,

  
SECRETARY



5/2/2025